

Product datasheet

Anti-DDDDK tag (Binds to FLAG® tag sequence) antibody ab95045

★★★★★ [1 Abreviews](#) [2 References](#)

Overview

Product name	Anti-DDDDK tag (Binds to FLAG® tag sequence) antibody
Description	Goat polyclonal to DDDDK tag (Binds to FLAG® tag sequence)
Host species	Goat
Tested applications	Suitable for: ELISA, WB, ICC, IHC-P
Species reactivity	Reacts with: Species independent
Immunogen	Synthetic peptide corresponding to DDDDK tag (Binds to FLAG® tag sequence) conjugated to keyhole limpet haemocyanin.

General notes

FLAG® is a registered trade mark of Sigma Aldrich Biotechnology LP. It is used here for informational purposes only.

The Life Science industry has been in the grips of a reproducibility crisis for a number of years. Abcam is leading the way in addressing this with our range of recombinant monoclonal antibodies and knockout edited cell lines for gold-standard validation. Please check that this product meets your needs before purchasing.

If you have any questions, special requirements or concerns, please send us an inquiry and/or contact our Support team ahead of purchase. Recommended alternatives for this product can be found below, along with publications, customer reviews and Q&As

Properties

Form	Liquid
Storage instructions	Shipped at 4°C. Store at +4°C short term (1-2 weeks). Store at -20°C or -80°C. Avoid freeze / thaw cycle.
Storage buffer	pH: 7.20 Preservative: 0.1% Sodium azide Constituent: PBS
Purity	Protein G purified
Clonality	Polyclonal
Isotype	IgG

Applications

The Abpromise guarantee Our **Abpromise guarantee** covers the use of ab95045 in the following tested applications. The application notes include recommended starting dilutions; optimal dilutions/concentrations should be determined by the end user.

Application	Abreviews	Notes
ELISA		Use a concentration of 0.03 - 10 µg/ml. Coating: 2-10 µg/ml; Primary: 0.03-1 µg/ml
WB		Use a concentration of 0.03 - 1 µg/ml. Predicted molecular weight: 1 kDa. Colorimetric: 0.1-1 µg/ml; Chemiluminescent: 0.03-1 µg/ml
ICC		Use a concentration of 0.5 - 5 µg/ml.
IHC-P		Use at an assay dependent dilution.

Target

Relevance This is a useful tool for the localisation and characterisation of DDDDK tagged proteins (Binds to FLAG® tag sequence).

Please note: All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <https://www.abcam.com/abpromise> or contact our technical team.

Terms and conditions

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors