abcam

Product datasheet

Anti-delta 1 Catenin/CAS antibody [SP63] - BSA and Azide free ab236441

Recombinant RabMAb

14 Images

Overview

Product name Anti-delta 1 Catenin/CAS antibody [SP63] - BSA and Azide free

Description Rabbit monoclonal [SP63] to delta 1 Catenin/CAS - BSA and Azide free

Host species Rabbit

Tested applications Suitable for: IHC-P, Flow Cyt (Intra), ICC/IF

Species reactivity Reacts with: Human

Predicted to work with: Mouse, Rat, Chicken, Cow, Pig

Immunogen Synthetic peptide within Human delta 1 Catenin/CAS aa 850-950 (C terminal). The exact

> sequence is proprietary. Database link: O60716

Positive control IHC-P: Human breast carcinoma, breast ductal cell carcinoma, bladder transitional cell

carcinoma, cervical squamous cell carcinoma, cervix, colon adenocarcinoma, endometrium

adenocarcinoma, hepatocellular carcinoma, stomach adenocarcinoma, prostate

adenocarcinoma and esophagus tissues. ICC/IF: A431 cells. Flow Cyt (intra): A431 cells.

General notes ab236441 is the carrier-free version of ab227638.

> Our carrier-free antibodies are typically supplied in a PBS-only formulation, purified and free of BSA, sodium azide and glycerol. The carrier-free buffer and high concentration allow for

increased conjugation efficiency.

This conjugation-ready format is designed for use with fluorochromes, metal isotopes, oligonucleotides, and enzymes, which makes them ideal for antibody labelling, functional and cellbased assays, flow-based assays (e.g. mass cytometry) and Multiplex Imaging applications.

Use our conjugation kits for antibody conjugates that are ready-to-use in as little as 20 minutes with <1 minute hands-on-time and 100% antibody recovery: available for fluorescent dyes, HRP, biotin and gold.

This product is compatible with the Maxpar® Antibody Labeling Kit from Fluidigm, without the need for antibody preparation. Maxpar $^{\mbox{\scriptsize (B)}}$ is a trademark of Fluidigm Canada Inc.

This product is a recombinant monoclonal antibody, which offers several advantages including:

- High batch-to-batch consistency and reproducibility
- Improved sensitivity and specificity
- Long-term security of supply

- Animal-free production

For more information see here.

This product is FOR RESEARCH USE ONLY. For commercial use, please contact partnerships@abcam.com.

Properties

Form Liquid

Storage instructions Shipped at 4°C. Store at +4°C. Do Not Freeze.

Storage buffer pH: 7.20

Constituent: PBS

Carrier free Yes

Purity Protein A/G purified

Purification notes Purified from TCS by protein A/G.

Clonality Monoclonal

Clone number SP63

Isotype ΙgG

Applications

The Abpromise guarantee

Our Abpromise guarantee covers the use of ab236441 in the following tested applications.

The application notes include recommended starting dilutions; optimal dilutions/concentrations should be determined by the end user.

Application	Abreviews	Notes
IHC-P		Use at an assay dependent concentration. Perform heat mediated antigen retrieval before commencing with IHC staining protocol. Perform heat mediated antigen retrieval with EDTA buffer pH 6.0 before commencing with IHC staining protocol. Primary antibody incubation for 30 minutes at room temperature.
Flow Cyt (Intra)		Use at an assay dependent concentration.
ICC/IF		Use at an assay dependent concentration.

Target

Function Binds to and inhibits the transcriptional repressor ZBTB33, which may lead to activation of target

> genes of the Wnt signaling pathway (By similarity). May associate with and regulate the cell adhesion properties of both C- and E-cadherins. Implicated both in cell transformation by SRC and in ligand-induced receptor signaling through the EGF, PDGF, CSF-1 and ERBB2 receptors.

Promotes GLIS2 C-terminal cleavage.

Belongs to the beta-catenin family.

Tissue specificity Expressed in vascular endothelium. Sequence similarities

Contains 10 ARM repeats.

A possible nuclear localization signal exists in all isoforms where Asp-626--631-Arg are deleted.

Phosphorylated by protein-tyrosine kinases. Dephosphorylated by PTPRJ.

Cytoplasm. Nucleus. Cell membrane. Interaction with GLIS2 promotes nuclear translocation (By similarity). NANOS1 induces its translocation from sites of cell-cell contact to the cytoplasm.


Images

Domain

Post-translational

Cellular localization


modifications



Immunocytochemistry/ Immunofluorescence - Antidelta 1 Catenin/CAS antibody [SP63] - BSA and Azide free (ab236441)


Immunocytochemistry/ Immunofluorescence analysis of A431 (human epidermoid carcinoma epithelial cell) cells labeling delta 1 Catenin/CAS with purified ab227638. Cells were fixed in 100% Methanol. Cells were counterstained with ab195889 Anti-alpha Tubulin antibody [DM1A] - Microtubule Marker (Alexa Fluor® 594). Goat anti rabbit lgG (Alexa Fluor® 488, ab150077) was used as the secondary antibody. DAPI was used as nuclear counterstain. PBS instead of the primary antibody was used as the secondary antibody only control.

This data was developed using the same antibody clone in a different buffer formulation containing PBS, BSA and sodium azide (ab227638).



Immunohistochemistry (Formalin/PFA-fixed paraffinembedded sections) - Anti-delta 1 Catenin/CAS antibody [SP63] - BSA and Azide free (ab236441)


Formalin-fixed, paraffin-embedded human breast carcinoma tissue stained for delta 1 Catenin/CAS using ab227638 at 1/100 dilution in immunohistochemical analysis.



Flow Cytometry (Intracellular) - Anti-delta 1 Catenin/CAS antibody [SP63] - BSA and Azide free (ab236441)


Intracellular Flow Cytometry analysis of A431 (human epidermoid carcinoma epithelial cell) labeling delta 1 Catenin/CAS with purified ab227638 at (red). Cells were fixed with 4% paraformaldehyde and permeabilised with 90% methanol. Goat anti rabbit lgG (Alexa Fluor[®]488, ab150077) was used as a secondary antibody. Isotype control -Rabbit monoclonal lgG (ab172730) / Black. Unlableled control -Unlabelled cells / blue.

This data was developed using the same antibody clone in a different buffer formulation containing PBS, BSA and sodium azide (ab227638).



Immunohistochemistry (Formalin/PFA-fixed paraffinembedded sections) - Anti-delta 1 Catenin/CAS antibody [SP63] - BSA and Azide free (ab236441)


Formalin-fixed, paraffin-embedded human breast ductal cell carcinoma tissue stained for delta 1 Catenin/CAS using **ab227638** at 1/100 dilution in immunohistochemical analysis.



Immunohistochemistry (Formalin/PFA-fixed paraffinembedded sections) - Anti-delta 1 Catenin/CAS antibody [SP63] - BSA and Azide free (ab236441)


Formalin-fixed, paraffin-embedded human bladder transitional cell carcinoma tissue stained for delta 1 Catenin/CAS using **ab227638** at 1/100 dilution in immunohistochemical analysis.

This data was developed using the same antibody clone in a different buffer formulation containing PBS, BSA, glycerol, and sodium azide (ab227638).



Immunohistochemistry (Formalin/PFA-fixed paraffinembedded sections) - Anti-delta 1 Catenin/CAS antibody [SP63] - BSA and Azide free (ab236441)


Formalin-fixed, paraffin-embedded human cervical squamous cell carcinoma tissue stained for delta 1 Catenin/CAS using <u>ab227638</u> at 1/100 dilution in immunohistochemical analysis.



Immunohistochemistry (Formalin/PFA-fixed paraffinembedded sections) - Anti-delta 1 Catenin/CAS antibody [SP63] - BSA and Azide free (ab236441)


Formalin-fixed, paraffin-embedded human cervix tissue stained for delta 1 Catenin/CAS using <u>ab227638</u> at 1/100 dilution in immunohistochemical analysis.

This data was developed using the same antibody clone in a different buffer formulation containing PBS, BSA, glycerol, and sodium azide (ab227638).



Immunohistochemistry (Formalin/PFA-fixed paraffinembedded sections) - Anti-delta 1 Catenin/CAS antibody [SP63] - BSA and Azide free (ab236441)


Formalin-fixed, paraffin-embedded human colon adenocarcinoma tissue stained for delta 1 Catenin/CAS using <u>ab227638</u> at 1/100 dilution in immunohistochemical analysis.



Immunohistochemistry (Formalin/PFA-fixed paraffinembedded sections) - Anti-delta 1 Catenin/CAS antibody [SP63] - BSA and Azide free (ab236441)


Formalin-fixed, paraffin-embedded human endometrium adenocarcinoma tissue stained for delta 1 Catenin/CAS using **ab227638** at 1/100 dilution in immunohistochemical analysis.

This data was developed using the same antibody clone in a different buffer formulation containing PBS, BSA, glycerol, and sodium azide (ab227638).



Immunohistochemistry (Formalin/PFA-fixed paraffinembedded sections) - Anti-delta 1 Catenin/CAS antibody [SP63] - BSA and Azide free (ab236441)


Formalin-fixed, paraffin-embedded human hepatocellular carcinoma tissue stained for delta 1 Catenin/CAS using <u>ab227638</u> at 1/100 dilution in immunohistochemical analysis.



Immunohistochemistry (Formalin/PFA-fixed paraffinembedded sections) - Anti-delta 1 Catenin/CAS antibody [SP63] - BSA and Azide free (ab236441)


Formalin-fixed, paraffin-embedded human stomach adenocarcinoma tissue stained for delta 1 Catenin/CAS using **ab227638** at 1/100 dilution in immunohistochemical analysis.

This data was developed using the same antibody clone in a different buffer formulation containing PBS, BSA, glycerol, and sodium azide (ab227638).



Immunohistochemistry (Formalin/PFA-fixed paraffinembedded sections) - Anti-delta 1 Catenin/CAS antibody [SP63] - BSA and Azide free (ab236441)


Formalin-fixed, paraffin-embedded human esophagus tissue stained for delta 1 Catenin/CAS using <u>ab227638</u> at 1/100 dilution in immunohistochemical analysis.



Immunohistochemistry (Formalin/PFA-fixed paraffinembedded sections) - Anti-delta 1 Catenin/CAS antibody [SP63] - BSA and Azide free (ab236441)

Formalin-fixed, paraffin-embedded human prostate adenocarcinoma tissue stained for delta 1 Catenin/CAS using **ab227638** at 1/100 dilution in immunohistochemical analysis.

This data was developed using the same antibody clone in a different buffer formulation containing PBS, BSA, glycerol, and sodium azide (ab227638).



Anti-delta 1 Catenin/CAS antibody [SP63] - BSA and Azide free (ab236441)

Please note: All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- · We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit https://www.abcam.com/abpromise or contact our technical team.

Terms and conditions

• Guarantee only valid for products bought direct from Abcam or one of our authorized distributors