


## Product datasheet

# Anti-E1 Ubiquitin Activating Enzyme 1/UBA1 antibody [EPR14203(B)] - BSA and Azide free ab250138

Recombinant RabMAb

[5 Images](#)

### Overview

<b>Product name</b>	Anti-E1 Ubiquitin Activating Enzyme 1/UBA1 antibody [EPR14203(B)] - BSA and Azide free
<b>Description</b>	Rabbit monoclonal [EPR14203(B)] to E1 Ubiquitin Activating Enzyme 1/UBA1 - BSA and Azide free
<b>Host species</b>	Rabbit
<b>Tested applications</b>	<b>Suitable for:</b> Flow Cyt (Intra), IHC-P, ICC/IF, WB
<b>Species reactivity</b>	<b>Reacts with:</b> Human <b>Predicted to work with:</b> Mouse, Rat 
<b>Immunogen</b>	Synthetic peptide. This information is proprietary to Abcam and/or its suppliers.
<b>General notes</b>	<p>ab250138 is the carrier-free version of <a href="#">ab180125</a>.</p> <p>Our <b>carrier-free</b> antibodies are typically supplied in a PBS-only formulation, purified and free of BSA, sodium azide and glycerol. The carrier-free buffer and high concentration allow for increased conjugation efficiency.</p> <p>This conjugation-ready format is designed for use with fluorochromes, metal isotopes, oligonucleotides, and enzymes, which makes them ideal for antibody labelling, functional and cell-based assays, flow-based assays (e.g. mass cytometry) and Multiplex Imaging applications.</p> <p>Use our <b>conjugation kits</b> for antibody conjugates that are ready-to-use in as little as 20 minutes with &lt;1 minute hands-on-time and 100% antibody recovery: available for fluorescent dyes, HRP, biotin and gold.</p> <p>This product is compatible with the Maxpar<sup>®</sup> Antibody Labeling Kit from Fluidigm, without the need for antibody preparation. Maxpar<sup>®</sup> is a trademark of Fluidigm Canada Inc.</p>

### Properties

<b>Form</b>	Liquid
<b>Storage instructions</b>	Shipped at 4°C. Store at +4°C. Do Not Freeze.
<b>Storage buffer</b>	pH: 7.2 Constituent: PBS

<b>Carrier free</b>	Yes
<b>Purity</b>	Protein A purified
<b>Clonality</b>	Monoclonal
<b>Clone number</b>	EPR14203(B)
<b>Isotype</b>	IgG

## Applications


**The Abpromise guarantee** Our **Abpromise guarantee** covers the use of ab250138 in the following tested applications. The application notes include recommended starting dilutions; optimal dilutions/concentrations should be determined by the end user.

Application	Abreviews	Notes
<b>Flow Cyt (Intra)</b>		Use at an assay dependent concentration.
<b>IHC-P</b>		Use at an assay dependent concentration. Perform heat mediated antigen retrieval with Tris/EDTA buffer pH 9.0 before commencing with IHC staining protocol.
<b>ICC/IF</b>		Use at an assay dependent concentration.
<b>WB</b>		Use at an assay dependent concentration. Detects a band of approximately 117 kDa (predicted molecular weight: 118 kDa).

## Target

<b>Function</b>	Activates ubiquitin by first adenylating its C-terminal glycine residue with ATP, and thereafter linking this residue to the side chain of a cysteine residue in E1, yielding an ubiquitin-E1 thioester and free AMP.
<b>Pathway</b>	Protein modification; protein ubiquitination.
<b>Involvement in disease</b>	Defects in UBA1 are the cause of spinal muscular atrophy X-linked type 2 (SMAX2) [MIM:301830]; also known as X-linked lethal infantile spinal muscular atrophy, distal X-linked arthrogryposis multiplex congenita or X-linked arthrogryposis type 1 (AMCX1). Spinal muscular atrophy refers to a group of neuromuscular disorders characterized by degeneration of the anterior horn cells of the spinal cord, leading to symmetrical muscle weakness and atrophy. SMAX2 is a lethal infantile form presenting with hypotonia, areflexia, and multiple congenital contractures.
<b>Sequence similarities</b>	Belongs to the ubiquitin-activating E1 family.
<b>Post-translational modifications</b>	ISGylated.

## Images



Western blot - Anti-E1 Ubiquitin Activating Enzyme 1/UBA1 antibody [EPR14203(B)] - BSA and Azide free (ab250138)

**All lanes** : Anti-E1 Ubiquitin Activating Enzyme 1/UBA1 antibody [EPR14203(B)] (**ab180125**) at 1/5000 dilution

**Lane 1** : Fetal brain tissue lysate

**Lane 2** : K562 cell lysate

**Lane 3** : HeLa cell lysate

**Lane 4** : Jurkat cell lysate

Lysates/proteins at 20 µg per lane.


**Secondary**

**All lanes** : Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/1000 dilution

**Predicted band size:** 118 kDa

**Observed band size:** 117 kDa


This data was developed using **ab180125**, the same antibody clone in a different buffer formulation.



Flow Cytometry (Intracellular) - Anti-E1 Ubiquitin Activating Enzyme 1/UBA1 antibody [EPR14203(B)] - BSA and Azide free (ab250138)


This data was developed using **ab180125**, the same antibody clone in a different buffer formulation.

Intracellular flow cytometric analysis of 2% paraformaldehyde-fixed K562 cells labeling E1 Ubiquitin Activating Enzyme with **ab180125** at 1/70 dilution (red) compared to a rabbit IgG control (green), followed by Goat anti rabbit IgG (FITC) secondary antibody at 1/75 dilution.



Immunocytochemistry/ Immunofluorescence - Anti-E1 Ubiquitin Activating Enzyme 1/UBA1 antibody [EPR14203(B)] - BSA and Azide free (ab250138)

This data was developed using **ab180125**, the same antibody clone in a different buffer formulation. Immunofluorescent analysis of acetone-fixed Jurkat cells labeling E1 Ubiquitin Activating Enzyme with **ab180125** at 1/100 dilution, followed by Goat anti rabbit IgG (Dylight 488) at 1/250 dilution. Counter stained with DAPI (blue).



Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-E1 Ubiquitin Activating Enzyme 1/UBA1 antibody [EPR14203(B)] - BSA and Azide free (ab250138)

This data was developed using **ab180125**, the same antibody clone in a different buffer formulation. Immunohistochemical analysis of paraffin-embedded Human ovarian carcinoma tissue labeling E1 Ubiquitin Activating Enzyme with **ab180125** at 1/250 dilution, followed by prediluted HRP Polymer for Rabbit IgG. Counter stained with Hematoxylin. Perform heat mediated antigen retrieval with EDTA buffer pH 9 before commencing with IHC staining protocol.

### Why choose a recombinant antibody?



**Research with confidence**  
Consistent and reproducible results



**Long-term and scalable supply**  
Recombinant technology



**Success from the first experiment**  
Confirmed specificity



**Ethical standards compliant**  
Animal-free production

Anti-E1 Ubiquitin Activating Enzyme 1/UBA1  
antibody [EPR14203(B)] - BSA and Azide free  
(ab250138)

**Please note:** All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

### Our Abpromise to you: Quality guaranteed and expert technical support

---

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <https://www.abcam.com/abpromise> or contact our technical team.

### Terms and conditions

---

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors