

Product datasheet

Anti-ErbB2 / HER2 antibody [ICR12] ab11710

Recombinant

[3 References](#) [5 Images](#)

Overview

Product name	Anti-ErbB2 / HER2 antibody [ICR12]
Description	Rat monoclonal [ICR12] to ErbB2 / HER2
Host species	Rat
Tested applications	Suitable for: IHC-P, ICC/IF, Flow Cyt Unsuitable for: IP or WB
Species reactivity	Reacts with: Human
Immunogen	Tissue, cells or virus. This information is proprietary to Abcam and/or its suppliers.
Positive control	IHC-P: Human breast carcinoma tissue. ICC/IF: SK-BR-3 cells. Flow Cyt: SK-BR-3 cells.
General notes	<p>This antibody clone is manufactured by Abcam. If you require a custom buffer formulation or conjugation for your experiments, please contact orders@abcam.com.</p> <p>This product is a recombinant monoclonal antibody, which offers several advantages including:</p> <ul style="list-style-type: none">- High batch-to-batch consistency and reproducibility- Improved sensitivity and specificity- Long-term security of supply- Animal-free production <p>For more information see here.</p>

Properties

Form	Liquid
Storage instructions	Shipped at 4°C. Store at +4°C short term (1-2 weeks). Upon delivery aliquot. Store at -20°C long term. Avoid freeze / thaw cycle.
Storage buffer	pH: 7.2 Preservative: 0.01% Sodium azide Constituents: PBS, 0.05% BSA, 40% Glycerol (glycerin, glycerine)
Purity	Ion Exchange Chromatography
Clonality	Monoclonal
Clone number	ICR12
Isotype	IgG2a

Light chain type kappa

Applications

The Abpromise guarantee Our **Abpromise guarantee** covers the use of ab11710 in the following tested applications. The application notes include recommended starting dilutions; optimal dilutions/concentrations should be determined by the end user.

Application	Abreviews	Notes
IHC-P		1/400. Perform heat mediated antigen retrieval with Tris/EDTA buffer pH 9.0 before commencing with IHC staining protocol.
ICC/IF		1/100.
Flow Cyt		1/1000.

Application notes Is unsuitable for IP or WB.

Target

Function Protein tyrosine kinase that is part of several cell surface receptor complexes, but that apparently needs a coreceptor for ligand binding. Essential component of a neuregulin-receptor complex, although neuregulins do not interact with it alone. GP30 is a potential ligand for this receptor. Regulates outgrowth and stabilization of peripheral microtubules (MTs). Upon ERBB2 activation, the MEMO1-RHOA-DIAPH1 signaling pathway elicits the phosphorylation and thus the inhibition of GSK3B at cell membrane. This prevents the phosphorylation of APC and CLASP2, allowing its association with the cell membrane. In turn, membrane-bound APC allows the localization of MACF1 to the cell membrane, which is required for microtubule capture and stabilization. In the nucleus is involved in transcriptional regulation. Associates with the 5'-TCAAATTC-3' sequence in the PTGS2/COX-2 promoter and activates its transcription. Implicated in transcriptional activation of CDKN1A; the function involves STAT3 and SRC. Involved in the transcription of rRNA genes by RNA Pol I and enhances protein synthesis and cell growth.

Tissue specificity Expressed in a variety of tumor tissues including primary breast tumors and tumors from small bowel, esophagus, kidney and mouth.

Involvement in disease Hereditary diffuse gastric cancer
Glioma
Ovarian cancer
Lung cancer
Gastric cancer
Chromosomal aberrations involving ERBB2 may be a cause gastric cancer. Deletions within 17q12 region producing fusion transcripts with CDK12, leading to CDK12-ERBB2 fusion leading to truncated CDK12 protein not in-frame with ERBB2.


Sequence similarities Belongs to the protein kinase superfamily. Tyr protein kinase family. EGF receptor subfamily. Contains 1 protein kinase domain.

Post-translational modifications Autophosphorylated. Autophosphorylation occurs in trans, i.e. one subunit of the dimeric receptor phosphorylates tyrosine residues on the other subunit (Probable). Ligand-binding increases phosphorylation on tyrosine residues (PubMed:27134172). Signaling via SEMA4C promotes phosphorylation at Tyr-1248 (PubMed:17554007). Dephosphorylated by PTPN12 (PubMed:27134172).

Cellular localization

Cytoplasm. Nucleus and Cell membrane. Cytoplasm, perinuclear region. Nucleus. Translocation to the nucleus requires endocytosis, probably endosomal sorting and is mediated by importin beta-1/KPNB1.

Images




Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-ErbB2 / HER2 antibody [ICR12] (ab11710)

Immunohistochemical analysis of paraffin-embedded human breast carcinoma tissue labeling ErbB2 / HER2 with ab11710 at 1/400 dilution followed by ready to use Goat Anti-rat IgG H&L (HRP polymer) ([ab214882](#)). Membranous staining on human breast carcinoma. Counterstained with Hematoxylin.

Secondary antibody only control: Secondary antibody is ready to use Goat Anti-rat IgG H&L (HRP polymer) ([ab214882](#)).

Heat mediated antigen retrieval using [ab93684](#) (Tris/EDTA buffer, pH 9.0).



Immunocytochemistry/ Immunofluorescence - Anti-ErbB2 / HER2 antibody [ICR12] (ab11710)


Immunofluorescent analysis of 4% Paraformaldehyde-fixed, 0.1% Triton X-100 permeabilized SK-BR-3 cells labelling ErbB 2 with ab11710 at 1/100 dilution, followed by [ab150157](#) Goat Anti-rat IgG H&L (Alexa Fluor[®] 488) antibody at 1/1000 dilution (Green).

Confocal image showing strong membranous staining in SK-BR-3 cells. [ab179513](#) Anti-beta Tubulin rabbit monoclonal antibody was used to counterstain tubulin at 1/200 dilution, followed by [ab150080](#) Goat Anti-Rabbit IgG H&L (Alexa Fluor[®] 594) at a 1/1000 dilution (Red). The nuclear counterstain was DAPI (Blue).

Negative control 1: ab11710 at 1/100 dilution followed by [ab150080](#) at a 1/1000 dilution.

Negative control 2: [ab179513](#) at a 1/200 dilution followed by [ab150157](#) at a 1/1000 dilution.

Negative control cells: MCF7 (PMID: 18288420).




Flow Cytometry - Anti-ErbB2 / HER2 antibody
[ICR12] (ab11710)

Flow cytometric analysis of MCF7 (human breast adenocarcinoma epithelial cell, Left panel) / SK-BR-3 (human breast adenocarcinoma epithelial cell, Right panel) cells labelling ErbB 2 with ab11710 at 1/1000 dilution (0.1µg) (Red) compared with a rat monoclonal IgG (Black) isotype control and an unlabelled control (cells without incubation with primary antibody and secondary antibody) (Blue). Goat anti rat IgG (Alexa Fluor® 488, **ab150157**) at 1/2000 dilution was used as the secondary antibody.

Low expression control: MCF7 (PMID: 17938260).

Gated on viable cells.



Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-ErbB2 / HER2 antibody
[ICR12] (ab11710)

Immunohistochemical analysis of paraffin-embedded human breast carcinoma tissue labeling ErbB 2 with ab11710 at 1/400 dilution followed by ready to use Goat Anti-rat IgG H&L (HRP polymer) (**ab214882**). No staining on human breast carcinoma without expression of HER2 is observed. Counterstained with Hematoxylin.

Secondary antibody only control: Secondary antibody is ready to use Goat Anti-rat IgG H&L (HRP polymer) (**ab214882**).

Heat mediated antigen retrieval using **ab93684** (Tris/EDTA buffer, pH 9.0).

Why choose a recombinant antibody?



Research with confidence
Consistent and reproducible results



Long-term and scalable supply
Recombinant technology



Success from the first experiment
Confirmed specificity



Ethical standards compliant
Animal-free production

Anti-ErbB2 / HER2 antibody [ICR12] (ab11710)

Please note: All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <https://www.abcam.com/abpromise> or contact our technical team.

Terms and conditions

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors