

Product datasheet

Anti-ErbB4 / HER4 (phospho Y1188) antibody [EPR2271Y] ab68479

Recombinant RabMAb

[1 References](#) [2 Images](#)

Overview

Product name	Anti-ErbB4 / HER4 (phospho Y1188) antibody [EPR2271Y]
Description	Rabbit monoclonal [EPR2271Y] to ErbB4 / HER4 (phospho Y1188)
Host species	Rabbit
Specificity	This antibody detects ErbB 4 phosphorylated on tyrosine 1188.
Tested applications	Suitable for: WB Unsuitable for: Flow Cyt, IHC-P or IP
Species reactivity	Reacts with: Human
Immunogen	Synthetic peptide. This information is proprietary to Abcam and/or its suppliers.
Positive control	A431 cell lysate, cells treated with EGF.
General notes	<p>This product is a recombinant monoclonal antibody, which offers several advantages including:</p> <ul style="list-style-type: none"> - High batch-to-batch consistency and reproducibility - Improved sensitivity and specificity - Long-term security of supply - Animal-free production <p>For more information see here.</p> <p>Our RabMAb[®] technology is a patented hybridoma-based technology for making rabbit monoclonal antibodies. For details on our patents, please refer to RabMAb[®] patents.</p> <p>Mouse, Rat: We have preliminary internal testing data to indicate this antibody may not react with these species. Please contact us for more information.</p>

Properties

Form	Liquid
Storage instructions	Shipped at 4°C. Upon delivery aliquot and store at -20°C. Avoid freeze / thaw cycles.
Storage buffer	<p>pH: 7.2</p> <p>Preservative: 0.05% Sodium azide</p> <p>Constituents: 0.1% BSA, 40% Glycerol (glycerin, glycerine), 9.85% Tris glycine, 50% Tissue culture supernatant</p>

Purity	Protein A purified
Clonality	Monoclonal
Clone number	EPR2271Y
Isotype	IgG

Applications

The Abpromise guarantee Our **Abpromise guarantee** covers the use of ab68479 in the following tested applications.

The application notes include recommended starting dilutions; optimal dilutions/concentrations should be determined by the end user.

Application	Abreviews	Notes
WB		1/1000 - 1/5000. Detects a band of approximately 170-200 kDa (predicted molecular weight: 147 kDa).

Application notes Is unsuitable for Flow Cyt, IHC-P or IP.

Target

Function Specifically binds and is activated by neuregulins, NRG-2, NRG-3, heparin-binding EGF-like growth factor, betacellulin and NTAK. Interaction with these factors induces cell differentiation. Not activated by EGF, TGF- α , and amphiregulin. The C-terminal fragment (CTF) of isoform JMA-A CYT-2 (containing E4ICD2) can stimulate transcription in the presence of YAP1. ERBB4 intracellular domain is involved in the regulation of cell growth. Conflicting reports are likely due at least in part to the opposing effects of the isoform-specific and nuclear-translocated ERBB4 intracellular domains (E4ICD1 and E4ICD2). Overexpression studies in epithelium show growth inhibition using E4ICD1 and increased proliferation using E4ICD2. E4ICD2 has greater in vitro kinase activity than E4ICD1. The kinase activity is required for the nuclear translocation of E4ICD2.

Tissue specificity Expressed at highest levels in brain, heart, kidney, in addition to skeletal muscle, parathyroid, cerebellum, pituitary, spleen, testis and breast. Lower levels in thymus, lung, salivary gland, and pancreas. Isoform JM-A CYT-1 and isoform JM-B CYT-1 are expressed in cerebellum, but only the isoform JM-B is expressed in the heart.

Sequence similarities Belongs to the protein kinase superfamily. Tyr protein kinase family. EGF receptor subfamily. Contains 1 protein kinase domain.


Post-translational modifications Isoform JM-A CYT-1 and isoform JM-A CYT-2 but not isoform JM-B CYT-1 and isoform JM-B CYT-2 are processed by ADAM17. Proteolytic processing in response to ligand or 12-O-tetradecanoylphorbol-13-acetate stimulation results in the production of 120 kDa soluble receptor forms and intermediate membrane-anchored 80 kDa fragments (m80HER4), which are further processed by a presenilin-dependent gamma-secretase to release the respective cytoplasmic intracellular domain E4ICD (either E4ICD1/s80Cyt1 or E4ICD2/s80Cyt2). Membrane-anchored 80 kDa fragments of the processed isoform JM-A CYT-1 are more readily degraded by the proteasome than fragments of isoform JM-A CYT-2 suggesting a prevalence of E4ICD2 over E4ICD1. Ligand-binding increases phosphorylation on tyrosine residues. Isoform JM-A CYT-2 is constitutively phosphorylated on tyrosine residues in a ligand-independent manner. E4ICD2 but not E4ICD1 is phosphorylated on tyrosine residues. Ubiquitinated. The ERBB4 intracellular domain is ubiquitinated and targeted to proteosomal

degradation during mitosis mediated by the APC/C complex. Isoform JM-A CYT-1 and isoform JM-B CYT-1 are ubiquitinated by WWP1. The ERBB4 intracellular domain (E4ICD1) is ubiquitinated, and this involves NEDD4.

Cellular localization

Membrane and Nucleus. Following proteolytical processing E4ICD (E4ICD1 or E4ICD2 generated from the respective isoforms) is translocated to the nucleus. Significantly more E4ICD2 than E4ICD1 is found in the nucleus. E4ICD2 colocalizes with YAP1 in the nucleus.

Images



All lanes : Anti-ErbB4 / HER4 (phospho Y1188) antibody [EPR2271Y] (ab68479) at 1/2000 dilution

Lane 1 : A431 cell lysate. Cells untreated.

Lane 2 : A431 cell lysate. Cells treated with EGF.

Lysates/proteins at 10 µg per lane.

Secondary

All lanes : HRP conjugated Goat anti-Rabbit at 1/2000 dilution

Predicted band size: 147 kDa

Observed band size: 170-200 kDa

Why choose a recombinant antibody?

Research with confidence
Consistent and reproducible results

Long-term and scalable supply
Recombinant technology

Success from the first experiment
Confirmed specificity

Ethical standards compliant
Animal-free production

Anti-ErbB4 / HER4 (phospho Y1188) antibody [EPR2271Y] (ab68479)

Please note: All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <https://www.abcam.com/abpromise> or contact our technical team.

Terms and conditions

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors