abcam

Product datasheet

Anti-Fibrinogen beta chain antibody [EPR3083] - BSA and Azide free ab248494

Recombinant

RabMAb

3 Images

Overview

Product name Anti-Fibrinogen beta chain antibody [EPR3083] - BSA and Azide free

Description Rabbit monoclonal [EPR3083] to Fibrinogen beta chain - BSA and Azide free

Host species Rabbit

Tested applications Suitable for: ICC/IF, WB

Unsuitable for: IHC-P or IP

Species reactivity Reacts with: Rat, Human

Predicted to work with: Mouse

Immunogen Synthetic peptide. This information is proprietary to Abcam and/or its suppliers.

General notes ab248494 is the carrier-free version of <u>ab133359</u>.

Our <u>carrier-free</u> antibodies are typically supplied in a PBS-only formulation, purified and free of BSA, sodium azide and glycerol. The carrier-free buffer and high concentration allow for increased conjugation efficiency.

This conjugation-ready format is designed for use with fluorochromes, metal isotopes, oligonucleotides, and enzymes, which makes them ideal for antibody labelling, functional and cell-based assays, flow-based assays (e.g. mass cytometry) and Multiplex Imaging applications.

Use our **conjugation kits** for antibody conjugates that are ready-to-use in as little as 20 minutes with <1 minute hands-on-time and 100% antibody recovery: available for fluorescent dyes, HRP, biotin and gold.

This product is compatible with the Maxpar[®] Antibody Labeling Kit from Fluidigm, without the need for antibody preparation. Maxpar[®] is a trademark of Fluidigm Canada Inc.

This product is a recombinant monoclonal antibody, which offers several advantages including:

- High batch-to-batch consistency and reproducibility
- Improved sensitivity and specificity
- Long-term security of supply
- Animal-free production

For more information see here.

Our RabMAb[®] technology is a patented hybridoma-based technology for making rabbit monoclonal antibodies. For details on our patents, please refer to **RabMAb**[®] **patents**.

1

Properties

Form Liquid

Storage instructions Shipped at 4°C. Store at +4°C. Do Not Freeze.

Storage buffer pH: 7.2

Constituent: PBS

Carrier free Yes

Purity Affinity purified

Clonality Monoclonal

Clone number EPR3083

Isotype IgG

Applications

The Abpromise guarantee Our Abpromise guarantee covers the use of ab248494 in the following tested applications.

The application notes include recommended starting dilutions; optimal dilutions/concentrations should be determined by the end user.

Application	Abreviews	Notes
ICC/IF		Use at an assay dependent concentration.
WB		Use at an assay dependent concentration. Detects a band of approximately 56 kDa (predicted molecular weight: 56 kDa).

Application notes Is unsuitable for IHC-P or IP.

Target

Function Fibrinogen has a double function: yielding monomers that polymerize into fibrin and acting as a

cofactor in platelet aggregation.

Involvement in disease Defects in FGB are a cause of congenital afibrinogenemia (CAFBN) [MIM:202400]. This rare

autosomal recessive disorder is characterized by bleeding that varies from mild to severe and by complete absence or extremely low levels of plasma and platelet fibrinogen. Note=Patients with congenital fibrinogen abnormalities can manifest different clinical pictures. Some cases are clinically silent, some show a tendency toward bleeding and some show a predisposition for

thrombosis with or without bleeding.

Sequence similarities Contains 1 fibrinogen C-terminal domain.

Domain A long coiled coil structure formed by 3 polypeptide chains connects the central nodule to the C-

terminal domains (distal nodules). The long C-terminal ends of the alpha chains fold back,

contributing a fourth strand to the coiled coil structure.

Post-translational

modifications from alpha and beta chains, and thus exposes the N-terminal polymerization sites responsible for


the formation of the soft clot. The soft clot is converted into the hard clot by factor XIIIA which catalyzes the epsilon-(gamma-glutamyl)lysine cross-linking between gamma chains (stronger)

Conversion of fibrinogen to fibrin is triggered by thrombin, which cleaves fibrinopeptides A and B

and between alpha chains (weaker) of different monomers.

Cellular localization Secreted.

Images



Western blot - Anti-Fibrinogen beta chain antibody [EPR3083] - BSA and Azide free (ab248494)

All lanes : Anti-Fibrinogen beta chain antibody [EPR3083] (ab133359) at 1/1000 dilution

Lane 1 : Human plasma lysate
Lane 3 : Rat platelet lysate

Lysates/proteins at 10 µg per lane.


Secondary

All lanes: HRP labelled goat anti-rabbit at 1/2000 dilution

Predicted band size: 56 kDa


This data was developed using <u>ab133359</u>, the same antibody clone in a different buffer formulation.

Actual band size: 56 kDa.



Immunocytochemistry/ Immunofluorescence - Anti-Fibrinogen beta chain antibody [EPR3083] - BSA and Azide free (ab248494)

This data was developed using <u>ab133359</u>, the same antibody clone in a different buffer formulation.lmmunofluorescence analysis of Fibrinogen beta chain in HeLa cells, using <u>ab133359</u> at a dilution of 1/100.



Please note: All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- · We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit https://www.abcam.com/abpromise or contact our technical team.

Terms and conditions

• Guarantee only valid for products bought direct from Abcam or one of our authorized distributors