

## Product datasheet

# Anti-Fibronectin antibody [EPR23110-46] - BSA and Azide free ab268022

KO VALIDATED Recombinant RabMAb

★★★★★ [3 Abreviews](#) [11 Images](#)

### Overview

Product name	Anti-Fibronectin antibody [EPR23110-46] - BSA and Azide free
Description	Rabbit monoclonal [EPR23110-46] to Fibronectin - BSA and Azide free
Host species	Rabbit
Tested applications	<b>Suitable for:</b> IHC-P, IHC-Fr, ICC/IF, WB <b>Unsuitable for:</b> Flow Cyt or IP
Species reactivity	<b>Reacts with:</b> Mouse, Rat, Human
Immunogen	Synthetic peptide. This information is proprietary to Abcam and/or its suppliers.
Positive control	WB: Human serum, kidney and stomach lysate. Mouse, human and rat plasma lysate. NIH/3T3 whole cell lysate; HAP1 whole cell lysate. IHC-P: Human mammary gland tissue. Human esophageal squamous cell carcinoma tissue. Mouse and rat stomach tissue, human pancreatic carcinoma tissue. ICC/IF: HepG2 and NIH/3T3 cells.
General notes	<p>ab268022 is the carrier-free version of <a href="#">ab268020</a>.</p> <p>Our <b>carrier-free</b> antibodies are typically supplied in a PBS-only formulation, purified and free of BSA, sodium azide and glycerol. The carrier-free buffer and high concentration allow for increased conjugation efficiency.</p> <p>This conjugation-ready format is designed for use with fluorochromes, metal isotopes, oligonucleotides, and enzymes, which makes them ideal for antibody labelling, functional and cell-based assays, flow-based assays (e.g. mass cytometry) and Multiplex Imaging applications.</p> <p>Use our <b>conjugation kits</b> for antibody conjugates that are ready-to-use in as little as 20 minutes with &lt;1 minute hands-on-time and 100% antibody recovery: available for fluorescent dyes, HRP, biotin and gold.</p> <p>This product is compatible with the Maxpar® Antibody Labeling Kit from Fluidigm, without the need for antibody preparation. Maxpar® is a trademark of Fluidigm Canada Inc.</p> <p>This product is a recombinant monoclonal antibody, which offers several advantages including:</p> <ul style="list-style-type: none"> <li>- High batch-to-batch consistency and reproducibility</li> <li>- Improved sensitivity and specificity</li> <li>- Long-term security of supply</li> <li>- Animal-free production</li> </ul> <p>For more information <a href="#">see here</a>.</p>

Our RabMAb<sup>®</sup> technology is a patented hybridoma-based technology for making rabbit monoclonal antibodies. For details on our patents, please refer to [RabMAb<sup>®</sup> patents](#).

## Properties

Form	Liquid
Storage instructions	Shipped at 4°C. Store at +4°C. Do Not Freeze.
Storage buffer	pH: 7.2 Constituent: PBS
Carrier free	Yes
Purity	Protein A purified
Clonality	Monoclonal
Clone number	EPR23110-46
Isotype	IgG

## Applications

**The Abpromise guarantee** Our **Abpromise guarantee** covers the use of ab268022 in the following tested applications. The application notes include recommended starting dilutions; optimal dilutions/concentrations should be determined by the end user.

Application	Abreviews	Notes
IHC-P	★★★★★ (1)	Use at an assay dependent concentration. Perform heat mediated antigen retrieval with Tris/EDTA buffer pH 9.0 before commencing with IHC staining protocol.
IHC-Fr	★★★★★ (1)	Use at an assay dependent concentration. Heat mediated antigen retrieval using sodium citrate buffer (10mM citrate pH 6.0 + 0.05% Tween-20).
ICC/IF		Use at an assay dependent concentration.
WB		Use at an assay dependent concentration. Detects a band of approximately 285 kDa (predicted molecular weight: 262 kDa).

**Application notes** Is unsuitable for Flow Cyt or IP.

## Target

**Function** Fibronectins bind cell surfaces and various compounds including collagen, fibrin, heparin, DNA, and actin. Fibronectins are involved in cell adhesion, cell motility, opsonization, wound healing, and maintenance of cell shape. Involved in osteoblast compaction through the fibronectin fibrillogenesis cell-mediated matrix assembly process, essential for osteoblast mineralization. Participates in the regulation of type I collagen deposition by osteoblasts. Anastellin binds fibronectin and induces fibril formation. This fibronectin polymer, named superfibronectin, exhibits enhanced adhesive properties. Both anastellin and superfibronectin inhibit tumor growth, angiogenesis and metastasis. Anastellin activates p38 MAPK and inhibits

lysophospholipid signaling.

### Tissue specificity

Plasma FN (soluble dimeric form) is secreted by hepatocytes. Cellular FN (dimeric or cross-linked multimeric forms), made by fibroblasts, epithelial and other cell types, is deposited as fibrils in the extracellular matrix. Ugl-Y1, Ugl-Y2 and Ugl-Y3 are found in urine.

### Involvement in disease

Glomerulopathy with fibronectin deposits 2

### Sequence similarities

Contains 12 fibronectin type-I domains.

Contains 2 fibronectin type-II domains.

Contains 16 fibronectin type-III domains.

### Developmental stage

Ugl-Y1, Ugl-Y2 and Ugl-Y3 are present in the urine from 0 to 17 years of age.

### Post-translational modifications

Sulfated.

It is not known whether both or only one of Thr-2064 and Thr-2065 are/is glycosylated.

Forms covalent cross-links mediated by a transglutaminase, such as F13A or TGM2, between a glutamine and the epsilon-amino group of a lysine residue, forming homopolymers and heteropolymers (e.g. fibrinogen-fibronectin, collagen-fibronectin heteropolymers).


Phosphorylated by FAM20C in the extracellular medium.

Proteolytic processing produces the C-terminal NC1 peptide, anastellin.

### Cellular localization

Secreted, extracellular space, extracellular matrix.

## Images




Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-Fibronectin antibody [EPR23110-46] - BSA and Azide free (ab268022)

Immunohistochemical analysis of paraffin-embedded Human pancreatic carcinoma tissue labeling Fibronectin with [ab268020](#) at 1/4000 dilution (0.146 µg/ml) followed by a ready to use Goat Anti-Rabbit IgG H&L (HRP). The section was incubated with [ab268020](#) at 4°C overnight. Positive staining in the human pancreatic carcinoma counterstained with hematoxylin.

Secondary antibody only control: Secondary antibody is a ready to use Goat Anti-Rabbit IgG H&L (HRP)([ab214880](#))

Heat mediated antigen retrieval using [ab93684](#) (Tris/EDTA buffer, pH 9.0).

This data was developed using the same antibody clone in a different buffer formulation containing PBS, BSA, glycerol, and sodium azide ([ab268020](#)).




Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-Fibronectin antibody [EPR23110-46] - BSA and Azide free (ab268022)

Immunohistochemical analysis of paraffin-embedded rat stomach tissue labeling Fibronectin with **ab268020** at 1/2000 dilution (0.29 µg/ml) followed by a ready to use Goat Anti-Rabbit IgG H&L (HRP). Positive staining in the smooth muscle of rat stomach (PMID:650101) Counterstained with hematoxylin.

Secondary antibody only control: Secondary antibody is a ready to use Goat Anti-Rabbit IgG H&L (HRP).

Heat mediated antigen retrieval using **ab93684** (Tris/EDTA buffer, pH 9.0).

This data was developed using the same antibody clone in a different buffer formulation containing PBS, BSA, glycerol, and sodium azide (**ab268020**).



Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-Fibronectin antibody [EPR23110-46] - BSA and Azide free (ab268022)


Immunohistochemical analysis of paraffin-embedded mouse stomach tissue labeling Fibronectin with **ab268020** at 1/2000 dilution (0.29 µg/ml) followed by a ready to use Goat Anti-Rabbit IgG H&L (HRP). Positive staining in the smooth muscle of mouse stomach (PMID:650101) Counterstained with hematoxylin.

Secondary antibody only control: Secondary antibody is a ready to use Goat Anti-Rabbit IgG H&L (HRP).

Heat mediated antigen retrieval using **ab93684** (Tris/EDTA buffer, pH 9.0).

This data was developed using the same antibody clone in a different buffer formulation containing PBS, BSA, glycerol, and sodium azide (**ab268020**).






Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-Fibronectin antibody [EPR23110-46] - BSA and Azide free (ab268022)

Immunohistochemical analysis of paraffin-embedded human esophageal squamous cell carcinoma tissue labeling Fibronectin with **ab268020** at 1/2000 dilution (0.29 µg/ml) followed by a ready to use Goat Anti-Rabbit IgG H&L (HRP). Positive staining in the stroma of human esophageal squamous cell carcinoma (PMID:30314454). Counterstained with hematoxylin.

Secondary antibody only control: Secondary antibody is a ready to use Goat Anti-Rabbit IgG H&L (HRP).

Heat mediated antigen retrieval using **ab93684** (Tris/EDTA buffer, pH 9.0).

This data was developed using the same antibody clone in a different buffer formulation containing PBS, BSA, glycerol, and sodium azide (**ab268020**).




Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-Fibronectin antibody [EPR23110-46] - BSA and Azide free (ab268022)

Immunohistochemical analysis of paraffin-embedded human mammary gland tissue labeling Fibronectin with **ab268020** at 1/2000 dilution (0.29 µg/ml) followed by a ready to use Goat Anti-Rabbit IgG H&L (HRP). Positive staining in the stroma of human mammary gland (PMID:650101). Counterstained with hematoxylin.

Secondary antibody only control: Secondary antibody is a ready to use Goat Anti-Rabbit IgG H&L (HRP).

Heat mediated antigen retrieval using **ab93684** (Tris/EDTA buffer, pH 9.0).

This data was developed using the same antibody clone in a different buffer formulation containing PBS, BSA, glycerol, and sodium azide (**ab268020**).




Immunohistochemistry (Frozen sections) - Anti-Fibronectin antibody [EPR23110-46] - BSA and Azide free (ab268022)

Immunohistochemical analysis of frozen section of 4% PFA-fixed, 0.2% Triton X-100 permeabilized rat colon tissue labeling Fibronectin with [ab268020](#) at 1/100 dilution followed by [ab150077](#) AlexaFluor®488 Goat anti-Rabbit secondary at 1/1000 dilution (Green). Positive staining mainly on smooth muscle of rat colon (PMID: 17881565) is observed. The nuclear counterstain is DAPI (Blue).

Secondary antibody only control: Secondary antibody is [ab150077](#) AlexaFluor®488 Goat anti-Rabbit secondary at 1/1000 dilution

Heat mediated antigen retrieval using sodium citrate buffer (10mM citrate pH 6.0 + 0.05% Tween-20).

This data was developed using the same antibody clone in a different buffer formulation containing PBS, BSA, glycerol, and sodium azide ([ab268020](#)).




Immunohistochemistry (Frozen sections) - Anti-Fibronectin antibody [EPR23110-46] - BSA and Azide free (ab268022)

Immunohistochemical analysis of frozen section of 4% PFA-fixed, 0.2% Triton X-100 permeabilized mouse colon tissue labeling Fibronectin with [ab268020](#) at 1/100 dilution followed by [ab150077](#) AlexaFluor®488 Goat anti-Rabbit secondary at 1/1000 dilution (Green). Positive staining mainly on smooth muscle of mouse colon (PMID: 17881565) is observed. The nuclear counterstain is DAPI (Blue).

Secondary antibody only control: Secondary antibody is [ab150077](#) AlexaFluor®488 Goat anti-Rabbit secondary at 1/1000 dilution

Heat mediated antigen retrieval using sodium citrate buffer (10mM citrate pH 6.0 + 0.05% Tween-20).

This data was developed using the same antibody clone in a different buffer formulation containing PBS, BSA, glycerol, and sodium azide ([ab268020](#)).



**All lanes :** Anti-Fibronectin antibody [EPR23110-46] (**ab268020**) at 0.585 µg/ml (1:1000 dilution)

**Lane 1 :** Human stomach tissue lysate at 10 µg

**Lane 2 :** NIH/3T3 (mouse embryonic fibroblast), whole cell lysate at 10 µg

**Lane 3 :** Fibronectin-1 knockout HAP1 whole cell lysate at 40 µg

**Lane 4 :** Wild-type HAP1 whole cell lysate at 40 µg

Developed using the ECL technique.

**Predicted band size:** 262 kDa


**Observed band size:** 285 kDa

This data was developed using the same antibody clone in a different buffer formulation containing PBS, BSA, glycerol, and sodium azide (**ab268020**).

**ab268020** was shown to specifically react with Fibronectin in wild-type HAP1 cells as signal was lost in Fibronectin knockout cells. Wild-type and Fibronectin knockout samples were subjected to SDS-PAGE. **ab268020** and **ab129002** (Rabbit anti-Vinculin loading control) were incubated 1 hour at room temperature at 1/1000 dilution and 1/5000 dilution respectively. Blots were developed with Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated (**ab97051**) secondary antibody at 1/100,000 dilution for 1 hour at room temperature before imaging. The blot was developed on a BIO-RAD® ChemiDoc™ MP instrument using the ECL technique.

Exposure time: Lanes 1-2: 37 seconds; Lanes 3-4: 3 minutes.

Blocking/diluting buffer and concentration: 5% NFDm/TBST.




Immunocytochemistry/ Immunofluorescence - Anti-Fibronectin antibody [EPR23110-46] - BSA and Azide free (ab268022)

Immunofluorescent analysis of 4% Paraformaldehyde-fixed, 0.1% TritonX-100 permeabilized NIH/3T3 (mouse embryonic fibroblast) cells labeling Fibronectin with **ab268020** at 1/50 dilution, followed by **ab150077** AlexaFluor®488 Goat anti-Rabbit secondary antibody at 1/1000 dilution (Green). Confocal image showing cytoplasmic staining in NIH/3T3 cells is observed. **ab195889** Anti-alpha Tubulin antibody [DM1A] - Microtubule Marker (Alexa Fluor® 594) was used to counterstain tubulin at 1/200 dilution (Red). The nuclear counterstain was DAPI (Blue).

Secondary antibody only control: Secondary antibody is **ab150077** AlexaFluor®488 Goat anti-Rabbit secondary at 1/1000 dilution.

This data was developed using the same antibody clone in a different buffer formulation containing PBS, BSA, glycerol, and sodium azide (**ab268020**).



Immunocytochemistry/ Immunofluorescence - Anti-Fibronectin antibody [EPR23110-46] - BSA and Azide free (ab268022)

Immunofluorescent analysis of 4% Paraformaldehyde-fixed, 0.1% TritonX-100 permeabilized HepG2 (human hepatocellular carcinoma epithelial cell) cells labeling Fibronectin with **ab268020** at 1/50 dilution, followed by **ab150077** AlexaFluor®488 Goat anti-Rabbit secondary antibody at 1/1000 dilution (Green). Confocal image showing cytoplasmic staining in HepG2 cells is observed. **ab195889** Anti-alpha Tubulin antibody [DM1A] - Microtubule Marker (Alexa Fluor® 594) was used to counterstain tubulin at 1/200 dilution (Red). The nuclear counterstain was DAPI (Blue).

Secondary antibody only control: Secondary antibody is **ab150077** AlexaFluor®488 Goat anti-Rabbit secondary at 1/1000 dilution.

This data was developed using the same antibody clone in a different buffer formulation containing PBS, BSA, glycerol, and sodium azide (**ab268020**).



### Why choose a recombinant antibody?



**Research with confidence**  
Consistent and reproducible results



**Long-term and scalable supply**  
Recombinant technology



**Success from the first experiment**  
Confirmed specificity



**Ethical standards compliant**  
Animal-free production

Anti-Fibronectin antibody [EPR23110-46] - BSA and Azide free (ab268022)

**Please note:** All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

### Our Abpromise to you: Quality guaranteed and expert technical support

---

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <https://www.abcam.com/abpromise> or contact our technical team.

### Terms and conditions

---

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors