abcam

Product datasheet

Anti-Filaggrin antibody [rFLG/1561] - BSA and Azide free ab236431



2 Images

Overview

Product name Anti-Filaggrin antibody [rFLG/1561] - BSA and Azide free

Description Mouse monoclonal [rFLG/1561] to Filaggrin - BSA and Azide free

Host species Mouse

Tested applications
Suitable for: IHC-P
Species reactivity
Reacts with: Human

Immunogen Recombinant fragment within Human Filaggrin aa 198-288. The exact sequence is proprietary.

Database link: P20930

Positive control IHC-P: Human skin tissue.

Properties

Form Liquid

Storage instructions Shipped at 4°C. Store at +4°C short term (1-2 weeks). Upon delivery aliquot. Store at -20°C long

term. Avoid freeze / thaw cycle.

Storage buffer pH: 7.2

Constituent: PBS

Carrier free Yes

Purity Protein A/G purified

Purification notes Purified from bioreactor concentrate by Protein A/G.

ClonalityMonoclonalClone numberrFLG/1561IsotypeIgG1

Applications

The Abpromise guarantee Our Abpromise guarantee covers the use of ab236431 in the following tested applications.

The application notes include recommended starting dilutions; optimal dilutions/concentrations should be determined by the end user.

1

Application	Abreviews	Notes
IHC-P		Use a concentration of 0.5 - 1 µg/ml. Staining of formalin-fixed tissues is enhanced by boiling tissue sections in 10 mM citrate buffer, pH 6.0, for 10-20 minutes followed by cooling at RT for 20 minutes Primary incubation for 30 minutes at room temperature.

Target

Function Aggregates keratin intermediate filaments and promotes disulfide-bond formation among the

intermediate filaments during terminal differentiation of mammalian epidermis.

Tissue specificity Keratohyalin granules.

Involvement in disease Ichthyosis vulgaris

Dermatitis atopic 2

Sequence similaritiesBelongs to the S100-fused protein family.

Contains 2 EF-hand domains. Contains 23 filaggrin repeats.

Post-translational

modifications

Filaggrin is initially synthesized as a large, insoluble, highly phosphorylated precursor containing many tandem copies of 324 AA, which are not separated by large linker sequences. During terminal differentiation it is dephosphorylated and proteolytically cleaved. The N-terminal of the mature protein is heterogeneous, and is blocked by the formation of pyroglutamate.

Undergoes deimination of some arginine residues (citrullination).


Images



Immunohistochemistry (Formalin/PFA-fixed paraffinembedded sections) - Anti-Filaggrin antibody
[rFLG/1561] - BSA and Azide free (ab236431)

Formalin-fixed, paraffin-embedded human skin tissue stained for Filaggrin using $\underline{ab234414}$ at 1 $\mu g/ml$ in immunohistochemical analysis.

This data was developed using the same antibody clone in a different buffer formulation containing PBS, BSA and sodium azide (ab234414).



Please note: All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

Our Abpromise to you: Quality guaranteed and expert technical support

- · Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit https://www.abcam.com/abpromise or contact our technical team.

Terms and conditions

· Guarantee only valid for products bought direct from Abcam or one of our authorized distributors