

FITC Anti-Rotavirus antibody ab31435

3 References

Overview

Product name	FITC Anti-Rotavirus antibody
Description	FITC Goat polyclonal to Rotavirus
Host species	Goat
Conjugation	FITC. Ex: 493nm, Em: 528nm
Tested applications	Suitable for: ICC/IF
Species reactivity	Reacts with: Rotavirus
Immunogen	Tissue, cells or virus corresponding to Rotavirus.
General notes	<p>The Life Science industry has been in the grips of a reproducibility crisis for a number of years. Abcam is leading the way in addressing this with our range of recombinant monoclonal antibodies and knockout edited cell lines for gold-standard validation. Please check that this product meets your needs before purchasing.</p> <p>If you have any questions, special requirements or concerns, please send us an inquiry and/or contact our Support team ahead of purchase. Recommended alternatives for this product can be found below, along with publications, customer reviews and Q&As</p>

Properties

Form	Liquid
Storage instructions	Shipped at 4°C. Store at +4°C.
Storage buffer	<p>pH: 7.20</p> <p>Preservative: 0.1% Sodium azide</p> <p>Constituents: 1% BSA, PBS</p>
Purity	Ion Exchange Chromatography
Purification notes	Ammonium sulfate fractionation and ion-exchange chromatography.
Clonality	Polyclonal
Isotype	IgG

Applications

The **Abpromise guarantee** Our **Abpromise guarantee** covers the use of ab31435 in the following tested applications.

The application notes include recommended starting dilutions; optimal dilutions/concentrations should be determined by the end user.

Application	Abreviews	Notes
ICC/IF		1/250. Acetone fixed.

Target

Relevance

Rotaviruses, members of the family Reoviridae, are a major cause of diarrhoea in young mammals. Rotavirus infections also result in economic losses in agriculture due to diarrhoea in calf, pig, sheep, and poultry rearing. Diarrhoea (or scours) due to the rotavirus Nebraska Calf Diarrhea Virus can affect calves up to 30 days of age or older. Diarrhoea begins 2 to 3 days after exposure. Diagnosis is by history, lesions (ulcers on the tongue, lips, and mouth) and diagnostic laboratory tests. Mortality rates may be as high as 50 percent, depending on the secondary bacteria present. Human rotaviruses, the major aetiological agents of severe infantile diarrhoea worldwide, display surprisingly diverse and complex serotypic specificities. Rotaviruses are 70 nm, non enveloped viruses comprised of a triple layered protein capsid; Outer capsid proteins are VP4 and VP7, Inner capsid -VP6 and Core -VP2. The immunity acquired from exposure to rotavirus appears to be type specific following initial infection; therefore, multiple serotypes of rotavirus mean multiple opportunities for infection. The combination of animal reservoirs for the virus and rotavirus gene reassortment provides the potential for dramatic genetic shifts (similar to influenza virus) which could give rise to altered host ranges and viral virulence.

Please note: All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <https://www.abcam.com/abpromise> or contact our technical team.

Terms and conditions

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors