

## Product datasheet

# Anti-Forkhead box protein O/dFOXO antibody - N-terminal ab195977

## 6 References

### Overview

<b>Product name</b>	Anti-Forkhead box protein O/dFOXO antibody - N-terminal
<b>Description</b>	Rabbit polyclonal to Forkhead box protein O/dFOXO - N-terminal
<b>Host species</b>	Rabbit
<b>Tested applications</b>	<b>Suitable for:</b> WB
<b>Species reactivity</b>	<b>Reacts with:</b> Drosophila melanogaster
<b>Immunogen</b>	Synthetic peptide corresponding to Drosophila melanogaster Forkhead box protein O/dFOXO (N terminal). Database link: <a href="#">Q95V55</a>

### General notes

The Life Science industry has been in the grips of a reproducibility crisis for a number of years. Abcam is leading the way in addressing this with our range of recombinant monoclonal antibodies and knockout edited cell lines for gold-standard validation. Please check that this product meets your needs before purchasing.

If you have any questions, special requirements or concerns, please send us an inquiry and/or contact our Support team ahead of purchase. Recommended alternatives for this product can be found below, along with publications, customer reviews and Q&As

### Properties

<b>Form</b>	Lyophilized:Reconstitute with 100 uL distilled water.
<b>Storage instructions</b>	Shipped at 4°C. Store at +4°C short term (1-2 weeks). Upon delivery aliquot. Store at -20°C long term. Avoid freeze / thaw cycle.
<b>Storage buffer</b>	Constituents: 1.21% Tris buffer, 0.75% Glycine, 2% Sucrose
<b>Purity</b>	Protein A purified
<b>Clonality</b>	Polyclonal
<b>Isotype</b>	IgG

### Applications

**The Abpromise guarantee** Our **Abpromise guarantee** covers the use of ab195977 in the following tested applications.

The application notes include recommended starting dilutions; optimal dilutions/concentrations should be determined by the end user.

Application	Abreviews	Notes
WB		Use at an assay dependent concentration.

## Target

<b>Function</b>	Transcription factor involved in the regulation of the insulin signaling pathway. Consistently activates both the downstream target Thor'd4EBP and the feedback control target InR. Involved in negative regulation of the cell cycle, modulating cell growth and proliferation. In response to cellular stresses, such as nutrient deprivation or increased levels of reactive oxygen species, foxo is activated and inhibits growth through the action of target genes such as Thor. Foxo activated in the adult fat body can regulate lifespan in adults; an insulin peptide itself may function as one secondary messenger of insulin-regulated aging. Also regulates Lip4, homolog of human acid lipases, thereby acting as a key modulator of lipid metabolism by insulin signaling and integrates insulin responses to glucose and lipid homeostasis.
<b>Sequence similarities</b>	Contains 1 fork-head DNA-binding domain.
<b>Cellular localization</b>	Cytoplasm. Nucleus. When phosphorylated, translocated from nucleus to cytoplasm. Dephosphorylation triggers nuclear translocation.

**Please note:** All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

## Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <https://www.abcam.com/abpromise> or contact our technical team.

## Terms and conditions

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors