

Product datasheet

Anti-GFP antibody ab5449

★★★★★ [1 Abreviews](#) [14 References](#) [2 Images](#)

Overview

| | |
|----------------------------|---|
| Product name | Anti-GFP antibody |
| Description | Goat polyclonal to GFP |
| Host species | Goat |
| Specificity | Reactive against all variants of <i>Aequorea victoria</i> GFP such as S65T-GFP, RS-GFP, YFP and EGFP. |
| Tested applications | Suitable for: WB, IP |
| Species reactivity | Reacts with: Species independent |
| Immunogen | Recombinant full length protein. This information is proprietary to Abcam and/or its suppliers. |
| Positive control | Pure GFP protein, or cells known to overexpress GFP. |
| General notes | <p>Protein A will not bind goat IgG, so use alternates (eg. protein G) in IP with this antibody. This antibody is available in an affinity purified form as ab5450.</p> <p>The Life Science industry has been in the grips of a reproducibility crisis for a number of years. Abcam is leading the way in addressing this with our range of recombinant monoclonal antibodies and knockout edited cell lines for gold-standard validation. Please check that this product meets your needs before purchasing.</p> <p>If you have any questions, special requirements or concerns, please send us an inquiry and/or contact our Support team ahead of purchase. Recommended alternatives for this product can be found below, along with publications, customer reviews and Q&As</p> |

Properties

| | |
|-----------------------------|---|
| Form | Liquid |
| Storage instructions | Shipped at 4°C. Upon delivery aliquot and store at -20°C or -80°C. Avoid repeated freeze / thaw cycles. |
| Storage buffer | Preservative: 0.05% Sodium azide |
| Purity | Whole antiserum |
| Clonality | Polyclonal |
| Isotype | IgG |

Applications

The Abpromise guarantee

Our **Abpromise guarantee** covers the use of ab5449 in the following tested applications.

The application notes include recommended starting dilutions; optimal dilutions/concentrations should be determined by the end user.

| Application | Abreviews | Notes |
|-------------|-----------|--------------------------------------|
| WB | ★★★★★ (1) | 1/5000 - 1/20000. |
| IP | | Use 0.5µl for 10 ⁶ cells. |

Target

Relevance

Function: Energy-transfer acceptor. Its role is to transduce the blue chemiluminescence of the protein aequorin into green fluorescent light by energy transfer. Fluoresces in vivo upon receiving energy from the Ca²⁺-activated photoprotein aequorin.

Subunit structure: Monomer.

Tissue specificity: Photocytes.

Post-translational modification: Contains a chromophore consisting of modified amino acid residues. The chromophore is formed by autocatalytic backbone condensation between Ser-65 and Gly-67, and oxidation of Tyr-66 to didehydrotyrosine. Maturation of the chromophore requires nothing other than molecular oxygen.


Biotechnological use: Green fluorescent protein has been engineered to produce a vast number of variously colored mutants, fusion proteins, and biosensors. Fluorescent proteins and its mutated allelic forms, blue, cyan and yellow have become a useful and ubiquitous tool for making chimeric proteins, where they function as a fluorescent protein tag. Typically they tolerate N- and C-terminal fusion to a broad variety of proteins. They have been expressed in most known cell types and are used as a noninvasive fluorescent marker in living cells and organisms. They enable a wide range of applications where they have functioned as a cell lineage tracer, reporter of gene expression, or as a measure of protein-protein interactions. Can also be used as a molecular thermometer, allowing accurate temperature measurements in fluids. The measurement process relies on the detection of the blinking of GFP using fluorescence correlation spectroscopy.

Sequence similarities: Belongs to the GFP family.

Biophysicochemical properties: Absorption: Abs(max)=395 nm

Exhibits a smaller absorbance peak at 470 nm. The fluorescence emission spectrum peaks at 509 nm with a shoulder at 540 nm.

Images



Western blot - Anti-GFP antibody (ab5449)

Lane 1 : parental YTS cells (negative control)

Lanes 2-5 : YTS cells transfected with KIR-EGFP (mw 88 kD)

KIR-EGFP IP's with Goat polyclonal to GFP (ab5449) using 0.1 ul for 2×10^6 cells. KIR-EGFP detected with a mouse monoclonal to KIR receptor (Borszcz et al EGI 2003, 33: 1084).

Lane 1 : parental YTS cells (negative control) Lanes 2-5 : YTS cells transfected with KIR-EGFP (mw 88 kD) KIR-EGFP IP's with Goat polyclonal to GFP (ab5449) using 0.1 ul for 2×10^6 cells. KIR-EGFP detected with a mouse monoclonal to KIR receptor (Borszcz et al EGI 2003, 33: 1084).



Western blot - Anti-GFP antibody (ab5449)

5 ng GFP on PVDF membrane QC. Goat polyclonal to GFP (ab5449) used at dilutions of:

Lane 1 : 1/2500

Lane 2 : 1/5000

Lane 3 : 1/10,000

Lane 4 : 1/20,000

5 ng GFP on PVDF membrane QC. Goat polyclonal to GFP (ab5449) used at dilutions of: Lane 1 : 1/2500 Lane 2 : 1/5000 Lane 3 : 1/10,000 Lane 4 : 1/20,000

Please note: All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <https://www.abcam.com/abpromise> or contact our technical team.

Terms and conditions

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors