

Anti-Glucose Transporter GLUT1 antibody [EPR3915] ab115730

KO **VALIDATED** Recombinant RabMAB


★★★★☆ [19 Abreviews](#) [235 References](#) [26 Images](#)

Overview

Product name	Anti-Glucose Transporter GLUT1 antibody [EPR3915]
Description	Rabbit monoclonal [EPR3915] to Glucose Transporter GLUT1
Host species	Rabbit
Specificity	We recommend not to boil the samples after lysis to get desired WB bands.
Tested applications	Suitable for: Flow Cyt (Intra), ICC/IF, WB, IHC-P
Species reactivity	Reacts with: Mouse, Rat, Human
Immunogen	Synthetic peptide. This information is proprietary to Abcam and/or its suppliers. (Peptide available as ab202335)
Positive control	WB: NIH/3T3, HepG2, HT-29, SW480, 3T3-L1 and PC-12 whole cell lysates. IHC-P: Rat kidney tissue; mouse liver tissue; human lung carcinoma, cervical carcinoma, colon carcinoma, liver, colon, kidney carcinoma, skeletal muscle, urinary bladder, heart and breast tissue. ICC/IF: HepG2 cells and A549 (SLC2A1 knockout A549 cells used as a negative control) cells. Flow Cyt (intra): HeLa and Jurkat cells.
General notes	<p>This product is a recombinant monoclonal antibody, which offers several advantages including:</p> <ul style="list-style-type: none">- High batch-to-batch consistency and reproducibility- Improved sensitivity and specificity- Long-term security of supply- Animal-free production <p>For more information see here.</p> <p>Our RabMAB[®] technology is a patented hybridoma-based technology for making rabbit monoclonal antibodies. For details on our patents, please refer to RabMAB[®] patents.</p> <p>We are constantly working hard to ensure we provide our customers with best in class antibodies. As a result of this work we are pleased to now offer this antibody in purified format. We are in the process of updating our datasheets. The purified format is designated 'PUR' on our product labels. If you have any questions regarding this update, please contact our Scientific Support team.</p>

Properties

Form	Liquid
Storage instructions	Shipped at 4°C. Store at +4°C short term (1-2 weeks). Upon delivery aliquot. Store at -20°C. Avoid freeze / thaw cycle.
Dissociation constant (K_D)	K _D = 7.70 x 10 ⁻¹² M



Storage buffer	pH: 7.20 Preservative: 0.01% Sodium azide Constituents: 49% PBS, 50% Glycerol (glycerin, glycerine), 0.05% BSA
Purity	Protein A purified
Clonality	Monoclonal
Clone number	EPR3915
Isotype	IgG

Applications


The Abpromise guarantee Our **Abpromise guarantee** covers the use of ab115730 in the following tested applications. The application notes include recommended starting dilutions; optimal dilutions/concentrations should be determined by the end user.

Application	Abreviews	Notes
Flow Cyt (Intra)		1/40. For unpurified, use 1/100 - 1/1000. ab172730 - Rabbit monoclonal IgG, is suitable for use as an isotype control with this antibody.
ICC/IF	★★★★★ (2)	Use a concentration of 1 µg/ml. This product gave a positive signal in A549 (SLC2A1 knockout A549 cells used as a negative control) fixed with 100% methanol (5 min).
WB	★★★★☆ (11)	1/100000. Detects a band of approximately 40-60 kDa (predicted molecular weight: 54 kDa). We would not recommend boiling due to the possible irreversible aggregation of glucose transporters. If samples are boiled it can prevent some of the protein from entering the gel or produce multimers which are often mistaken for background. Samples should be solubilized in standard SDS Laemmli buffer and
IHC-P	★★★★☆ (4)	1/250 - 1/500. Perform heat mediated antigen retrieval before commencing with IHC staining protocol. See <u>IHC antigen retrieval protocols</u> .

Target

Function	Facilitative glucose transporter. This isoform may be responsible for constitutive or basal glucose uptake. Has a very broad substrate specificity; can transport a wide range of aldoses including both pentoses and hexoses.
Tissue specificity	Expressed at variable levels in many human tissues.
Involvement in disease	Defects in SLC2A1 are the cause of glucose transporter type 1 deficiency syndrome (GLUT1DS) [MIM:606777]; also known as blood-brain barrier glucose transport defect. This disease causes a defect in glucose transport across the blood-brain barrier. It is characterized by infantile seizures, delayed development, and acquired microcephaly. Defects in SLC2A1 are the cause of dystonia type 18 (DYT18) [MIM:612126]. DYT18 is an exercise-induced paroxysmal dystonia/dyskinesia. Dystonia is defined by the presence of sustained involuntary muscle contraction, often leading to abnormal postures. DYT18 is characterized by attacks of involuntary movements triggered by certain stimuli such as sudden movement or prolonged exercise. In some patients involuntary exertion-induced dystonic, choreoathetotic, and ballistic movements may be associated with macrocytic hemolytic anemia.
Sequence similarities	Belongs to the major facilitator superfamily. Sugar transporter (TC 2.A.1.1) family. Glucose transporter subfamily.
Post-translational modifications	Phosphorylated upon DNA damage, probably by ATM or ATR.
Cellular localization	Cell membrane. Melanosome. Localizes primarily at the cell surface (By similarity). Identified by mass spectrometry in melanosome fractions from stage I to stage IV.

Images



All lanes : Anti-Glucose Transporter GLUT1 antibody [EPR3915] (ab115730) at 1/100000 dilution

Lane 1 : Wild-type HepG2 cell lysate

Lane 2 : LC2A1 knockout HepG2 cell lysate

Lane 3 : A549 cell lysate

Lane 4 : Jurkat cell lysate

Lysates/proteins at 20 mg/ml per lane.


Performed under reducing conditions.

Predicted band size: 54 kDa

Observed band size: 50-300 kDa

Western blot: Anti-Glucose Transporter GLUT1 antibody [EPR3915] staining at 1/100000 dilution, shown in green; Mouse anti-GAPDH antibody [6C5] ([ab8245](#)) loading control staining at 1/20000 dilution, shown in red. In Western blot, ab115730 was shown to bind specifically to Glucose Transporter GLUT1. A band was observed

at 50-300 kDa in wild-type HepG2 cell lysates with no signal observed at this size in SLC2A1 knockout cell line [ab280797](#) (knockout cell lysate [ab284224](#)). To generate this image, wild-type and SLC2A1 knockout HepG2 cell lysates were analysed. First, samples were run on an SDS-PAGE gel then transferred onto a nitrocellulose membrane. Membranes were blocked in 3% milk in TBS-0.1% Tween[®] 20 (TBS-T) before incubation with primary antibodies overnight at 4°C. Blots were washed four times in TBS-T, incubated with secondary antibodies for 1 h at room temperature, washed again four times then imaged. Secondary antibodies used were Goat anti-Rabbit IgG H&L 800CW and Goat anti-Mouse IgG H&L 680RD at 1/20000 dilution.



Western blot - Anti-Glucose Transporter GLUT1 antibody [EPR3915] (ab115730)

All lanes : Anti-Glucose Transporter GLUT1 antibody [EPR3915] (ab115730) at 1 µg/ml

Lane 1 : Wild-type A549 whole cell lysate


Lane 2 : SLC2A1 knockout A549 whole cell lysate

Lysates/proteins at 20 µg per lane.

Predicted band size: 54 kDa

Lanes 1 - 2: Merged signal (red and green). Green - [ab196357](#) observed at 54 kDa. Red - loading control, [ab8245](#), observed at 37 kDa.

[ab196357](#) was shown to recognize in wild-type A549 cells as signal was lost at the expected MW in SLC2A1 knockout cells. Additional cross-reactive bands were observed in the wild-type and knockout cells. Wild-type and SLC2A1 knockout samples were subjected to SDS-PAGE. Ab196357 and [ab8245](#) (Mouse anti-GAPDH loading control) were incubated overnight at 4°C at 1 µg/ml and 1/20000 dilution respectively. Blots were developed with Goat anti-Rabbit IgG H&L (IRDye[®] 800CW) preabsorbed [ab216773](#) and Goat anti-Mouse IgG H&L (IRDye[®] 680RD) preabsorbed [ab216776](#) secondary antibodies at 1/20000 dilution for 1 hour at room temperature before imaging.




GLUT1 and GLUT3 are downregulated in KSHV-infected cells in human KS tumors

Representative illustration of dual immunofluorescence detection of LANA and GLUT1 or in a normal human skin section and a Kaposi Sarcoma (KS) tumor section. Tissues were fixed with paraformaldehyde and paraffin-embedded.

Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-Glucose Transporter

GLUT1 antibody [EPR3915] (ab115730)

Zhu, Y. et al PLoS Pathog. 2016 May 17;12(5):e1005648. doi: 10.1371/journal.ppat.1005648. eCollection 2016 May Reproduced under the Creative Commons license <http://creativecommons.org/licenses/by/4.0/>




Immunohistochemical expression of Glut1 in normal tongue epithelium and tongue cancer. Expression was greatest in lymphocytes (arrows in left upper and lower panels). In the normal oral epithelium, Glut1 was weakly expressed in the basal and spinous cells (left upper panel). In OSCC, Glut1 was upregulated, showing a level of expression comparable with lymphocytes (left and right lower panels). Scale bar, 100 μ m.

Note: Glut1 = SLC2A (alternative names for the same target).

Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-Glucose Transporter

GLUT1 antibody [EPR3915] (ab115730)

Khaom, R. et al PLoS One. 2016 Aug 11;11(8):e0161163. doi: 10.1371/journal.pone.0161163. eCollection 2016 Reproduced under the Creative Commons license <http://creativecommons.org/licenses/by/4.0/>



Western blot - Anti-Glucose Transporter GLUT1 antibody [EPR3915] (ab115730)

All lanes : Anti-Glucose Transporter GLUT1 antibody [EPR3915] (ab115730) at 1/50000 dilution

Lane 1 : HT-29 (Human colorectal adenocarcinoma epithelial cell) whole cell lysates unboiled with 5% NFDN/TBST

Lane 2 : HT-29 (Human colorectal adenocarcinoma epithelial cell) whole cell lysates boiled with 5% NFDN/TBST

Lane 3 : 3T3-L1 (Mouse embryonic fibroblast) whole cell lysates unboiled with 5% NFDN/TBST

Lane 4 : 3T3-L1 (Mouse embryonic fibroblast) whole cell lysates boiled with 5% NFDN/TBST

Lysates/proteins at 20 µg per lane.

Secondary

All lanes : Goat Anti-Rabbit IgG H&L (HRP) (**ab97051**) at 1/20000 dilution

Predicted band size: 54 kDa


Observed band size: 40-60 kDa

Exposure time

Lane 1 to 2: 10 seconds


Lane 3 to 4: 30 seconds

We recommend not to boil the samples after lysis to get desired WB bands.




Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-Glucose Transporter GLUT1 antibody [EPR3915] (ab115730)

Immunohistochemical staining of paraffin embedded rat kidney with purified ab115730 at a working dilution of 1/500. The secondary antibody used is **ab97051**, a goat anti-rabbit IgG (H&L) at a dilution of 1/500. The sample is counter-stained with hematoxylin. Antigen retrieval was performed using Tris-EDTA buffer, pH 9.0. PBS was used instead of the primary antibody as the negative control, and is shown in the inset.




Immunocytochemistry/ Immunofluorescence - Anti-Glucose Transporter GLUT1 antibody [EPR3915] (ab115730)

ab115730 staining SLC2A1 in wild-type A549 cells, with negative expression in SLC2A1 knockout A549 cells. The cells were fixed with 100% methanol (5 min), permeabilised with 0.1% Triton x-100 for 5 minutes and then blocked with 1% BSA/10% normal goat serum/0.3M glycine in 0.1% PBS-Tween for 1h. The cells were then incubated overnight at +4°C with ab115730 at 1 µg/ml and **ab7291**, Mouse monoclonal [DM1A] to alpha Tubulin at 0.5 µg/ml. Cells were then incubated with **ab150081**, Goat polyclonal Secondary Antibody to Rabbit IgG - H&L (Alexa Fluor® 488), pre-adsorbed at 1/1000 dilution (shown in green) and **ab150119**, Goat polyclonal Secondary Antibody to Mouse IgG - H&L (Alexa Fluor® 647), pre-adsorbed at 1/1000 dilution (shown in red). Nuclear DNA was labelled with DAPI (shown in blue). Image was acquired with a confocal microscope (Leica-Microsystems TCS SP8) and a single confocal section is shown.




Immunocytochemistry/ Immunofluorescence - Anti-Glucose Transporter GLUT1 antibody [EPR3915] (ab115730)

Immunofluorescence staining of HepG2 cells with purified ab115730 at a working dilution of 1/100, counter-stained with DAPI. The secondary antibody was Alexa Fluor[®] 488 goat anti-rabbit (**ab150077**), used at a dilution of 1/1000. **ab7291**, a mouse anti-tubulin antibody (1/1000), was used to stain tubulin along with **ab150120** (Alexa Fluor[®] 594 goat anti-mouse, 1/1000), shown in the top right hand panel. The cells were fixed in 4% PFA and permeabilized using 0.1% Triton X 100. The negative controls are shown in bottom middle and right hand panels - for negative control 1, purified ab115730 was used at a dilution of 1/500 followed by an Alexa Fluor[®] 594 goat anti-mouse antibody (**ab150120**) at a dilution of 1/500. For negative control 2, **ab7291** (mouse anti-tubulin) was used at a dilution of 1/500 followed by an Alexa Fluor[®] 488 goat anti-rabbit antibody (**ab150077**) at a dilution of 1/400. Alexa Fluor[®] 488 (**ab195359**) and Alexa Fluor[®] 647 (**ab195020**) conjugated versions are available for this clone.




Flow Cytometry (Intracellular) - Anti-Glucose Transporter GLUT1 antibody [EPR3915] (ab115730)

Overlay histogram showing Jurkat cells fixed in 4% PFA and stained with purified ab115730 at a dilution of 1/40 (red line). The secondary antibody used was Alexa Fluor[®] 488 goat anti-rabbit at a dilution of 1/500. Rabbit monoclonal IgG was used as an isotype control (black line) and cells incubated in the absence of both primary and secondary antibody were used as a negative control (blue line). Alexa Fluor[®]488 (**ab195359**) and Alexa Fluor[®]647 (**ab195020**) conjugated versions are available for this clone.




Immunohistochemical staining of paraffin embedded mouse liver with purified ab115730 at a working dilution of 1/500. The secondary antibody used is **ab97051**, a goat anti-rabbit IgG (H&L) at a dilution of 1/500. The sample is counter-stained with hematoxylin. Antigen retrieval was performed using Tris-EDTA buffer, pH 9.0. PBS was used instead of the primary antibody as the negative control, and is shown in the inset.

Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-Glucose Transporter GLUT1 antibody [EPR3915] (ab115730)




Immunohistochemical staining of paraffin embedded human lung carcinoma with purified ab115730 at a working dilution of 1/500. The secondary antibody used is **ab97051**, a goat anti-rabbit IgG (H&L) at a dilution of 1/500. The sample is counter-stained with hematoxylin. Antigen retrieval was performed using Tris-EDTA buffer, pH 9.0. PBS was used instead of the primary antibody as the negative control, and is shown in the inset.

Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-Glucose Transporter GLUT1 antibody [EPR3915] (ab115730)



Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-Glucose Transporter GLUT1 antibody [EPR3915] (ab115730)

Immunohistochemical staining of paraffin embedded human cervical carcinoma with purified ab115730 at a working dilution of 1/500. The secondary antibody used is **ab97051**, a goat anti-rabbit IgG (H&L) at a dilution of 1/500. The sample is counter-stained with hematoxylin. Antigen retrieval was performed using Tris-EDTA buffer, pH 9.0. PBS was used instead of the primary antibody as the negative control, and is shown in the inset.



Western blot - Anti-Glucose Transporter GLUT1 antibody [EPR3915] (ab115730)

All lanes : Anti-Glucose Transporter GLUT1 antibody [EPR3915] (ab115730) at 1/1000000 dilution (purified)

Lane 1 : HepG2 whole cell lysate

Lane 2 : Human fetal liver lysate

Lane 3 : HT-29 whole cell lysate

Lane 4 : SW480 whole cell lysate

Lysates/proteins at 10 µg per lane.

Secondary


All lanes : Anti-rabbit IgG (HRP), specific to the non-reduced form of IgG at 1/1000 dilution

Predicted band size: 54 kDa

Observed band size: 40-60 kDa

Blocking buffer: 5% NFDm/TBST

Dilution buffer: 5% NFDm/TBST



Western blot - Anti-Glucose Transporter GLUT1 antibody [EPR3915] (ab115730)

All lanes : Anti-Glucose Transporter GLUT1 antibody [EPR3915] (ab115730) at 1/1000 dilution (Unpurified)

Lane 1 : Jurkat lysate

Lane 2 : Mouse brain lysate

Lane 3 : Human fetal brain lysate


Lane 4 : 3T3L1 lysate

Lane 5 : Human fetal liver lysate

Lane 6 : HepG2 lysate

Lysates/proteins at 10 µg per lane.


Predicted band size: 54 kDa



Flow Cytometry (Intracellular) - Anti-Glucose Transporter GLUT1 antibody [EPR3915] (ab115730)

Overlay histogram showing HeLa cells stained with unpurified ab115730 (red line). The cells were fixed with 80% methanol (5 min) and then permeabilized with 0.1% PBS-Tween for 20 min. The cells were then incubated in 1x PBS / 10% normal goat serum / 0.3M glycine to block non-specific protein-protein interactions followed by the antibody (ab115730, 1/1000 dilution) for 30 min at 22°C. The secondary antibody used was Alexa Fluor® 488 goat anti-rabbit IgG (H+L) (**ab150077**) at 1/2000 dilution for 30 min at 22°C. Isotype control antibody (black line) was rabbit IgG (monoclonal) (1µg/1x10⁶ cells) used under the same conditions. Unlabelled sample (blue line) was also used as a control.


Acquisition of >5,000 events were collected using a 20mW Argon ion laser (488nm) and 525/30 bandpass filter. This antibody gave a positive signal in HeLa cells fixed with 4% paraformaldehyde (10 min)/permeabilized with 0.1% PBS-Tween for 20 min used under the same conditions.



Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-Glucose Transporter GLUT1 antibody [EPR3915] (ab115730)

Unpurified ab115730 at 1/250 dilution staining Glucose Transporter GLUT1 in Paraffin-embedded human cervical carcinoma tissue by Immunohistochemistry.


Perform heat mediated antigen retrieval before commencing with IHC staining protocol.



Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-Glucose Transporter GLUT1 antibody [EPR3915] (ab115730)

Unpurified ab115730 at 1/250 dilution staining Glucose Transporter GLUT1 in Paraffin-embedded human colonic adenocarcinoma tissue by Immunohistochemistry.


Perform heat mediated antigen retrieval before commencing with IHC staining protocol.



Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-Glucose Transporter GLUT1 antibody [EPR3915] (ab115730)

Unpurified ab115730 showing positive staining in human normal liver tissue.


Perform heat mediated antigen retrieval before commencing with IHC staining protocol.



Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-Glucose Transporter GLUT1 antibody [EPR3915] (ab115730)

Unpurified ab115730 showing positive staining in human normal breast tissue.


Perform heat mediated antigen retrieval before commencing with IHC staining protocol.



Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-Glucose Transporter GLUT1 antibody [EPR3915] (ab115730)

Unpurified ab115730 showing positive staining in human normal colon tissue.


Perform heat mediated antigen retrieval before commencing with IHC staining protocol.



Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-Glucose Transporter GLUT1 antibody [EPR3915] (ab115730)

Unpurified ab115730 showing positive staining in human kidney carcinoma tissue.


Perform heat mediated antigen retrieval before commencing with IHC staining protocol.



Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-Glucose Transporter GLUT1 antibody [EPR3915] (ab115730)

Unpurified ab115730 showing negative staining in human skeletal muscle tissue.


Perform heat mediated antigen retrieval before commencing with IHC staining protocol.



Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-Glucose Transporter GLUT1 antibody [EPR3915] (ab115730)

Unpurified ab115730 showing positive staining in human urinary bladder transitional carcinoma tissue.


Perform heat mediated antigen retrieval before commencing with IHC staining protocol.



Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-Glucose Transporter GLUT1 antibody [EPR3915] (ab115730)

Unpurified ab115730 showing negative staining in human normal heart tissue.

Perform heat mediated antigen retrieval before commencing with IHC staining protocol.



OI-RD Scanning - Anti-Glucose Transporter GLUT1 antibody [EPR3915] (ab115730)

Equilibrium disassociation constant (K_D)

Learn more about K_D

[Click here to learn more about \$K_D\$](#)

Why choose a recombinant antibody?



Research with confidence
Consistent and reproducible results



Long-term and scalable supply
Recombinant technology



Success from the first experiment
Confirmed specificity



Ethical standards compliant
Animal-free production

Anti-Glucose Transporter GLUT1 antibody
[EPR3915] (ab115730)

Please note: All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <https://www.abcam.com/abpromise> or contact our technical team.

Terms and conditions

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors