

Product datasheet

Anti-Glycophorin A antibody [EPR8200] α b129024

Recombinant RabMAb

★★★★★ [1 Abreviews](#) [11 References](#) [14 Images](#)


Overview

Product name	Anti-Glycophorin A antibody [EPR8200]
Description	Rabbit monoclonal [EPR8200] to Glycophorin A
Host species	Rabbit
Tested applications	Suitable for: Flow Cyt (Intra), WB, IHC-P
Species reactivity	Reacts with: Human
Immunogen	Synthetic peptide within Human Glycophorin A aa 100 to the C-terminus. The exact sequence is proprietary. Database link: P02724
Positive control	Fetal liver lysate, Human lung tissue, Human spleen tissue.
General notes	<p>This product is a recombinant monoclonal antibody, which offers several advantages including:</p> <ul style="list-style-type: none"> - High batch-to-batch consistency and reproducibility - Improved sensitivity and specificity - Long-term security of supply - Animal-free production <p>For more information see here.</p> <p>Our RabMAb[®] technology is a patented hybridoma-based technology for making rabbit monoclonal antibodies. For details on our patents, please refer to RabMAb[®] patents.</p> <p>We are constantly working hard to ensure we provide our customers with best in class antibodies. As a result of this work we are pleased to now offer this antibody in purified format. We are in the process of updating our datasheets. The purified format is designated 'PUR' on our product labels. If you have any questions regarding this update, please contact our Scientific Support team.</p>

Properties

Form	Liquid
Storage instructions	Shipped at 4°C. Store at -20°C. Stable for 12 months at -20°C.
Dissociation constant (K_D)	K _D = 2.38 x 10 ⁻¹⁰ M





[Learn more about K_D](#)

Storage buffer	pH: 7.20 Preservative: 0.01% Sodium azide Constituents: 59% PBS, 0.05% BSA, 40% Glycerol
Purity	Protein A purified
Clonality	Monoclonal
Clone number	EPR8200
Isotype	IgG

Applications

The Abpromise guarantee Our **Abpromise guarantee** covers the use of ab129024 in the following tested applications. The application notes include recommended starting dilutions; optimal dilutions/concentrations should be determined by the end user.

Application	Abreviews	Notes
Flow Cyt (Intra)		1/90. ab172730 - Rabbit monoclonal IgG, is suitable for use as an isotype control with this antibody.
WB		1/1000. Detects a band of approximately 38 kDa (predicted molecular weight: 16 kDa).
IHC-P	★★★★★ (1)	1/2500 - 1/5000. Perform heat mediated antigen retrieval before commencing with IHC staining protocol. See IHC antigen retrieval protocols . For unpurified use at 1/100 - 1/250.

Target

Function Glycophorin A is the major intrinsic membrane protein of the erythrocyte. The N-terminal glycosylated segment, which lies outside the erythrocyte membrane, has MN blood group receptors. Appears to be important for the function of SLC4A1 and is required for high activity of SLC4A1. May be involved in translocation of SLC4A1 to the plasma membrane. Is a receptor for influenza virus. Is a receptor for Plasmodium falciparum erythrocyte-binding antigen 175 (EBA-175); binding of EBA-175 is dependent on sialic acid residues of the O-linked glycans. Appears to be a receptor for Hepatitis A virus (HAV).

Sequence similarities Belongs to the glycophorin A family.


Post-translational modifications The major O-linked glycan are NeuAc-alpha-(2-3)-Gal-beta-(1-3)-[NeuAc-alpha-(2-6)]-GalNAcOH (about 78 %) and NeuAc-alpha-(2-3)-Gal-beta-(1-3)-GalNAcOH (17 %). Minor O-glycans (5 %) include NeuAc-alpha-(2-3)-Gal-beta-(1-3)-[NeuAc-alpha-(2-6)]-GalNAcOH NeuAc-alpha-(2-8)-NeuAc-alpha-(2-3)-Gal-beta-(1-3)-GalNAcOH. About 1% of all O-linked glycans carry blood group A, B and H determinants. They derive from a type-2 precursor core structure, Gal-beta-(1,3)-GlcNAc-beta-1-R, and the antigens are synthesized by addition of fucose (H antigen-specific) and then N-acetylgalactosamine (A antigen-specific) or galactose (B antigen-specific). Specifically O-

linked-glycans are NeuAc-alpha-(2-3)-Gal-beta-(1-3)-GalNAcOH-(6-1)-GlcNAc-beta-(4-1)-[Fuc-alpha-(1-2)]-Gal-beta-(3-1)-GalNAc-alpha (about 1%, B antigen-specific) and NeuAc-alpha-(2-3)-Gal-beta-(1-3)-GalNAcOH-(6-1)-GlcNAc-beta-(4-1)-[Fuc-alpha-(1-2)]-Gal-beta (1 %, O antigen-, A antigen- and B antigen-specific).

Cellular localization


Cell membrane. Appears to be colocalized with SLC4A1.

Images




Flow Cytometry (Intracellular) - Anti-Glycophorin A antibody [EPR8200] (ab129024)

Overlay histogram showing K562 cells stained with unpurified ab129024 (red line). The cells were fixed with 80% methanol (5 min) and then permeabilized with 0.1% PBS-Tween for 20 min. The cells were then incubated in 1x PBS / 10% normal goat serum / 0.3M glycine to block non-specific protein-protein interactions followed by the antibody (ab129024, 1/100 dilution) for 30 min at 22°C. The secondary antibody used was Alexa Fluor® 488 goat anti-rabbit IgG (H&L) ([ab150077](#)) at 1/2000 dilution for 30 min at 22°C. Isotype control antibody (black line) was rabbit IgG (monoclonal) (1µg/1x10⁶ cells) used under the same conditions. Unlabelled sample (blue line) was also used as a control. Acquisition of >5,000 events were collected using a 20mW Argon ion laser (488nm) and 525/30 bandpass filter.



Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-Glycophorin A antibody [EPR8200] (ab129024)

ab129024 staining Glycophorin A in Human placenta tissue sections by Immunohistochemistry (IHC-P - paraformaldehyde-fixed, paraffin-embedded sections). Tissue was fixed and paraffin-embedded, antigen retrieval was by heat mediation in Tris/EDTA buffer pH9. Samples were incubated with primary antibody (1/2500). [ab97051](#)(1/500) HRP-conjugated goat anti-rabbit IgG(H&L) was used as the secondary antibody. Tissue counterstained with Hematoxylin. PBS was used in the negative control rather than the Primary antibody.




Western blot - Anti-Glycophorin A antibody
[EPR8200] (ab129024)

Anti-Glycophorin A antibody [EPR8200] (ab129024) at 1/1000
dilution + Human fetal kidney at 10 µg

Secondary

Goat Anti-Rabbit IgG, (H+L), HRP- conjugated at 1/1000 dilution

Predicted band size: 16 kDa




Western blot - Anti-Glycophorin A antibody
[EPR8200] (ab129024)

Anti-Glycophorin A antibody [EPR8200] (ab129024) at 1/2000
dilution + Human fetal liver lysate at 20 µg

Secondary

Goat Anti-Rabbit IgG, (H+L), HRP- conjugated at 1/1000 dilution

Predicted band size: 16 kDa




Western blot - Anti-Glycophorin A antibody
[EPR8200] (ab129024)

Anti-Glycophorin A antibody [EPR8200] (ab129024) at 1/1000 dilution (unpurified) + Fetal liver lysate at 10 μ g

Predicted band size: 16 kDa


Observed band size: 38 kDa



Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-Glycophorin A antibody
[EPR8200] (ab129024)

ab129024, at 1/100 dilution staining Glycophorin A in formalin fixed paraffin embedded Human lung tissue by immunohistochemistry.


Perform heat mediated antigen retrieval before commencing with IHC staining protocol.



Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-Glycophorin A antibody
[EPR8200] (ab129024)

ab129024, unpurified, at 1/100 dilution staining Glycophorin A in formalin fixed paraffin embedded Human spleen tissue by immunohistochemistry.


Perform heat mediated antigen retrieval before commencing with IHC staining protocol.



Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-Glycophorin A antibody [EPR8200] (ab129024)

ab129024, unpurified, showing positive staining in Thyroid gland erythrocytes tissue.


Perform heat mediated antigen retrieval before commencing with IHC staining protocol.



Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-Glycophorin A antibody [EPR8200] (ab129024)

ab129024, unpurified, showing negative staining in Normal brain tissue.


Perform heat mediated antigen retrieval before commencing with IHC staining protocol.



Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-Glycophorin A antibody [EPR8200] (ab129024)

ab129024, unpurified, showing positive staining in Normal colon erythrocytes tissue.


Perform heat mediated antigen retrieval before commencing with IHC staining protocol.



ab129024, unpurified, showing positive staining in Normal kidney erythrocytes tissue.

Perform heat mediated antigen retrieval before commencing with IHC staining protocol.


Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-Glycophorin A antibody [EPR8200] (ab129024)



ab129024, unpurified, showing positive staining in Normal placenta erythrocytes tissue.

Perform heat mediated antigen retrieval before commencing with IHC staining protocol.

Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-Glycophorin A antibody [EPR8200] (ab129024)



Equilibrium disassociation constant (K_D)

Learn more about K_D

[Click here to learn more about \$K_D\$](#)

OIR-D Scanning - Anti-Glycophorin A antibody [EPR8200] (ab129024)

Why choose a recombinant antibody?



Research with confidence
Consistent and reproducible results



Long-term and scalable supply
Recombinant technology



Success from the first experiment
Confirmed specificity



Ethical standards compliant
Animal-free production

Anti-Glycophorin A antibody [EPR8200] (ab129024)

Please note: All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <https://www.abcam.com/abpromise> or contact our technical team.

Terms and conditions

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors