

Anti-GTPase HRAS antibody [Y132] - BSA and Azide free ab239815


KO VALIDATED

Recombinant

RabMAb

5 Images

Overview

Product name	Anti-GTPase HRAS antibody [Y132] - BSA and Azide free
Description	Rabbit monoclonal [Y132] to GTPase HRAS - BSA and Azide free
Host species	Rabbit
Specificity	Reactivity with other RAS members has not been tested.
Tested applications	Suitable for: IP, WB Unsuitable for: IHC
Species reactivity	Reacts with: Mouse, Rat, Human Predicted to work with: Chicken 
Immunogen	Synthetic peptide. This information is proprietary to Abcam and/or its suppliers.
Positive control	MCF7 and PC12 cell lysates and MCF7 cells.
General notes	<p>ab239815 is the carrier-free version of ab32417.</p> <p>Our carrier-free antibodies are typically supplied in a PBS-only formulation, purified and free of BSA, sodium azide and glycerol. The carrier-free buffer and high concentration allow for increased conjugation efficiency.</p> <p>This conjugation-ready format is designed for use with fluorochromes, metal isotopes, oligonucleotides, and enzymes, which makes them ideal for antibody labelling, functional and cell-based assays, flow-based assays (e.g. mass cytometry) and Multiplex Imaging applications.</p> <p>Use our conjugation kits for antibody conjugates that are ready-to-use in as little as 20 minutes with <1 minute hands-on-time and 100% antibody recovery: available for fluorescent dyes, HRP, biotin and gold.</p> <p>This product is compatible with the Maxpar[®] Antibody Labeling Kit from Fluidigm, without the need for antibody preparation. Maxpar[®] is a trademark of Fluidigm Canada Inc.</p> <p>This product is a recombinant monoclonal antibody, which offers several advantages including:</p> <ul style="list-style-type: none"> - High batch-to-batch consistency and reproducibility - Improved sensitivity and specificity - Long-term security of supply - Animal-free production <p>For more information see here.</p>

Our RabMAb[®] technology is a patented hybridoma-based technology for making rabbit monoclonal antibodies. For details on our patents, please refer to [RabMAb[®] patents](#).

Properties

Form	Liquid
Storage instructions	Shipped at 4°C. Store at +4°C. Do Not Freeze.
Storage buffer	pH: 7.2 Constituent: PBS
Carrier free	Yes
Purity	Protein A purified
Clonality	Monoclonal
Clone number	Y132
Isotype	IgG

Applications

The Abpromise guarantee Our [Abpromise guarantee](#) covers the use of ab239815 in the following tested applications. The application notes include recommended starting dilutions; optimal dilutions/concentrations should be determined by the end user.

Application	Abreviews	Notes
IP		1/50 - 1/60.
WB		1/500 - 1/1000. Detects a band of approximately 21 kDa (predicted molecular weight: 21 kDa).

Application notes Is unsuitable for IHC.

Target

Function	Ras proteins bind GDP/GTP and possess intrinsic GTPase activity.
Involvement in disease	<p>Defects in HRAS are the cause of faciocutaneoskeletal syndrome (FCSS) [MIM:218040]. A rare condition characterized by prenatally increased growth, postnatal growth deficiency, mental retardation, distinctive facial appearance, cardiovascular abnormalities (typically pulmonic stenosis, hypertrophic cardiomyopathy and/or atrial tachycardia), tumor predisposition, skin and musculoskeletal abnormalities.</p> <p>Defects in HRAS are the cause of congenital myopathy with excess of muscle spindles (CMEMS) [MIM:218040]. CMEMS is a variant of Costello syndrome.</p> <p>Defects in HRAS may be a cause of susceptibility to Hurtle cell thyroid carcinoma (HCTC) [MIM:607464]. Hurtle cell thyroid carcinoma accounts for approximately 3% of all thyroid cancers. Although they are classified as variants of follicular neoplasms, they are more often multifocal and somewhat more aggressive and are less likely to take up iodine than are other follicular neoplasms.</p> <p>Note=Mutations which change positions 12, 13 or 61 activate the potential of HRAS to transform cultured cells and are implicated in a variety of human tumors.</p>

Defects in HRAS are a cause of susceptibility to bladder cancer (BLC) [MIM:109800]. A malignancy originating in tissues of the urinary bladder. It often presents with multiple tumors appearing at different times and at different sites in the bladder. Most bladder cancers are transitional cell carcinomas. They begin in cells that normally make up the inner lining of the bladder. Other types of bladder cancer include squamous cell carcinoma (cancer that begins in thin, flat cells) and adenocarcinoma (cancer that begins in cells that make and release mucus and other fluids). Bladder cancer is a complex disorder with both genetic and environmental influences.

Note=Defects in HRAS are the cause of oral squamous cell carcinoma (OSCC).

Sequence similarities

Belongs to the small GTPase superfamily. Ras family.

Post-translational modifications


Palmitoylated by the ZDHHC9-GOLGA7 complex. A continuous cycle of de- and re-palmitoylation regulates rapid exchange between plasma membrane and Golgi.

S-nitrosylated; critical for redox regulation. Important for stimulating guanine nucleotide exchange. No structural perturbation on nitrosylation.

Cellular localization

Cell membrane. Golgi apparatus membrane. The active GTP-bound form is localized most strongly to membranes than the inactive GDP-bound form (By similarity). Shuttles between the plasma membrane and the Golgi apparatus.

Images



Western blot - Anti-GTPase HRAS antibody [Y132] - BSA and Azide free (ab239815)

All lanes : Anti-GTPase HRAS antibody [Y132] ([ab32417](#)) at 1/1000 dilution (purified)

Lane 1 : MCF7 cell lysate

Lane 2 : HeLa cell lysate

Lysates/proteins at 10 µg per lane.

Secondary


All lanes : HRP goat anti-rabbit IgG (H+L) at 1/20000 dilution

Predicted band size: 21 kDa

This data was developed using [ab32417](#), the same antibody clone in a different buffer formulation.

Blocking buffer: 5% NFDm/TBST

Dilution buffer: 5% NFDm/TBST



All lanes : Anti-GTPase HRAS antibody [Y132] ([ab32417](#)) at 1/1000 dilution

Lane 1 : MCF7 cell lysate

Lane 2 : HeLa cell lysate

Lysates/proteins at 10 µg per lane.

Secondary

All lanes : HRP goat anti-rabbit IgG (H+L) at 1/20000 dilution


Predicted band size: 21 kDa

Observed band size: 21 kDa

This data was developed using [ab32417](#), the same antibody clone in a different buffer formulation.

Blocking buffer: 5% NFDM/TBST

Dilution buffer: 5% NFDM/TBST



All lanes : Anti-GTPase HRAS antibody [Y132] ([ab32417](#)) at 1/500 dilution

Lane 1 : Wild-type HEK-293 whole cell lysate

Lane 2 : HRAS knockout HEK-293 whole cell lysate

Lane 3 : HeLa whole cell lysate

Lane 4 : MCF7 whole cell lysate

Lysates/proteins at 20 µg/ml per lane.


Predicted band size: 21 kDa

This data was developed using [ab32417](#), the same antibody clone in a different buffer formulation.

Lanes 1 -4: Merged signal (red and green). Green - [ab32417](#) observed at 21 kDa. Red - loading control, [ab8245](#), observed at 37

kDa.

ab32417 was shown to recognize HRAS in wild-type HEK-293 cells as signal was lost at the expected MW in HRAS knockout cells. Additional cross-reactive bands were observed in the wild-type and knockout cells. Wild-type and HRAS knockout samples were subjected to SDS-PAGE. The membrane was blocked with 3% Milk. Ab32417 and **ab8245** (Mouse anti-GAPDH loading control) were incubated overnight at 4°C at 1/500 dilution and 1/20000 dilution respectively. Blots were developed with Goat anti-Rabbit IgG H&L (IRDye® 800CW) preabsorbed **ab216773** and Goat anti-Mouse IgG H&L (IRDye® 680RD) preabsorbed **ab216776** secondary antibodies at 1/20000 dilution for 1 hour at room temperature before imaging.




Immunoprecipitation - Anti-GTPase HRAS antibody
[Y132] - BSA and Azide free (ab239815)

ab32417 (purified) at 1/60 immunoprecipitating GTPase in 10 µg mouse brain whole cell lysate (Lanes 1 and 2, observed at 21 kDa). Lane 3 - PBS. For western blotting, VeriBlot for IP Detection Reagent (HRP) (**ab131366**), was used for detection at 1/10,000 dilution. Blocking buffer and concentration: 5% NFDM/TBST Dilution buffer and concentration: 5% NFDM/TBST

This data was developed using the same antibody clone in a different buffer formulation containing PBS, BSA, glycerol, and sodium azide (**ab32417**).

Why choose a recombinant antibody?



Anti-GTPase HRAS antibody [Y132] - BSA and
Azide free (ab239815)

Please note: All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <https://www.abcam.com/abpromise> or contact our technical team.

Terms and conditions

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors