

## Product datasheet

### Anti-Hairy Cell Leukemia antibody [DBA.44] ab779

[3 References](#) [1 Image](#)

#### Overview

<b>Product name</b>	Anti-Hairy Cell Leukemia antibody [DBA.44]
<b>Description</b>	Mouse monoclonal [DBA.44] to Hairy Cell Leukemia
<b>Host species</b>	Mouse
<b>Tested applications</b>	<b>Suitable for:</b> IHC-Fr, IHC-P
<b>Species reactivity</b>	<b>Reacts with:</b> Human
<b>Immunogen</b>	Tissue, cells or virus corresponding to Hairy Cell Leukemia. Deau cell line established from a large cell lymphoma
<b>General notes</b>	<p>The Life Science industry has been in the grips of a reproducibility crisis for a number of years. Abcam is leading the way in addressing this with our range of recombinant monoclonal antibodies and knockout edited cell lines for gold-standard validation. Please check that this product meets your needs before purchasing.</p> <p>If you have any questions, special requirements or concerns, please send us an inquiry and/or contact our Support team ahead of purchase. Recommended alternatives for this product can be found below, along with publications, customer reviews and Q&amp;As</p>

#### Properties

<b>Form</b>	Liquid
<b>Storage instructions</b>	Shipped at 4°C. Store at +4°C short term (1-2 weeks). Store at -20°C or -80°C. Avoid freeze / thaw cycle.
<b>Storage buffer</b>	pH: 7.3 Preservative: 0.05% Sodium azide Constituents: Tissue culture supernatant, 1% BSA
<b>Purity</b>	Tissue culture supernatant
<b>Clonality</b>	Monoclonal
<b>Clone number</b>	DBA.44
<b>Isotype</b>	IgM
<b>Light chain type</b>	kappa

#### Applications

## The Abpromise guarantee

Our **Abpromise guarantee** covers the use of ab779 in the following tested applications.

The application notes include recommended starting dilutions; optimal dilutions/concentrations should be determined by the end user.


Application	Abreviews	Notes
IHC-Fr		Use at an assay dependent concentration.
IHC-P		1/15 - 1/30. Perform heat mediated antigen retrieval with citrate buffer pH 6 before commencing with IHC staining protocol. Primary incubation for 1 hour at room temperature.

## Target

### Cellular localization

Nuclear

## Images



Formalin fixed paraffin embedded human tonsil tissue, staining Hairy Cell Leukemia with ab779 in immunohistochemical analysis

Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-Hairy Cell Leukemia antibody [DBA.44] (ab779)

**Please note:** All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

## Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you

- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <https://www.abcam.com/abpromise> or contact our technical team.

#### **Terms and conditions**

---

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors