# abcam

### Product datasheet

## Anti-Hepatitis G Virus antibody ab 1037

Overview

Product name Anti-Hepatitis G Virus antibody

**Description** Rabbit polyclonal to Hepatitis G Virus

Host species Rabbit

**Specificity** This antibody recognizes the NS5 polyprotein of HGV, also known as GBV-C/HBV.

Tested applications Suitable for: WB

Species reactivity Reacts with: GB virus C

Immunogen Peptide corresponding to amino acids 2268-2276 of the NS5 non-structural polyprotein of

Hepatitis G Virus (HGV).

**General notes**The Life Science industry has been in the grips of a reproducibility crisis for a number of years.

Abcam is leading the way in addressing this with our range of recombinant monoclonal antibodies and knockout edited cell lines for gold-standard validation. Please check that this product meets

your needs before purchasing.

If you have any questions, special requirements or concerns, please send us an inquiry and/or contact our Support team ahead of purchase. Recommended alternatives for this product can be

found below, along with publications, customer reviews and Q&As

**Properties** 

Form Liquid

**Storage instructions** Shipped at 4°C. Upon delivery aliquot and store at -20°C or -80°C. Avoid repeated freeze / thaw

cycles.

Storage buffer pH: 7.4

Preservative: 0.05% Sodium azide

**Purity** Immunogen affinity purified

Clonality Polyclonal Isotype unknown
Light chain type unknown

**Applications** 

The Abpromise guarantee Our Abpromise guarantee covers the use of ab1037 in the following tested applications.

1

The application notes include recommended starting dilutions; optimal dilutions/concentrations should be determined by the end user.

Application	Abreviews	Notes
WB		1/500 - 1/1000. Detects a band of approximately 41 kDa.

#### **Target**

#### Relevance

Hepatitis G virus is from a group of viruses called GB viruses (GB virus A, GB virus B and GB virus C) and is also designated as GB virus C (GBV C). HGV is a positive stranded RNA virus, with a genome of 10 KB and appears, on the basis of genome length and organisation, to be a member of the Flaviviridae and distinct from the hepatitis C virus (HCV). HGV infection appears to be transmitted following receipt of blood from HGV RNA positive donors, however, these patients have only moderate liver enzyme elevations and do not develop chronic liver disease.

Please note: All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

#### Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- · Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <a href="https://www.abcam.com/abpromise">https://www.abcam.com/abpromise</a> or contact our technical team.

#### Terms and conditions

· Guarantee only valid for products bought direct from Abcam or one of our authorized distributors