

Product datasheet

Anti-HNF-1B antibody [EPR18644-13] ab213149

Recombinant **RabMAb**

★★★★★ **1 Abreviews** [9 Images](#)

Overview

Product name	Anti-HNF-1B antibody [EPR18644-13]
Description	Rabbit monoclonal [EPR18644-13] to HNF-1B
Host species	Rabbit
Tested applications	Suitable for: IHC-P
Species reactivity	Reacts with: Mouse, Rat, Human
Immunogen	Recombinant fragment. This information is proprietary to Abcam and/or its suppliers.
Positive control	IHC-P: Human liver, liver bile duct carcinoma and ovarian clear cell carcinoma tissues. Mouse and rat liver tissues.
General notes	<p>This product is a recombinant monoclonal antibody, which offers several advantages including:</p> <ul style="list-style-type: none"> - High batch-to-batch consistency and reproducibility - Improved sensitivity and specificity - Long-term security of supply - Animal-free production <p>For more information see here.</p> <p>Our RabMAb[®] technology is a patented hybridoma-based technology for making rabbit monoclonal antibodies. For details on our patents, please refer to RabMAb[®] patents.</p>

Properties

Form	Liquid
Storage instructions	Shipped at 4°C. Store at +4°C short term (1-2 weeks). Upon delivery aliquot. Store at -20°C long term. Avoid freeze / thaw cycle.
Storage buffer	<p>pH: 7.2</p> <p>Preservative: 0.01% Sodium azide</p> <p>Constituents: 59% PBS, 40% Glycerol (glycerin, glycerine), 0.05% BSA</p>
Purity	Protein A purified
Clonality	Monoclonal
Clone number	EPR18644-13
Isotype	IgG

Applications


The Abpromise guarantee Our **Abpromise guarantee** covers the use of ab213149 in the following tested applications. The application notes include recommended starting dilutions; optimal dilutions/concentrations should be determined by the end user.

Application	Abreviews	Notes
IHC-P		1/2000. Perform heat mediated antigen retrieval with Tris/EDTA buffer pH 9.0 before commencing with IHC staining protocol.

Target

Function	Transcription factor, probably binds to the inverted palindrome 5'-GTTAATNATTAAC-3'.
Involvement in disease	<p>Defects in HNF1B are the cause of renal cysts and diabetes syndrome (RCAD) [MIM:137920]; also called maturity-onset diabetes of the young type 5 (MODY5) or familial hypoplastic glomerulocystic kidney disease (GCKD). RCAD is an autosomal dominant disorder comprising non-diabetic renal disease resulting from abnormal renal development, and diabetes, which in some cases occurs earlier than age 25 years and is thus consistent with a diagnosis of maturity-onset diabetes of the young (MODY5). The renal disease is highly variable and includes renal cysts, glomerular tufts, aberrant nephrogenesis, primitive tubules, irregular collecting systems, oligomeganephronia, enlarged renal pelvis, abnormal calyces, small kidney, single kidney, horseshoe kidney, and hyperuricemic nephropathy.</p> <p>Defects in HNF1B may be rare genetic risk factor contributing to the development of non-insulin-dependent diabetes mellitus (NIDDM) [MIM:125853]. NIDDM is characterized by an autosomal dominant mode of inheritance, onset during adulthood and insulin resistance.</p> <p>Defects in HNF1B may be a cause of susceptibility to prostate cancer hereditary type 11 (HPC11) [MIM:611955]. It is a condition associated with familial predisposition to cancer of the prostate. Most prostate cancers are adenocarcinomas that develop in the acini of the prostatic ducts. Other rare histopathologic types of prostate cancer that occur in approximately 5% of patients include small cell carcinoma, mucinous carcinoma, prostatic ductal carcinoma, transitional cell carcinoma, squamous cell carcinoma, basal cell carcinoma, adenoid cystic carcinoma (basaloid), signet-ring cell carcinoma and neuroendocrine carcinoma.</p>
Sequence similarities	<p>Belongs to the HNF1 homeobox family.</p> <p>Contains 1 homeobox DNA-binding domain.</p>
Cellular localization	Nucleus.

Images




Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-HNF-1B antibody
[EPR18644-13] (ab213149)

Immunohistochemical analysis of paraffin-embedded mouse liver tissue labeling HNF-1B with ab213149 at 1/4000 dilution, followed by LeicaDS9800 (Bond™ Polymer Refine Detection). Nuclear staining on mouse liver bile duct is observed. Counter stained with Hematoxylin.

Secondary antibody only control: Used PBS instead of primary antibody

Perform heat mediated antigen retrieval with Tris/EDTA buffer pH 9.0 before commencing with IHC staining protocol.




Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-HNF-1B antibody
[EPR18644-13] (ab213149)

Immunohistochemical analysis of paraffin-embedded rat liver tissue labeling HNF-1B with ab213149 at 1/4000 dilution, followed by LeicaDS9800 (Bond™ Polymer Refine Detection). Nuclear staining on rat liver bile duct is observed. Counter stained with Hematoxylin.

Secondary antibody only control: Used PBS instead of primary antibody

Perform heat mediated antigen retrieval with Tris/EDTA buffer pH 9.0 before commencing with IHC staining protocol.




Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-HNF-1B antibody [EPR18644-13] (ab213149)

Immunohistochemical analysis of paraffin-embedded human liver tissue labeling HNF-1B with ab213149 at 1/2000 dilution, followed by Goat Anti-Rabbit IgG H&L (HRP) at 1/500 dilution. Nuclear staining on human liver bile duct is observed. Counter stained with Hematoxylin.

Secondary antibody only control: Used PBS instead of primary antibody, secondary antibody is Goat Anti-Rabbit IgG H&L (HRP) at 1/500 dilution.

Perform heat mediated antigen retrieval with Tris/EDTA buffer pH 9.0 before commencing with IHC staining protocol.




Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-HNF-1B antibody [EPR18644-13] (ab213149)

Immunohistochemical analysis of paraffin-embedded human liver bile duct carcinoma tissue labeling HNF-1B with ab213149 at 1/2000 dilution, followed by Goat Anti-Rabbit IgG H&L (HRP) at 1/500 dilution. Nuclear staining on human liver bile duct carcinoma is observed. Counter stained with Hematoxylin.

Secondary antibody only control: Used PBS instead of primary antibody, secondary antibody is Goat Anti-Rabbit IgG H&L (HRP) at 1/500 dilution.


Perform heat mediated antigen retrieval with Tris/EDTA buffer pH 9.0 before commencing with IHC staining protocol.



Immunohistochemical analysis of paraffin-embedded human hepatocellular carcinoma tissue labeling HNF-1B with ab213149 at 1/2000 dilution, followed by Goat Anti-Rabbit IgG H&L (HRP) at 1/500 dilution. Negative staining on human hepatocellular carcinoma. Counter stained with Hematoxylin.

Perform heat mediated antigen retrieval with Tris/EDTA buffer pH 9.0 before commencing with IHC staining protocol.

Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-HNF-1B antibody [EPR18644-13] (ab213149)




Immunohistochemical analysis of paraffin-embedded human ovarian clear cell carcinoma tissue labeling HNF-1B with ab213149 at 1/2000 dilution, followed by Goat Anti-Rabbit IgG H&L (HRP) at 1/500 dilution. Nuclear staining on human ovarian clear cell carcinoma is observed. Counter stained with Hematoxylin.

Secondary antibody only control: Used PBS instead of primary antibody, secondary antibody is Goat Anti-Rabbit IgG H&L (HRP) at 1/500 dilution.

Perform heat mediated antigen retrieval with Tris/EDTA buffer pH 9.0 before commencing with IHC staining protocol.


Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-HNF-1B antibody [EPR18644-13] (ab213149)



Immunohistochemical analysis of paraffin-embedded human serous ovarian carcinoma tissue labeling HNF-1B with ab213149 at 1/2000 dilution, followed by Goat Anti-Rabbit IgG H&L (HRP) at 1/500 dilution. Negative staining on human serous ovarian carcinoma. Counter stained with Hematoxylin.

Perform heat mediated antigen retrieval with Tris/EDTA buffer pH 9.0 before commencing with IHC staining protocol.

Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-HNF-1B antibody [EPR18644-13] (ab213149)



Immunohistochemical analysis of paraffin-embedded human endometrioid ovary adenocarcinoma tissue labeling HNF-1B with ab213149 at 1/2000 dilution, followed by Goat Anti-Rabbit IgG H&L (HRP) at 1/500 dilution. Negative staining on human endometrioid ovary adenocarcinoma. Counter stained with Hematoxylin.

Perform heat mediated antigen retrieval with Tris/EDTA buffer pH 9.0 before commencing with IHC staining protocol.

Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-HNF-1B antibody [EPR18644-13] (ab213149)

Why choose a recombinant antibody?



Research with confidence
Consistent and reproducible results



Long-term and scalable supply
Recombinant technology



Success from the first experiment
Confirmed specificity



Ethical standards compliant
Animal-free production

Anti-HNF-1B antibody [EPR18644-13] (ab213149)

Please note: All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <https://www.abcam.com/abpromise> or contact our technical team.

Terms and conditions

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors