abcam

Product datasheet

Anti-HRP antibody ab34961

1 References

Overview

Product name Anti-HRP antibody

Description Mouse polyclonal to HRP

Host species Mouse

Tested applications
Suitable for: ELISA, WB
Species reactivity
Reacts with: Horseradish

Immunogen Full length Horseradish peroxidase.

General notesThe Life Science industry has been in the grips of a reproducibility crisis for a number of years.

Abcam is leading the way in addressing this with our range of recombinant monoclonal antibodies and knockout edited cell lines for gold-standard validation. Please check that this product meets

your needs before purchasing.

If you have any questions, special requirements or concerns, please send us an inquiry and/or contact our Support team ahead of purchase. Recommended alternatives for this product can be

found below, along with publications, customer reviews and Q&As

Properties

Form Liquid

Storage instructions Shipped at 4°C. Upon delivery aliquot and store at -20°C or -80°C. Avoid repeated freeze / thaw

cycles.

Storage buffer Preservative: 0.01% Sodium azide

Constituents: 0.87% Sodium chloride, 0.42% Potassium phosphate

Purity Whole antiserum

Purification notes This product was prepared from monospecific antiserum by delipidation and defibrination.

Clonality Polyclonal

Isotype IgG

Applications

The Abpromise guarantee Our Abpromise guarantee covers the use of ab34961 in the following tested applications.

The application notes include recommended starting dilutions; optimal dilutions/concentrations should be determined by the end user.

1

Application	Abreviews	Notes
ELISA		Use at an assay dependent concentration.
WB		Use at an assay dependent concentration.

Target

Relevance	Horseradish Peroxidase (HRP) is an enzyme commonly used as an indicator for chemical
	reactions which produce peroxide. The enzyme is routinely conjugated to antibodies for use in
	enzyme-based immunoassay systems. HRP functions in the removal of H2O2 (hydrogen
	peroxide), oxidation of toxic reductants, biosynthesis and degradation of lignin, suberization, auxin
	catabolism, response to environmental stresses such as wounding, pathogen attack and oxidative
	stress. These functions might be dependent on each isozyme/isoform in each plant tissue.
Cellular localization	Secreted Probable. Vacuole Probable. Note: Carboxy-terminal extension appears to target the protein to vacuoles.

Please note: All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit https://www.abcam.com/abpromise or contact our technical team.

Terms and conditions

· Guarantee only valid for products bought direct from Abcam or one of our authorized distributors