abcam

Product datasheet

HRP Anti-Protein G antibody ab18599

Overview

Product name HRP Anti-Protein G antibody

Description HRP Chicken polyclonal to Protein G

Host species Chicken
Conjugation HRP

Tested applications Suitable for: WB, ELISA

Species reactivity Reacts with: Streptococcus sp. group G

Immunogen Recombinant fragment corresponding to Protein G. Purified recombinant protein G lacking the

albumin binding region.

General notesThe Life Science industry has been in the grips of a reproducibility crisis for a number of years.

Abcam is leading the way in addressing this with our range of recombinant monoclonal antibodies and knockout edited cell lines for gold-standard validation. Please check that this product meets

your needs before purchasing.

If you have any questions, special requirements or concerns, please send us an inquiry and/or contact our Support team ahead of purchase. Recommended alternatives for this product can be

found below, along with publications, customer reviews and Q&As

Properties

Form Liquid

Storage instructions Shipped at 4°C. Store at +4°C.

Storage buffer pH: 7.00

Preservative: 0.01% Thimerosal (merthiolate)

Constituents: 1.19% HEPES, 0.58% Sodium chloride, 0.2% BSA

Purity Immunogen affinity purified

Purification notesAfter multiple immunizations in Freund's adjuvant, eggs were collected and delipidated.

Antibodies were immunoaffinity purified off protein G agarose and conjugated with horseradish

peroxidase (HRP).

Clonality Polyclonal

Isotype IgY

Applications

1

The Abpromise quarantee

Our Abpromise guarantee covers the use of ab18599 in the following tested applications.

The application notes include recommended starting dilutions; optimal dilutions/concentrations should be determined by the end user.

| Application | Abreviews | Notes |
|-------------|-----------|-------------------|
| WB | | 1/1000 - 1/10000. |
| ELISA | | 1/5000 - 1/20000. |

Target

Relevance

Protein G is a bacterial protein derived from the cell wall of certain strains of b-hemolytic Streptococcci. It binds with high affinity to the Fc portion of various classes and subclasses of immunoglobulins from a variety of species. Protein G binds to all IgG subclasses from human, mouse and rat species. It also binds to total IgG from guinea pig, rabbit, goat, cow, sheep, and horse. Protein G binds preferentially to the Fc portion of IgG, but can also bind to the Fab region, making it useful for purification of F(ab') fragments of IgG. Due to it's affinity for the Fc region of many mammalian immunoglobulins, protein G is considered a universal reagent in biochemistry and immunology.

Cellular localization

Cell Wall and Secreted

Please note: All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit https://www.abcam.com/abpromise or contact our technical team.

Terms and conditions

• Guarantee only valid for products bought direct from Abcam or one of our authorized distributors