abcam

Product datasheet

HRP Anti-S6K1 antibody ab9367

2 Images

Overview

Product name HRP Anti-S6K1 antibody

Description HRP Rabbit polyclonal to S6K1

Host species Rabbit
Conjugation HRP

Specificity Reacts strongly with the 70 kDa of p70 S6K but only weakly to the 80 kDa S6K.

Tested applications Suitable for: WB

Species reactivity Reacts with: Mouse

Immunogen Synthetic peptide corresponding to Human S6K1 (C terminal) conjugated to keyhole limpet

haemocyanin.

Run BLAST with EXPASY M Run BLAST with S NCBI

General notesThe Life Science industry has been in the grips of a reproducibility crisis for a number of years.

Abcam is leading the way in addressing this with our range of recombinant monoclonal antibodies and knockout edited cell lines for gold-standard validation. Please check that this product meets

your needs before purchasing.

If you have any questions, special requirements or concerns, please send us an inquiry and/or contact our Support team ahead of purchase. Recommended alternatives for this product can be

found below, along with publications, customer reviews and Q&As

Properties

Form Liquid

Storage instructions Shipped at 4°C. Store at +4°C.

Storage buffer pH: 6

Preservative: 0.02% Sodium azide Constituent: Tris buffered saline

Purity Immunogen affinity purified

Purification notes Immunoaffinity chromatography with S6K c-terminal peptide on agarose.

Clonality Polyclonal

Isotype IgG

1

Applications

The Abpromise guarantee

Our **Abpromise guarantee** covers the use of ab9367 in the following tested applications.

The application notes include recommended starting dilutions; optimal dilutions/concentrations should be determined by the end user.

Application	Abreviews	Notes
WB		1/250.

_			
	2	ra	At.
	а	ıu	CL

Function

Acts to integrate nutrient and growth factor signals in regulation of protein synthesis, cell proliferation, cell growth, cell cycle progression and cell survival. Downstream effector of the mTOR signaling pathway. Phosphorylates specifically ribosomal protein S6 in response to insulin or several classes of mitogens. During translation initiation, the inactive form associatess with the eIF-3 complex under conditions of nutrient depletion. Mitogenic stimulation leads to phosphorylation and dissociation from the eIF-3 complex and the free activated form can phosphorylate other translational targets including EIF4B. Promotes protein synthesis by phosphorylating PDCD4 at 'Ser-67' and targeting it for degradation. Phosphorylates RICTOR leading to regulation of mammalian target of rapamycin complex 2 (mTORC2) signaling; probably phosphorylates RICTOR at 'Thr-1135'. Phosphorylates IRS1 at multiple serine residues coupled with insulin resistance; probably phosphorylates IRS1 at 'Ser-270'. Required for TNF-alpha induced IRS-1 degradation. Phosphorylates EEF2K in response to IGF1 and inhibits EEF2K activity. Phosphorylates BAD at 'Ser-99' in response to IGF1 leading to BAD inactivation and inhibition of BAD-induced apoptosis. Phosphorylates mitochondrial RMP leading to dissociation of a RMP:PPP1CC complex; probably phosphorylates RMP at 'Ser-99'. The free mitochondrial PPP1CC can dephosphorylate RPS6KB1 at Thr-412 which is proposed to be a negative feed back mechanism for the RPS6KB1 antiapoptotic function. Phosphorylates GSK3B at 'Ser-9' under conditions leading to loss of the TSC1-TSC2 complex. Phosphorylates POLDIP3.

Tissue specificity

Widely expressed.

Sequence similarities

Belongs to the protein kinase superfamily. AGC Ser/Thr protein kinase family. S6 kinase

subfamily.

Contains 1 AGC-kinase C-terminal domain.

Contains 1 protein kinase domain.

Domain

The autoinhibitory domain is believed to block phosphorylation within the AGC-kinase C-terminal

domain and the activation loop.

The TOS (TOR signaling) motif is essential for activation by mTORC1.

Post-translational modifications

Phosphorylation at Thr-412 is regulated by mTORC1. The phosphorylation at this site is


maintained by an agonist-dependent autophosphorylation mechanism.

Cellular localization

Cytoplasm; Nucleus. Cytoplasm and Cell junction > synapse > synaptosome. Mitochondrion outer

membrane.

Images




One-step immunoblotting with cross-antibodies of the immunoprecipitated kinases

Immunoprecipitation with anti-p90 RSK1 (Lane A and C)
Immunoprecipitation with anti-p70 S6K (Lane B and D)
Immunoblotting with anti-p70 S6K HRPconjugates (Lane A and B)

Immunoblotting with anti-RSK1 HRP conjugates (Lane C and D)

One-step immunoblotting with cross-antibodies of the immunoprecipitated kinases

Immunoprecipitation with anti-p90 RSK1 (Lane A and C)
Immunoprecipitation with anti-p70 S6K (Lane B and D)
Immunoblotting with anti-p70 S6K HRPconjugates (Lane A and B)
Immunoblotting with anti-RSK1 HRP conjugates (Lane C and D)



Lane A: mouse Muscle

Lane B: mouse Brain

One-step immunoblotting of crude mouse brain and muscle samples with anti-S6K HRP conjugates (ECL system).

One-step immunoblotting of crude mouse brain and muscle samples with anti-S6K HRP conjugates (ECL system).

Lane A: mouse Muscle
Lane B: mouse Brain

western blot - HRP Anti-Sor i antibody (ab9507)

Please note: All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- · Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- · Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit https://www.abcam.com/abpromise or contact our technical team.

Terms and conditions

• Guarantee only valid for products bought direct from Abcam or one of our authorized distributors