abcam

Product datasheet

Anti-ICAM1 antibody [EPR16608] - BSA and Azide free ab271933



3 Images

Overview

Product name Anti-ICAM1 antibody [EPR16608] - BSA and Azide free

Description Rabbit monoclonal [EPR16608] to ICAM1 - BSA and Azide free

Host species Rabbit

Tested applications Suitable for: WB, IHC-P

Species reactivity Reacts with: Mouse

Immunogen Recombinant fragment. This information is proprietary to Abcam and/or its suppliers.

Positive control IHC-P: Mouse liver and kidney tissues.

General notes ab271933 is the carrier-free version of ab179707.

> Our carrier-free antibodies are typically supplied in a PBS-only formulation, purified and free of BSA, sodium azide and glycerol. The carrier-free buffer and high concentration allow for increased conjugation efficiency.

This conjugation-ready format is designed for use with fluorochromes, metal isotopes, oligonucleotides, and enzymes, which makes them ideal for antibody labelling, functional and cellbased assays, flow-based assays (e.g. mass cytometry) and Multiplex Imaging applications.

Use our conjugation kits for antibody conjugates that are ready-to-use in as little as 20 minutes with <1 minute hands-on-time and 100% antibody recovery: available for fluorescent dyes, HRP, biotin and gold.

This product is compatible with the Maxpar® Antibody Labeling Kit from Fluidigm, without the need for antibody preparation. Maxpar[®] is a trademark of Fluidigm Canada Inc.

This product is a recombinant monoclonal antibody, which offers several advantages including:

- High batch-to-batch consistency and reproducibility
- Improved sensitivity and specificity
- Long-term security of supply
- Animal-free production

For more information see here.

Our RabMAb® technology is a patented hybridoma-based technology for making rabbit monoclonal antibodies. For details on our patents, please refer to **RabMAb**® **patents**.

Properties

Form Liquid

Storage instructions Shipped at 4°C. Store at +4°C. Do Not Freeze.

Storage buffer pH: 7.2

Constituent: PBS

Carrier free Yes

Purity Protein A purified

Clonality Monoclonal
Clone number EPR16608

Isotype IgG

Applications

The Abpromise guarantee Our Abpromise guarantee covers the use of ab271933 in the following tested applications.

The application notes include recommended starting dilutions; optimal dilutions/concentrations should be determined by the end user.

Application	Abreviews	Notes
WB		Use at an assay dependent concentration. Predicted molecular weight: 57 kDa.
IHC-P		Use at an assay dependent concentration.

Target

Function ICAM proteins are ligands for the leukocyte adhesion protein LFA-1 (integrin alpha-L/beta-2).

During leukocyte trans-endothelial migration, ICAM1 engagement promotes the assembly of endothelial apical cups through ARHGEF26/SGEF and RHOG activation. In case of rhinovirus

infection acts as a cellular receptor for the virus.

Sequence similaritiesBelongs to the immunoglobulin superfamily. ICAM family.

Contains 5 lg-like C2-type (immunoglobulin-like) domains.


Post-translational

modifications

Monoubiquitinated, which is promoted by MARCH9 and leads to endocytosis.

Cellular localization Membrane.

Images



Immunohistochemistry (Formalin/PFA-fixed paraffinembedded sections) - Anti-ICAM1 antibody

[EPR16608] - BSA and Azide free (ab271933)

Secondary antibody only control Copyright (O) 20 16 Abcam plc

Immunohistochemistry (Formalin/PFA-fixed paraffinembedded sections) - Anti-ICAM1 antibody

[EPR16608] - BSA and Azide free (ab271933)

Immunohistochemical analysis of paraffin-embedded mouse liver tissue labeling ICAM1 with <u>ab179707</u> at 1/2000 dilution, followed by Goat Anti-Rabbit IgG H&L (HRP) (<u>ab97051</u>) at 1/500 dilution.

Positive staining on endothelial cells of mouse liver is observed [PMID: 17058232].

Counter stained with Hematoxylin.

Secondary antibody only control: Used PBS instead of primary antibody, secondary antibody is Goat Anti-Rabbit lgG H&L (HRP) (ab97051) at 1/500 dilution.

Perform heat mediated antigen retrieval with Tris/EDTA buffer pH 9.0 before commencing with IHC staining protocol.

This data was developed using the same antibody clone in a different buffer formulation containing PBS, BSA, glycerol, and sodium azide (<u>ab179707</u>).

Immunohistochemical analysis of paraffin-embedded mouse kidney tissue labeling ICAM1 with <u>ab179707</u> at 1/2000 dilution, followed by Goat Anti-Rabbit IgG H&L (HRP) (ab97051) at 1/500 dilution.


Positive staining on endothelial cells of mouse kidney is observed [PMID: 17058232].

Counter stained with Hematoxylin.

Secondary antibody only control: Used PBS instead of primary antibody, secondary antibody is Goat Anti-Rabbit lgG H&L (HRP) (ab97051) at 1/500 dilution.

Perform heat mediated antigen retrieval with Tris/EDTA buffer pH 9.0 before commencing with IHC staining protocol.

This data was developed using the same antibody clone in a different buffer formulation containing PBS, BSA, glycerol, and sodium azide (<u>ab179707</u>).



Please note: All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit https://www.abcam.com/abpromise or contact our technical team.

Terms and conditions

· Guarantee only valid for products bought direct from Abcam or one of our authorized distributors