# abcam

# Product datasheet

# Anti-IL-2 antibody [EPR16615-341] ab243650

Recombinant RabMAb

★★★★★ 1 Abreviews 1 References 4 Images

#### Overview

**Product name** Anti-IL-2 antibody [EPR16615-341]

**Description** Rabbit monoclonal [EPR16615-341] to IL-2

**Host species** Rabbit

**Tested applications** Suitable for: Flow Cyt (Intra), ICC/IF

Unsuitable for: IHC-P.IP or WB

Species reactivity Reacts with: Mouse

**Immunogen** Recombinant fragment. This information is proprietary to Abcam and/or its suppliers.

Positive control ICC/IF: EL4.IL-2 cells. Flow Cyt (intra):EL4.IL-2 (mouse lymphoma T lymphocyte) treated with

> 80nM PMA + 1.34uM Ionomycin + 10.6uM Brefeldin A + 2uM Monensin for 6 h, Mouse splenocytes treated with 80nM PMA+1.34uM lonomycin+10.6uM Brefeldin A+2uM Monensin.

General notes This product is a recombinant monoclonal antibody, which offers several advantages including:

- High batch-to-batch consistency and reproducibility

- Improved sensitivity and specificity

- Long-term security of supply

- Animal-free production

For more information see here.

Our RabMAb® technology is a patented hybridoma-based technology for making rabbit monoclonal antibodies. For details on our patents, please refer to **RabMAb**® **patents**.

#### **Properties**

**Form** Liquid

Storage instructions Shipped at 4°C. Store at +4°C short term (1-2 weeks). Upon delivery aliquot. Store at -20°C long

term. Avoid freeze / thaw cycle.

Storage buffer Preservative: 0.01% Sodium azide

Constituents: 59.94% PBS, 40% Glycerol (glycerin, glycerine), 0.05% BSA

**Purity** Protein A purified

Clonality Monoclonal Clone number EPR16615-341

**Isotype** IgG

#### **Applications**

The Abpromise guarantee

Our **Abpromise guarantee** covers the use of ab243650 in the following tested applications.

The application notes include recommended starting dilutions; optimal dilutions/concentrations should be determined by the end user.

Application	Abreviews	Notes
Flow Cyt (Intra)		1/50.
ICC/IF	★★☆☆☆ (1)	1/50.

**Application notes** 

Is unsuitable for IHC-P,IP or WB.

#### **Target**

Function Produced by T-cells in response to antigenic or mitogenic stimulation, this protein is required for T-cell proliferation and other activities crucial to regulation of the immune response. Can stimulate B-cells, monocytes, lymphokine-activated killer cells, natural killer cells, and glioma cells.


Involvement in disease Note=A chromosomal aberration involving IL2 is found in a form of T-cell acute lymphoblastic

leukemia (T-ALL). Translocation t(4;16)(q26;p13) with involves TNFRSF17.

**Sequence similarities** Belongs to the IL-2 family.


Cellular localization Secreted.

### **Images**




Immunocytochemistry/ Immunofluorescence - Anti-IL-2 antibody [EPR16615-341] (ab243650) Immunofluorescent analysis of 4% Paraformaldehyde-fixed, 0.1% TritonX-100 permeabilized EL4.IL-2 cells labelling IL-2 with ab243650 at 1/50 (9.1 ug/ml) dilution, followed by **ab150077** Goat Anti-Rabbit lgG H&L (Alexa Fluor® 488) antibody at 1/1000 dilution (Green) Confocal image showing cytoplasmic staining in EL4.IL-2 cell line treated with Cell Stimulation Cocktail (80nM PMA + 1.34uM lonomycin + 10.6uM Brefeldin A + 2uM Monensin) for 6 h. **ab195889** Anti-alpha Tubulin mouse monoclonal antibody - Microtubule Marker (Alexa Fluor® 594) was used to counterstain tubulin at 1/200 dilution (Red). The Nuclear counterstain was DAPI (Blue).

Secondary antibody only control: Secondary antibody is <u>ab150077</u> Goat Anti-Rabbit IgG H&L (Alexa Fluor® 488) at 1/1000 dilution.




Flow Cytometry (Intracellular) - Anti-IL-2 antibody [EPR16615-341] (ab243650)

Intracellular flow cytometric analysis of 2% paraformaldehyde fixed 0.1% Tween-20 permeabilized Mouse splenocytes treated with cell stimulation cocktail (80nM PMA+1.34uM lonomycin+10.6uM Brefeldin A+2uM Monensin) for 6 hours (Right)/ Untreated control (Left). cells labelling IL-2 with ab243650 at 1/500 dilution (1ug)/ Left and Right (Red) compared with a isotype control. A Goat anti rabbit IgG (Alexa Fluor® 647, **ab150079**) at 1/2000 dilution was used as the secondary antibody. Cells were surface stained with anti-CD3 conjugated to FITC. Then fixed with 2% PFA for 10min followed by intracellularly stained with ab243650. Gated on lymphocytes population. Data was kindly provided by an anonymous collaborator.



Flow Cytometry (Intracellular) - Anti-IL-2 antibody [EPR16615-341] (ab243650)

Intracellular flow cytometric analysis of 4% paraformaldehyde fixed 90% methanol permeabilized EL4.IL-2 (mouse lymphoma T lymphocyte) treated with cell stimulation cocktail (80nM PMA + 1.34uM lonomycin + 10.6uM Brefeldin A + 2uM Monensin) for 6 hours (Red)/ Untreated control (Green) cells labelling IL-2 with ab243650 at 1/50 dilution (1ug) (Red) compared with a Rabbit monoclonal IgG (ab172730) (Black) isotype control and an unlabelled control (cells without incubation with primary antibody and secondary antibody) (Blue). A Goat anti rabbit IgG (Alexa Fluor® 488, ab150077) at 1/2000 dilution was used as the secondary antibody.



Please note: All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

## Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <a href="https://www.abcam.com/abpromise">https://www.abcam.com/abpromise</a> or contact our technical team.

#### Terms and conditions

• Guarantee only valid for products bought direct from Abcam or one of our authorized distributors