

Product datasheet

Anti-IRF8 antibody [EPR20441] - BSA and Azide free ab251475

Recombinant RabMAb

6 Images

Overview

Product name	Anti-IRF8 antibody [EPR20441] - BSA and Azide free
Description	Rabbit monoclonal [EPR20441] to IRF8 - BSA and Azide free
Host species	Rabbit
Tested applications	Suitable for: WB, IP, IHC-P
Species reactivity	Reacts with: Human
Immunogen	Recombinant fragment. This information is proprietary to Abcam and/or its suppliers.
General notes	<p>ab251475 is the carrier-free version of ab207418.</p> <p>Our carrier-free antibodies are typically supplied in a PBS-only formulation, purified and free of BSA, sodium azide and glycerol. The carrier-free buffer and high concentration allow for increased conjugation efficiency.</p> <p>This conjugation-ready format is designed for use with fluorochromes, metal isotopes, oligonucleotides, and enzymes, which makes them ideal for antibody labelling, functional and cell-based assays, flow-based assays (e.g. mass cytometry) and Multiplex Imaging applications.</p> <p>Use our conjugation kits for antibody conjugates that are ready-to-use in as little as 20 minutes with <1 minute hands-on-time and 100% antibody recovery: available for fluorescent dyes, HRP, biotin and gold.</p> <p>This product is compatible with the Maxpar[®] Antibody Labeling Kit from Fluidigm, without the need for antibody preparation. Maxpar[®] is a trademark of Fluidigm Canada Inc.</p> <p>This product is a recombinant monoclonal antibody, which offers several advantages including:</p> <ul style="list-style-type: none">- High batch-to-batch consistency and reproducibility- Improved sensitivity and specificity- Long-term security of supply- Animal-free production <p>For more information see here.</p> <p>Our RabMAb[®] technology is a patented hybridoma-based technology for making rabbit monoclonal antibodies. For details on our patents, please refer to RabMAb[®] patents.</p>

Properties

Form	Liquid
Storage instructions	Shipped at 4°C. Store at +4°C. Do Not Freeze.
Storage buffer	pH: 7.2 Constituent: PBS
Carrier free	Yes
Purity	Protein A purified
Clonality	Monoclonal
Clone number	EPR20441
Isotype	IgG

Applications


The Abpromise guarantee Our **Abpromise guarantee** covers the use of ab251475 in the following tested applications. The application notes include recommended starting dilutions; optimal dilutions/concentrations should be determined by the end user.

Application	Abreviews	Notes
WB		Use at an assay dependent concentration. Detects a band of approximately 48 kDa (predicted molecular weight: 48 kDa).
IP		Use at an assay dependent concentration.
IHC-P		Use at an assay dependent concentration. Perform heat mediated antigen retrieval with Tris/EDTA buffer pH 9.0 before commencing with IHC staining protocol.

Target


Function	Specifically binds to the upstream regulatory region of type I IFN and IFN-inducible MHC class I genes (the interferon consensus sequence (ICS)). Plays a negative regulatory role in cells of the immune system.
Tissue specificity	Predominantly in lymphoid tissues.
Sequence similarities	Belongs to the IRF family. Contains 1 IRF tryptophan pentad repeat DNA-binding domain.
Post-translational modifications	Ubiquitinated. Ubiquitination by TRIM21 in macrophages, a process that is strongly increased upon interferon gamma stimulation, leads to the enhanced transcriptional activity of target cytokine genes.
Cellular localization	Nucleus.

Images




Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-IRF8 antibody [EPR20441] - BSA and Azide free (ab251475)

This data was developed using **ab207418**, the same antibody clone in a different buffer formulation. Immunohistochemical analysis of paraffin-embedded human tonsil tissue labeling IRF8 with **ab207418** at 1/1000 dilution, followed by Goat Anti-Rabbit IgG H&L (HRP) Ready to use. Nuclear staining on human tonsil ($\times 200$) [PMID: 18663414] [PMID: 16380510]. Counter stained with Hematoxylin. Secondary antibody only control: Used PBS instead of primary antibody, secondary antibody is Goat Anti-Rabbit IgG H&L (HRP) Ready to use. Perform heat mediated antigen retrieval with Tris/EDTA buffer pH 9.0 before commencing with IHC staining protocol.



Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-IRF8 antibody [EPR20441] - BSA and Azide free (ab251475)

This data was developed using **ab207418**, the same antibody clone in a different buffer formulation. Immunohistochemical analysis of paraffin-embedded human non-hodgkin's lymphoma tissue labeling IRF8 with **ab207418** at 1/1000 dilution, followed by Goat Anti-Rabbit IgG H&L (HRP) ready to use. Nuclear staining on human non-hodgkin's lymphoma. Counter stained with Hematoxylin. Secondary antibody only control: Used PBS instead of primary antibody, secondary antibody is Goat Anti-Rabbit IgG H&L (HRP) ready to use. Perform heat mediated antigen retrieval with Tris/EDTA buffer pH 9.0 before commencing with IHC staining protocol.



Western blot - Anti-IRF8 antibody [EPR20441] - BSA and Azide free (ab251475)

All lanes : Anti-IRF8 antibody [EPR20441] ([ab207418](#)) at 1/1000 dilution

Lane 1 : THP-1 (Human monocytic leukemia cell line) whole cell lysate

Lane 2 : Raji (Human Burkitt's lymphoma cell line) whole cell lysate

Lysates/proteins at 20 µg per lane.

Secondary

All lanes : Goat Anti-Rabbit IgG H&L (HRP) ([ab97051](#)) at 1/100000 dilution


Predicted band size: 48 kDa

Observed band size: 48 kDa

Exposure time: 1 minute

This data was developed using [ab207418](#), the same antibody clone in a different buffer formulation.

Blocking and dilution buffer: 5% NFD/MBST.



All lanes : Anti-IRF8 antibody [EPR20441] ([ab207418](#)) at 1/1000 dilution

Lane 1 : Human fetal spleen lysate

Lane 2 : Human tonsil lysate

Lane 3 : Human colon lysate

Lysates/proteins at 10 µg per lane.

Secondary

All lanes : VeriBlot for IP Detection Reagent (HRP) ([ab131366](#)) at 1/1000 dilution


Predicted band size: 48 kDa

Observed band size: 48 kDa

This data was developed using [ab207418](#), the same antibody clone in a different buffer formulation.





Blocking and dilution buffer: 5% NFDN/TBST.

Exposure time: Lane 1: 30 seconds; Lane 2: 2 seconds; Lane 3: 15 seconds.



This data was developed using **ab207418**, the same antibody clone in a different buffer formulation. IRF8 was immunoprecipitated from 0.35 mg of Raji (Human Burkitt's lymphoma cell line) whole cell lysate with **ab207418** at 1/40 dilution. Western blot was performed from the immunoprecipitate using **ab207418** at 1/500 dilution. VeriBlot for IP Detection Reagent (HRP) (**ab131366**), was used for detection at 1/1000 dilution. Lane 1: Raji whole cell lysate 10 µg (Input). Lane 2: **ab207418** IP in Raji whole cell lysate. Lane 3: Rabbit monoclonal IgG (**ab172730**) instead of **ab207418** in Raji whole cell lysate. Blocking and dilution buffer and concentration: 5% NFDm/TBST. Exposure time: 1 second.

Why choose a recombinant antibody?

 Research with confidence Consistent and reproducible results	 Long-term and scalable supply Recombinant technology
 Success from the first experiment Confirmed specificity	 Ethical standards compliant Animal-free production

Anti-IRF8 antibody [EPR20441] - BSA and Azide free (ab251475)

Please note: All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours

- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <https://www.abcam.com/abpromise> or contact our technical team.

Terms and conditions

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors