abcam

Product datasheet

Anti-Kappa light chain antibody [rKLC709] - BSA and Azide free ab237850

2 Images

Overview

Storage buffer

Purification notes

Clone number

Carrier free

Purity

Clonality

pH: 7.2

Yes

Constituent: PBS

Protein A/G purified

Monoclonal

rKLC709

Product name	Anti-Kappa light chain antibody [rKLC709] - BSA and Azide free		
Description	Mouse monoclonal [rKLC709] to Kappa light chain - BSA and Azide free		
Host species	Mouse		
Tested applications	Suitable for: Protein Array, IHC-P		
Species reactivity	Reacts with: Human		
Immunogen	Recombinant full length protein corresponding to Human Kappa light chain aa 1-150. Database link: P01601		
	Run BLAST with Run BLAST with		
Positive control	IHC-P: Human tonsil tissue.		
General notes	The Life Science industry has been in the grips of a reproducibility crisis for a number of years. Abcam is leading the way in addressing this with our range of recombinant monoclonal antibodies and knockout edited cell lines for gold-standard validation. Please check that this product meets your needs before purchasing.		
	If you have any questions, special requirements or concerns, please send us an inquiry and/or contact our Support team ahead of purchase. Recommended alternatives for this product can be found below, along with publications, customer reviews and Q&As		
Properties			
Form	Liquid		
Storage instructions	Shipped at 4°C. Store at +4°C short term (1-2 weeks). Upon delivery aliquot. Store at -20°C long term. Avoid freeze / thaw cycle.		

Purified from Bioreactor Concentrate by Protein A/G.

lsotype	lgG1
Light chain type	kappa

Applications

The Abpromise guarantee Our <u>Abpromise guarantee</u> covers the use of ab237850 in the following tested applications.


The application notes include recommended starting dilutions; optimal dilutions/concentrations should be determined by the end user.

Application	Abreviews	Notes
Protein Array		Use at an assay dependent concentration.
IHC-P		Use a concentration of 0.5 - 1 μ g/ml. Perform heat mediated antigen retrieval with citrate buffer pH 6 before commencing with IHC staining protocol. Incubate with primary antibody for 30 minutes at RT

Target

Relevance Immunoglobulins belong to a group of related glyco proteins which make up 20% of serum proteins. Antigens and immunoglobulins react to confer immunity to individuals. Immunoglobulins have similar structures of two identical heavy chains and two identical light chains. Both the heavy chains and the light chains are divided into constant and variable regions. The constant regions have the same amino acid sequences between all the immunoglobulin classes. The variable regions have approximately 110 amino acids with high sequence variability. The amino acid sequence of the heavy chain determines the class of an immunoglobulin. The five types of immunoglobulin heavy chains are known as: IgG, IgA, IgM, IgD, and IgE. IgG is divided into four subclasses, and IgA is divided into two subclasses. In serum IgA and IgG are monomers with a single 4 polypeptide unit; while, IgM is a pen tamer. IgA may also form polymers. Kappa light chain antibody can be used for the identification of leukemias, plasmacytomas and certain non Hodgkin's lymphomas. Kappa light chain contains one immunoglobulin like domain. The EU sequence has the INV allotypic marker, Ala 45 and Val 83. The ROY sequence has the INV allotypic marker, Ala 45 and Leu 83. **Cellular localization** Cytoplasmic and Secreted

Images




Protein Array - Anti-Kappa light chain antibody [rKLC709] - BSA and Azide free (ab237850) This data was produced with **<u>ab238006</u>**, the same antibody in a different formulation with BSA and Azide.

<u>ab238006</u> was tested in protein array against over 19000 different full-length human proteins.

Z- and S- Score: The Z-score represents the strength of a signal that a monoclonal antibody (MAb) (in combination with a fluorescently-tagged anti-IgG secondary antibody) produces when binding to a particular protein on the HuProtTM array. Z-scores are described in units of standard deviations (SD's) above the mean value of all signals generated on that array. If targets on HuProtTM are arranged in descending order of the Z-score, the S-score is the difference (also in units of SD's) between the Z-score. S-score therefore represents the relative target specificity of a MAb to its intended target.

A MAb is specific to its intended target if the MAb has an S-score of at least 2.5. For example, if a MAb binds to protein X with a Z-score of 43 and to protein Y with a Z-score of 14, then the S-score for the binding of that MAb to protein X is equal to 29.

Formalin-fixed, paraffin-embedded human tonsil tissue stained for Kappa light chain using ab237850 at 1 μ g/ml in immunohistochemical analysis.



Immunohistochemistry (Formalin/PFA-fixed paraffinembedded sections) - Anti-Kappa light chain antibody [rKLC709] - BSA and Azide free (ab237850)

Our Abpromise to you: Quality guaranteed and expert technical support

- · Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- · We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <u>https://www.abcam.com/abpromise</u> or contact our technical team.

Terms and conditions

• Guarantee only valid for products bought direct from Abcam or one of our authorized distributors