

Anti-L1CAM antibody [EPR18750] - Low endotoxin, Azide free ab213611

KO VALIDATED

Recombinant

RabMAb

13 Images

Overview

Product name	Anti-L1CAM antibody [EPR18750] - Low endotoxin, Azide free
Description	Rabbit monoclonal [EPR18750] to L1CAM - Low endotoxin, Azide free
Host species	Rabbit
Tested applications	Suitable for: IHC-P, WB, IP
Species reactivity	Reacts with: Mouse, Rat, Human
Immunogen	Recombinant fragment. This information is proprietary to Abcam and/or its suppliers.
Positive control	WB: Human fetal brain and cerebellum lysates; HeLa and A-375 whole cell lysates; Rat brain, cerebellum and hippocampus lysates. Mouse cerebellum and brain lysates. IHC-P: Human kidney, Human stomach cancer, mouse cerebrum, mouse colon, rat cerebellum and rat colon tissues. IP: Human cerebellum lysate; Rat brain whole lysate.
General notes	<p>ab213611 is the carrier-free version of ab208155.</p> <p>Our carrier-free antibodies are typically supplied in a PBS-only formulation, purified and free of BSA, sodium azide and glycerol. The carrier-free buffer and high concentration allow for increased conjugation efficiency.</p> <p>This conjugation-ready format is designed for use with fluorochromes, metal isotopes, oligonucleotides, and enzymes, which makes them ideal for antibody labelling, functional and cell-based assays, flow-based assays (e.g. mass cytometry) and Multiplex Imaging applications.</p> <p>Use our conjugation kits for antibody conjugates that are ready-to-use in as little as 20 minutes with <1 minute hands-on-time and 100% antibody recovery: available for fluorescent dyes, HRP, biotin and gold.</p> <p>This product is compatible with the Maxpar[®] Antibody Labeling Kit from Fluidigm, without the need for antibody preparation. Maxpar[®] is a trademark of Fluidigm Canada Inc.</p> <p>This product is a recombinant monoclonal antibody, which offers several advantages including:</p> <ul style="list-style-type: none"> - High batch-to-batch consistency and reproducibility - Improved sensitivity and specificity - Long-term security of supply - Animal-free production <p>For more information see here.</p> <p>Our RabMAb[®] technology is a patented hybridoma-based technology for making rabbit</p>

monoclonal antibodies. For details on our patents, please refer to [**RabMAb® patents**](#).

Our **Low endotoxin, azide-free formats** have low endotoxin level (≤ 1 EU/ml, determined by the LAL assay) and are free from azide, to achieve consistent experimental results in functional assays.

Properties

Form	Liquid
Storage instructions	Shipped at 4°C. Store at +4°C. Do Not Freeze.
Storage buffer	pH: 7.2 Constituent: PBS
Carrier free	Yes
Purity	Protein A purified
Clonality	Monoclonal
Clone number	EPR18750
Isotype	IgG

Applications

The Abpromise guarantee Our **Abpromise guarantee** covers the use of ab213611 in the following tested applications. The application notes include recommended starting dilutions; optimal dilutions/concentrations should be determined by the end user.

Application	Abreviews	Notes
IHC-P		Use at an assay dependent concentration. Perform heat mediated antigen retrieval with Tris/EDTA buffer pH 9.0 before commencing with IHC staining protocol.
WB		Use at an assay dependent concentration. Predicted molecular weight: 140 kDa.
IP		Use at an assay dependent concentration.

Target

Function	Cell adhesion molecule with an important role in the development of the nervous system. Involved in neuron-neuron adhesion, neurite fasciculation, outgrowth of neurites, etc. Binds to axonin on neurons.
Involvement in disease	Defects in L1CAM are the cause of hydrocephalus due to stenosis of the aqueduct of Sylvius (HSAS) [MIM:307000]. Hydrocephalus is a condition in which abnormal accumulation of cerebrospinal fluid in the brain causes increased intracranial pressure inside the skull. This is usually due to blockage of cerebrospinal fluid outflow in the brain ventricles or in the subarachnoid space at the base of the brain. In children is typically characterized by enlargement of the head, prominence of the forehead, brain atrophy, mental deterioration, and convulsions. In adults the syndrome includes incontinence, imbalance, and dementia. HSAS is characterized by mental

retardation and enlarged brain ventricles.

Defects in L1CAM are the cause of mental retardation-aphasia-shuffling gait-adducted thumbs syndrome (MASA) [MIM:303350]; also known as corpus callosum hypoplasia, psychomotor retardation, adducted thumbs, spastic paraparesis, and hydrocephalus or CRASH syndrome. MASA is an X-linked recessive syndrome with a highly variable clinical spectrum. Main clinical features include spasticity and hyperreflexia of lower limbs, shuffling gait, mental retardation, aphasia and adducted thumbs. The features of spasticity have been referred to as complicated spastic paraplegia type 1 (SPG1). Some patients manifest corpus callosum hypoplasia and hydrocephalus. Inter- and intrafamilial variability is very wide, such that patients with hydrocephalus, MASA, SPG1, and agenesis of corpus callosum can be present within the same family.

Defects in L1CAM are the cause of spastic paraplegia X-linked type 1 (SPG1) [MIM:303350]. Spastic paraplegia is a degenerative spinal cord disorder characterized by a slow, gradual, progressive weakness and spasticity of the lower limbs.

Note=Defects in L1CAM may contribute to Hirschsprung disease by modifying the effects of Hirschsprung disease-associated genes to cause intestinal aganglionosis.

Defects in L1CAM are a cause of partial agenesis of the corpus callosum (ACCPX) [MIM:304100]. A syndrome characterized by partial corpus callosum agenesis, hypoplasia of inferior vermis and cerebellum, mental retardation, seizures and spasticity. Other features include microcephaly, unusual facies, and Hirschsprung disease in some patients.

Sequence similarities

Belongs to the immunoglobulin superfamily. L1/neurofascin/NgCAM family.


Contains 5 fibronectin type-III domains.

Contains 6 Ig-like C2-type (immunoglobulin-like) domains.

Cellular localization

Cell membrane.

Images



All lanes : Anti-L1CAM antibody [EPR18750] ([ab208155](#)) at 1/1000 dilution

Lane 1 : Wild-type HeLa cell lysate

Lane 2 : L1CAM knockout HeLa cell lysate

Lysates/proteins at 20 µg per lane.

Performed under reducing conditions.

Predicted band size: 140 kDa


Observed band size: 220 kDa

This data was developed using the same antibody clone in a different buffer formulation ([ab208155](#)).

Lanes 1 - 2: Merged signal (red and green). Green - [ab208155](#) observed at 220 kDa. Red - loading control, [ab130007](#) observed at

125 kDa.

ab208155 was shown to react with L1CAM in wild-type HeLa. Loss of signal was observed when knockout cell line **ab255401** (knockout cell lysate **ab263786**) was used. Wild-type and L1CAM knockout samples were subjected to SDS-PAGE. **ab208155** and Anti-Vinculin antibody [VIN-54] (**ab130007**) were incubated overnight at 4°C at 1 in 1000 dilution and 1 in 20000 dilution respectively. Blots were developed with Goat anti-Rabbit IgG H&L (IRDye® 800CW) preadsorbed (**ab216773**) and Goat anti-Mouse IgG H&L (IRDye® 680RD) preadsorbed (**ab216776**) secondary antibodies at 1 in 20000 dilution for 1 hour at room temperature before imaging.



Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-L1CAM antibody [EPR18750] - Low endotoxin, Azide free (ab213611)

Immunohistochemical analysis of paraffin-embedded Human kidney tissue labeling L1CAM with **ab208155** at 1/500 dilution, followed by Goat Anti-Rabbit IgG H&L (HRP) (**ab97051**) at 1/500 dilution.


Membrane staining on a part of Human kidney tubules is observed. L1CAM specific staining most abundant on nervous system, distal kidney tubules, and tumor cells. [PMID: 16867862, PMID: 20044598].

Counter stained with Hematoxylin.

Secondary antibody only control: Used PBS instead of primary antibody, secondary antibody is Goat Anti-Rabbit IgG H&L (HRP) (**ab97051**) at 1/500 dilution.

This data was developed using the same antibody clone in a different buffer formulation containing PBS, BSA, glycerol, and sodium azide (**ab208155**).

Heat mediated antigen retrieval was performed with Tris/EDTA buffer pH 9.0 before commencing with IHC staining protocol.



Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-L1CAM antibody [EPR18750] - Low endotoxin, Azide free (ab213611)

Immunohistochemical analysis of paraffin-embedded Human liver tissue labeling L1CAM with **ab208155** at 1/500 dilution, followed by Goat Anti-Rabbit IgG H&L (HRP) (**ab97051**) at 1/500 dilution.


Negative staining on the Human liver.

L1CAM specific staining most abundant on nervous system, distal kidney tubules, and tumor cells. [PMID: 16867862, PMID: 20044598].

Counter stained with Hematoxylin.

This data was developed using the same antibody clone in a different buffer formulation containing PBS, BSA, glycerol, and sodium azide (**ab208155**).

Heat mediated antigen retrieval was performed with Tris/EDTA buffer pH 9.0 before commencing with IHC staining protocol.



Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-L1CAM antibody [EPR18750] - Low endotoxin, Azide free (ab213611)

Immunohistochemical analysis of paraffin-embedded Mouse colon tissue labeling L1CAM with **ab208155** at 1/500 dilution, followed by Goat Anti-Rabbit IgG H&L (HRP) (**ab97051**) at 1/500 dilution.

Mainly membrane staining on the nerve tract of mouse colon is observed.


L1CAM specific staining most abundant on nervous system, distal kidney tubules, and tumor cells. [PMID: 16867862, PMID: 20044598].

Counter stained with Hematoxylin.

Secondary antibody only control: Used PBS instead of primary antibody, secondary antibody is Goat Anti-Rabbit IgG H&L (HRP) (**ab97051**) at 1/500 dilution.

This data was developed using the same antibody clone in a different buffer formulation containing PBS, BSA, glycerol, and sodium azide (**ab208155**).

Heat mediated antigen retrieval was performed with Tris/EDTA buffer pH 9.0 before commencing with IHC staining protocol.



Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-L1CAM antibody [EPR18750] - Low endotoxin, Azide free (ab213611)

Immunohistochemical analysis of paraffin-embedded Mouse testis tissue labeling L1CAM with **ab208155** at 1/500 dilution, followed by Goat Anti-Rabbit IgG H&L (HRP) (**ab97051**) at 1/500 dilution.


Negative staining on the mouse testis.

L1CAM specific staining most abundant on nervous system, distal kidney tubules, and tumor cells. [PMID: 16867862, PMID: 20044598].

Counter stained with Hematoxylin.

This data was developed using the same antibody clone in a different buffer formulation containing PBS, BSA, glycerol, and sodium azide (**ab208155**).

Heat mediated antigen retrieval was performed with Tris/EDTA buffer pH 9.0 before commencing with IHC staining protocol.



Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-L1CAM antibody [EPR18750] - Low endotoxin, Azide free (ab213611)

Immunohistochemical analysis of paraffin-embedded Rat cerebellum tissue labeling L1CAM with **ab208155** at 1/500 dilution, followed by Goat Anti-Rabbit IgG H&L (HRP) (**ab97051**) at 1/500 dilution.

Cytoplasm staining on the molecular layer of the rat cerebellar cortex is observed.


L1CAM specific staining most abundant on nervous system, distal kidney tubules, and tumor cells. [PMID: 16867862, PMID: 20044598].

Counter stained with Hematoxylin.

Secondary antibody only control: Used PBS instead of primary antibody, secondary antibody is Goat Anti-Rabbit IgG H&L (HRP) (**ab97051**) at 1/500 dilution.

This data was developed using the same antibody clone in a different buffer formulation containing PBS, BSA, glycerol, and sodium azide (**ab208155**).

Heat mediated antigen retrieval was performed with Tris/EDTA buffer pH 9.0 before commencing with IHC staining protocol.



Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-L1CAM antibody [EPR18750] - Low endotoxin, Azide free (ab213611)

Immunohistochemical analysis of paraffin-embedded Rat colon tissue labeling L1CAM with **ab208155** at 1/500 dilution, followed by Goat Anti-Rabbit IgG H&L (HRP) (**ab97051**) at 1/500 dilution.


Cytoplasm staining on the nerve tract of the rat colon is observed. L1CAM specific staining most abundant on nervous system, distal kidney tubules, and tumor cells. [PMID: 16867862, PMID: 20044598].

Counter stained with Hematoxylin.

Secondary antibody only control: Used PBS instead of primary antibody, secondary antibody is Goat Anti-Rabbit IgG H&L (HRP) (**ab97051**) at 1/500 dilution.

This data was developed using the same antibody clone in a different buffer formulation containing PBS, BSA, glycerol, and sodium azide (**ab208155**).

Heat mediated antigen retrieval was performed with Tris/EDTA buffer pH 9.0 before commencing with IHC staining protocol.



Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-L1CAM antibody [EPR18750] - Low endotoxin, Azide free (ab213611)

Immunohistochemical analysis of paraffin-embedded Rat skeletal muscle tissue labeling L1CAM with **ab208155** at 1/500 dilution, followed by Goat Anti-Rabbit IgG H&L (HRP) (**ab97051**) at 1/500 dilution.

Negative staining on the rat skeletal muscle.


L1CAM specific staining most abundant on nervous system, distal kidney tubules, and tumor cells. [PMID: 16867862, PMID: 20044598].

Counter stained with Hematoxylin.

Secondary antibody only control: Used PBS instead of primary antibody, secondary antibody is **ab97051** at 1/500 dilution.

This data was developed using the same antibody clone in a different buffer formulation containing PBS, BSA, glycerol, and sodium azide (**ab208155**).

Heat mediated antigen retrieval was performed with Tris/EDTA buffer pH 9.0 before commencing with IHC staining protocol.



Immunoprecipitation - Anti-L1CAM antibody
[EPR18750] - Low endotoxin, Azide free (ab213611)

L1CAM was immunoprecipitated from 1mg of Human cerebellum lysate with **ab208155** at 1/40 dilution. Western blot was performed from the immunoprecipitate using **ab208155** at 1/1000 dilution. VeriBlot for IP Detection Reagent (HRP) (**ab131366**), was used for detection at 1/10,000 dilution.

Lane 1: Human cerebellum lysate 10µg (Input).


Lane 2: **ab208155** IP in Human cerebellum lysate.

Lane 3: Rabbit monoclonal IgG (**ab172730**) instead of **ab208155** in Human cerebellum lysate.

Blocking and dilution buffer and concentration: 5% NFDM/TBST.

Exposure time: 3 minutes.

This data was developed using the same antibody clone in a different buffer formulation containing PBS, BSA, glycerol, and sodium azide (**ab208155**).



Immunoprecipitation - Anti-L1CAM antibody
[EPR18750] - Low endotoxin, Azide free (ab213611)

L1CAM was immunoprecipitated from 1mg of Rat brain whole cell lysate with **ab208155** at 1/40 dilution. Western blot was performed from the immunoprecipitate using **ab208155** at 1/1000 dilution. VeriBlot for IP Detection Reagent (HRP) (**ab131366**), was used for detection at 1/10,000 dilution.

Lane 1: Rat brain whole cell lysate 10µg (Input).


Lane 2: **ab208155** IP in Rat brain whole cell lysate.

Lane 3: Rabbit monoclonal IgG (**ab172730**) instead of **ab208155** in Rat brain whole cell lysate.

Blocking and dilution buffer and concentration: 5% NFDM/TBST.

Exposure time: 3 minutes.

This data was developed using the same antibody clone in a different buffer formulation containing PBS, BSA, glycerol, and sodium azide (**ab208155**).



Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-L1CAM antibody [EPR18750] - Low endotoxin, Azide free (ab213611)

This IHC data was generated using the same anti-L1CAM antibody clone, EPR18750, in a different buffer formulation (cat# [ab208155](#)).


Immunohistochemical analysis of paraffin-embedded Mouse cerebrum tissue labeling L1CAM with [ab208155](#) at 1/500 dilution, followed by Goat Anti-Rabbit IgG H&L (HRP) ([ab97051](#)) at 1/500 dilution.

Cytoplasm staining on the mouse cerebrum is observed. L1CAM specific staining most abundant on nervous system, distal kidney tubules, and tumor cells. [PMID: 16867862, PMID: 20044598].

Counter stained with Hematoxylin.

Secondary antibody only control: Used PBS instead of primary antibody, secondary antibody is Goat Anti-Rabbit IgG H&L (HRP) ([ab97051](#)) at 1/500 dilution.

Heat mediated antigen retrieval was performed with Tris/EDTA buffer pH 9.0 before commencing with IHC staining protocol.



Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-L1CAM antibody [EPR18750] - Low endotoxin, Azide free (ab213611)

This IHC data was generated using the same anti-L1CAM antibody clone, EPR18750, in a different buffer formulation (cat# [ab208155](#)).

Immunohistochemical analysis of paraffin-embedded Human stomach cancer tissue labeling L1CAM with [ab208155](#) at 1/500 dilution, followed by Goat Anti-Rabbit IgG H&L (HRP) ([ab97051](#)) at 1/500 dilution.

Membrane staining on the tumor cells of Human stomach cancer is observed.

L1CAM specific staining most abundant on nervous system, distal kidney tubules, and tumor cells. [PMID: 16867862, PMID: 20044598].

Counter stained with Hematoxylin.

Secondary antibody only control: Used PBS instead of primary antibody, secondary antibody is Goat Anti-Rabbit IgG H&L (HRP) ([ab97051](#)) at 1/500 dilution.

Heat mediated antigen retrieval was performed with Tris/EDTA buffer pH 9.0 before commencing with IHC staining protocol.

Why choose a recombinant antibody?



Research with confidence
Consistent and reproducible results



Long-term and scalable supply
Recombinant technology



Success from the first experiment
Confirmed specificity



Ethical standards compliant
Animal-free production

Anti-L1CAM antibody [EPR18750] - Low endotoxin,
Azide free (ab213611)

Please note: All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <https://www.abcam.com/abpromise> or contact our technical team.

Terms and conditions

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors