

Product datasheet

Anti-LAG-3 antibody [EPR20261] α b209236

Recombinant RabMAb

★★★★★ 1 Abreviews 10 References 12 Images

Overview

| | |
|---------------------|---|
| Product name | Anti-LAG-3 antibody [EPR20261] |
| Description | Rabbit monoclonal [EPR20261] to LAG-3 |
| Host species | Rabbit |
| Tested applications | Suitable for: WB, IHC-P, ICC/IF, Flow Cyt, IP |
| Species reactivity | Reacts with: Human |
| Immunogen | Recombinant fragment. This information is proprietary to Abcam and/or its suppliers. |
| Positive control | WB: HDLM-2 cells and Human LAG3 Fc chimera recombinant protein (aa23-450). 293T transfected with LAG3 expression vector containing a GFP tag, whole cell lysate. IHC-P: Human tonsil and Hodgkin's lymphoma tissues. ICC/IF: HEK-293T cells transfected with a GFP-tagged LAG3 expression construct. Flow Cyt: HEK-293T transfected with a GFP-tagged human LAG3 construct, Human PBMCs. IP: HEK-293T transfected with a GFP-tagged human LAG3 construct whole cell lysate. |
| General notes | <p>This product is a recombinant monoclonal antibody, which offers several advantages including:</p> <ul style="list-style-type: none"> - High batch-to-batch consistency and reproducibility - Improved sensitivity and specificity - Long-term security of supply - Animal-free production <p>For more information see here.</p> <p>Our RabMAb[®] technology is a patented hybridoma-based technology for making rabbit monoclonal antibodies. For details on our patents, please refer to RabMAb[®] patents.</p> |

Properties

| | |
|----------------------|---|
| Form | Liquid |
| Storage instructions | Shipped at 4°C. Store at +4°C short term (1-2 weeks). Upon delivery aliquot. Store at -20°C long term. Avoid freeze / thaw cycle. |
| Storage buffer | <p>pH: 7.2</p> <p>Preservative: 0.01% Sodium azide</p> <p>Constituents: PBS, 40% Glycerol, 0.05% BSA</p> |
| Purity | Protein A purified |

| | |
|---------------------|------------|
| Clonality | Monoclonal |
| Clone number | EPR20261 |
| Isotype | IgG |

Applications


The Abpromise guarantee Our **Abpromise guarantee** covers the use of ab209236 in the following tested applications. The application notes include recommended starting dilutions; optimal dilutions/concentrations should be determined by the end user.

| Application | Abreviews | Notes |
|-----------------|-----------|---|
| WB | | 1/500. Detects a band of approximately 70 kDa (predicted molecular weight: 54 kDa). |
| IHC-P | ★★★★★ (1) | 1/1000. Perform heat mediated antigen retrieval with Tris/EDTA buffer pH 9.0 before commencing with IHC staining protocol. Different mRNA expression levels of LAG3 in brain have been reported in the literature (PMID: 1692078; PMID: 12825348). In IHC, under our experimental conditions, this antibody showed no positive staining on human cerebral cortex. |
| ICC/IF | | 1/100. |
| Flow Cyt | | 1/500. |
| IP | | 1/30. |

Target

| | |
|------------------------------|--|
| Function | Involved in lymphocyte activation. Binds to HLA class-II antigens. |
| Tissue specificity | On cell surface of activated NK and T-lymphocytes. |
| Sequence similarities | Contains 3 Ig-like C2-type (immunoglobulin-like) domains. Contains 1 Ig-like V-type (immunoglobulin-like) domain. |
| Cellular localization | Membrane. |

Images




Flow Cytometry - Anti-LAG-3 antibody [EPR20261] (ab209236)

Flow cytometry staining of human peripheral blood mononuclear cells (PBMCs) (top) or PBMCs treated with 5 µg/ml phytohaemagglutinin (PHA) for 72 hours (bottom), with ab209236 (right) or Recombinant Rabbit IgG, monoclonal [EPR25A] - Isotype Control (left). PBMCs were incubated for 30 min on ice in 1x PBS containing 10 µg/ml human IgG and 10 % normal goat serum to block FC receptors and non-specific protein-protein interaction followed by the antibody ab209236 or Recombinant Rabbit IgG, monoclonal [EPR25A] - Isotype Control (1×10^6 in 100 µl at 5.0 µg/ml (1/402)) for 30min on ice. The cells were simultaneously stained with CD3.

The secondary antibody Goat Anti-Rabbit IgG H&L (Alexa Fluor® 488) preadsorbed was incubated at 1/4000 for 30min on ice

Acquisition of >30000 events were collected using a 50 mW Blue laser (488nm) and 525/40 bandpass filter. Events were gated on viable cells.



Western blot - Anti-LAG-3 antibody [EPR20261] (ab209236)

All lanes : Anti-LAG-3 antibody [EPR20261] (ab209236) at 1/5000 dilution

Lane 1 : 293T (Human embryonic kidney epithelial cell) transfected with an empty vector (vector control), containing a GFP tag, whole cell lysate

Lane 2 : 293T transfected with LAG3 expression vector containing a GFP tag, whole cell lysate

Lysates/proteins at 15 µg per lane.


Secondary

All lanes : Goat Anti-Rabbit IgG H&L (HRP) (**ab97051**) at 1/100000 dilution

Predicted band size: 54 kDa

Exposure time: 5 seconds

Blocking/Diluting buffer: 5% NFDM/TBST.



All lanes : Anti-LAG-3 antibody [EPR20261] (ab209236) at 1/500 dilution

Lane 1 : HDLM-2 (Human Hodgkin lymphoma) whole cell lysate

Lane 2 : Jurkat (Human T cell leukemia T lymphocyte) whole cell lysate

Lysates/proteins at 20 µg per lane.

Secondary

All lanes : Goat anti-Rabbit IgG H&L (IRDye® 800CW) preadsorbed (**ab216773**) at 1/10000 dilution (Goat anti-Rabbit IgG H&L (IRDye® 800RCW) preadsorbed)

Predicted band size: 54 kDa

Primary loading control and concentration: Anti-GAPDH antibody [6C5] - Loading Control (**ab8245**) at 1/20000 dilution

Secondary loading control and concentration: Goat anti-Mouse IgG H&L (IRDye® 680RD) preadsorbed (**ab216776**) at 1/10000 dilution


Lanes 1-2: Merged signal (red and green). Green – ab209236 observed at 54-70 kDa. Red - loading control **ab8245** observed at 36 kDa.

ab209236 and Anti-GAPDH antibody [6C5] - Loading Control (**ab8245**) were incubated overnight at 4°C at 1 in 500 dilution and 1 in 20000 dilution respectively. Blots were developed with Goat anti-Rabbit IgG H&L (IRDye® 800RCW) preadsorbed (**ab216773**) and Goat anti-Mouse IgG H&L (IRDye® 680RD) preadsorbed (**ab216776**) secondary antibodies at 1 in 10000 dilution for 1 hour at room temperature before imaging.

The expression profile observed in Jurkat is consistent with the literature (PMID: 25108024).

Negative control: Jurkat (PMID: 25108024)

Observed MW: 54-70 kDa




Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-LAG-3 antibody [EPR20261] (ab209236)

Immunohistochemical analysis of paraffin-embedded human tonsil Hodgkin's lymphoma labeling LAG-3 with ab209236 at 1/1000 dilution, followed by Goat Anti-Rabbit IgG H&L (HRP) Ready to use. Cytoplasmic staining on immunocytes of the human Hodgkin's lymphoma [PMID: 11527700; PMID: 16757686].

Counter stained with Hematoxylin.

Secondary antibody only control: Used PBS instead of primary antibody, secondary antibody is Goat Anti-Rabbit IgG H&L (HRP) Ready to use.

Perform heat mediated antigen retrieval with Tris/EDTA buffer pH 9.0 before commencing with IHC staining protocol.




Immunocytochemistry/ Immunofluorescence - Anti-LAG-3 antibody [EPR20261] (ab209236)

Immunofluorescent analysis of 4% paraformaldehyde-fixed, 0.1% Triton X-100 permeabilized HEK-293T (Human epithelial cell line from embryonic kidney) cells transfected with GFP-tagged LAG3 expression construct or GFP only, labeling LAG-3 with ab209236 at 1/100 dilution, followed by Goat anti-rabbit IgG (Alexa Fluor® 647) ([ab150079](#)) secondary antibody at 1/1000 dilution (green).

Confocal image showing positive staining on HEK-293T cells transfected with a GFP-tagged LAG3 expression construct.

The nuclear counter stain is DAPI (blue). Tubulin is detected with [ab195889](#) (Anti-alpha Tubulin antibody [DM1A] - Microtubule Marker (Alexa Fluor® 594)) at 1/200 dilution (red).

Secondary antibody only control: Used PBS instead of primary antibody, secondary antibody is Goat anti-rabbit IgG (Alexa Fluor® 647) ([ab150079](#)) at 1/1000 dilution.




Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-LAG-3 antibody [EPR20261] (ab209236)

Immunohistochemical analysis of paraffin-embedded human tonsil tissue labeling LAG-3 with ab209236 at 1/1000 dilution, followed by Goat Anti-Rabbit IgG H&L (HRP) Ready to use. Sporadic cytoplasmic staining on immunocytes of human tonsil [PMID: 11527700; PMID: 16757686].

Counter stained with Hematoxylin.

Secondary antibody only control: Used PBS instead of primary antibody, secondary antibody is Goat Anti-Rabbit IgG H&L (HRP) Ready to use.


Perform heat mediated antigen retrieval with Tris/EDTA buffer pH 9.0 before commencing with IHC staining protocol.



Flow Cytometry - Anti-LAG-3 antibody [EPR20261] (ab209236)

Flow cytometric analysis of Human peripheral blood mononuclear cells treated with 1 µg/mL PHA for 3 days cells with ab209236 at 1/50 dilution (right) compared with a rabbit monoclonal IgG isotype control (**ab172730**; left). **ab150077** at 1/2000 dilution was used as the secondary antibody.


Only the CD3+ population are also positive for LAG3. Gated on total viable cells.



Flow Cytometry - Anti-LAG-3 antibody [EPR20261] (ab209236)

Flow cytometric analysis of HEK-293T (Human epithelial cell line from embryonic kidney) cells transfected with a GFP-tagged human LAG-3 construct labeling LAG-3 with ab209236 at 1/500 dilution (right) compared with a rabbit monoclonal IgG isotype control (**ab172730**; left). Goat anti rabbit IgG (Alexa Fluor® 647) **ab150079** at 1/2000 dilution was used as the secondary antibody.

Note: Fresh cells without fixation and permeabilization were used to perform FC testing. Only GFP positive population results in LAG3 positive staining (Q2, right panel).



Western blot - Anti-LAG-3 antibody [EPR20261]
(ab209236)

Anti-LAG-3 antibody [EPR20261] (ab209236) at 1/1000 dilution +
Human LAG3 Fc chimera recombinant protein (aa23-450) at 0.01
µg

Secondary


Goat Anti-Rabbit IgG H&L (HRP) ([ab97051](#)) at 1/100000 dilution

Predicted band size: 54 kDa

Observed band size: 90 kDa

Exposure time: 1 second

Blocking/Dilution buffer: 5% NFDM/TBST.



Immunoprecipitation - Anti-LAG-3 antibody
[EPR20261] (ab209236)

LAG-3 was immunoprecipitated from 0.35 mg of HEK-293T
(Human epithelial cell line from embryonic kidney) transfected with a
GFP-tagged human LAG3 construct whole cell lysate with
ab209236 at 1/30 dilution. Western blot was performed from the
immunoprecipitate using ab209236 at 1/1000 dilution. VeriBlot for
IP Detection Reagent (HRP) ([ab131366](#)), was used for detection at
1/10000 dilution.

Lane 1: HEK-293T transfected with a GFP-tagged human LAG3
construct whole cell lysate 10 µg (Input).

Lane 2: ab209236 IP in HEK-293T transfected with a GFP-tagged
human LAG3 construct whole cell lysate.

Lane 3: Rabbit monoclonal IgG ([ab172730](#)) instead of ab209236 in
HEK-293T transfected with a GFP-tagged human LAG3 construct
whole cell lysate.

Blocking and dilution buffer and concentration: 5% NFDM/TBST.

Exposure time: 1 second.

| Tissue Microarray (TMA) data for ab209236 | | | | | |
|---|---|-----------------------|--------------------------|--------------------------------------|---|
| Normal tissue samples | | | Malignant tissue samples | | |
| Human cardiac muscle | x | Human placenta | x | Clear cell carcinoma of human kidney | x |
| Human cerebrum | x | Human skeletal muscle | x | Human bladder cancer | x |
| Human colon | x | Human skin | x | Human breast carcinoma | x |
| Human endometrium | x | Human spleen | x | Human cervical carcinoma | x |
| Human kidney | x | Human stomach | x (immune cells) | Human colon carcinoma | x |
| Human liver | x | Human testis | x | Human endometrial carcinoma | x |
| Human lung | x | Human thyroid | x | Human gastric carcinoma | x |
| Human mammary gland | x | Human tonsil | ✓ | Human glioma | x |
| Human pancreas | x | | | | |

Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-LAG-3 antibody [EPR20261] (ab209236)

Tissue Microarrays stained for "Anti-LAG-3 antibody [EPR20261]" using "ab209236" in immunohistochemical analysis. This table provides a detailed overview of positive (tick mark) and negative (cross mark) staining per sample type tested. The sections were pre-treated using Heat mediated antigen retrieval using [ab93684](#) (Tris/EDTA buffer, pH 9.0). The sections were incubated with ab209236 at +4°C overnight followed by a ready to use Goat Anti-Rabbit IgG H&L (HRP polymer).

Why choose a recombinant antibody?

Research with confidence
Consistent and reproducible results

Long-term and scalable supply
Recombinant technology

Success from the first experiment
Confirmed specificity

Ethical standards compliant
Animal-free production

Anti-LAG-3 antibody [EPR20261] (ab209236)

Please note: All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery

8

- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <https://www.abcam.com/abpromise> or contact our technical team.

Terms and conditions

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors