

Product datasheet

Anti-Lambda Light chain antibody [EPR5367-62] ab124719

Recombinant RabMAb

[3 References](#) [9 Images](#)

Overview

Product name	Anti-Lambda Light chain antibody [EPR5367-62]
Description	Rabbit monoclonal [EPR5367-62] to Lambda Light chain
Host species	Rabbit
Tested applications	Suitable for: Flow Cyt (Intra), WB, ELISA, IHC-P, ICC/IF
Species reactivity	Reacts with: Human
Immunogen	Full length native protein (purified) corresponding to Human Lambda Light chain. Purified Human IgA.
Positive control	WB: Human tonsil, plasma, thymus and spleen lysates. IHC-P: Human tonsil and colon tissues. ICC/IF: Ramos cells.
General notes	<p>This product is a recombinant monoclonal antibody, which offers several advantages including:</p> <ul style="list-style-type: none">- High batch-to-batch consistency and reproducibility- Improved sensitivity and specificity- Long-term security of supply- Animal-free production <p>For more information see here.</p> <p>Our RabMAb[®] technology is a patented hybridoma-based technology for making rabbit monoclonal antibodies. For details on our patents, please refer to RabMAb[®] patents.</p> <p>We are constantly working hard to ensure we provide our customers with best in class antibodies. As a result of this work we are pleased to now offer this antibody in purified format. We are in the process of updating our datasheets. The purified format is designated 'PUR' on our product labels. If you have any questions regarding this update, please contact our Scientific Support team.</p> <p>Mouse, Rat: We have preliminary internal testing data to indicate this antibody may not react with these species. Please contact us for more information.</p>

Properties

Form Liquid

Storage instructions	Shipped at 4°C. Store at +4°C short term (1-2 weeks). Upon delivery aliquot. Store at -20°C. Stable for 12 months at -20°C.
Storage buffer	pH: 7.20 Preservative: 0.01% Sodium azide Constituents: 40% Glycerol (glycerin, glycerine), 0.05% BSA, 59% PBS
Purity	Protein A purified
Clonality	Monoclonal
Clone number	EPR5367-62
Isotype	IgG

Applications

The Abpromise guarantee Our **Abpromise guarantee** covers the use of ab124719 in the following tested applications. The application notes include recommended starting dilutions; optimal dilutions/concentrations should be determined by the end user.

Application	Abreviews	Notes
Flow Cyt (Intra)		Use at an assay dependent concentration.
WB		1/20000. Predicted molecular weight: 25 kDa. For unpurified use at 1/50000 - 1/200000.
ELISA		1/64000.
IHC-P		1/250 - 1/500. Perform heat mediated antigen retrieval before commencing with IHC staining protocol. See <u>IHC antigen retrieval protocols</u> .
ICC/IF		1/250. For unpurified use at 1/500 - 1/1000.

Target


Relevance

All five immunoglobulin classes share the same basic four polypeptide chain structure of two heavy-chains and two light chains. There are five heavy chain types, and two light-chain types (Kappa and Lambda) both having a molecular weight of 22.5kDa. Any heavy-chain type can associate with either light-chain type, but on any immunoglobulin molecule both light-chains are of the same type. Kappa and Lambda consist of a variable region and a constant region and can easily be differentiated by the antigenic properties of the constant region. The ratio of Kappa to Lambda is 70:30, the vast majority of which is bound to heavy-chain in immunoglobulin. In normal individuals low levels of free light-chain are present in serum (Kappa, 1.6-15.2 mg/L; Lambda, 0.4-4.2mg/L), with the occurrence of multiple myeloma or other B-cell malignancies these levels can be greatly elevated and can be found at high levels in the urine (Bence-Jones proteins).

Cellular localization


Cytoplasmic

Images



Flow Cytometry (Intracellular) - Anti-Lambda Light chain antibody [EPR5367-62] (ab124719)

Intracellular Flow Cytometry analysis of Ramos (human Burkitt's lymphoma) cells labeling Lambda Light chain with unpurified ab124719 at 1/50 dilution (10ug/ml) (red). Cells were fixed with 4% paraformaldehyde and permeabilised with 90% methanol. A Goat anti rabbit IgG (Alexa Fluor® 488) (1/2000 dilution) was used as the secondary antibody. Rabbit monoclonal IgG (Black) was used as the isotype control, cells without incubation with primary antibody and secondary antibody (Blue) were used as the unlabeled control.



Western blot - Anti-Lambda Light chain antibody [EPR5367-62] (ab124719)

Anti-Lambda Light chain antibody [EPR5367-62] (ab124719) at 1/100000 dilution (purified) + Human plasma lysate at 20 µg


Secondary

Peroxidase-conjugated goat anti-rabbit IgG, (H+L) at 1/1000 dilution

Predicted band size: 25 kDa

Observed band size: 25 kDa

Blocking and dilution buffer: 5% NFDm/TBST.



Western blot - Anti-Lambda Light chain antibody [EPR5367-62] (ab124719)

Anti-Lambda Light chain antibody [EPR5367-62] (ab124719) at 1/20000 dilution (purified) + Human thymus tissue lysate at 20 µg


Secondary

Peroxidase-conjugated goat anti-rabbit IgG, (H+L) at 1/1000 dilution

Predicted band size: 25 kDa


Observed band size: 25 kDa

Blocking and dilution buffer: 5% NFDm/TBST.



Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-Lambda Light chain antibody [EPR5367-62] (ab124719)

Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) analysis of human colon tissue labelling Lambda Light chain with purified ab124719 at 1/300. Heat mediated antigen retrieval was performed using Tris/EDTA buffer pH 9. **ab97051**, a HRP-conjugated goat anti-rabbit IgG (H+L) was used as the secondary antibody (1/500). Negative control using PBS instead of primary antibody. Counterstained with hematoxylin.




Immunocytochemistry/ Immunofluorescence - Anti-Lambda Light chain antibody [EPR5367-62] (ab124719)

Immunocytochemistry/Immunofluorescence analysis of Ramos cells labelling Lambda Light chain with purified ab124719 at 1/250. Cells were fixed with 4% paraformaldehyde and permeabilized with 0.1% Triton X-100. **ab150077**, an Alexa Fluor[®] 488-conjugated goat anti-rabbit IgG (1/500) was used as the secondary antibody. DAPI (blue) was used as the nuclear counterstain. **ab7291**, a mouse anti-tubulin (1/1000) and **ab150120**, an Alexa Fluor[®] 594-conjugated goat anti-mouse IgG (1/500) were also used.

Control 1: primary antibody (1/250) and secondary antibody, **ab150120**, an Alexa Fluor[®] 594-conjugated goat anti-mouse IgG (1/500).


Control 2: **ab7291** (1/1000) and secondary antibody, **ab150077**, an Alexa Fluor[®] 488-conjugated goat anti-rabbit IgG (1/500).

**Indirect ELISA antibody dose-response curve
antigen at 1000 ng/ml**



ELISA - Anti-Lambda Light chain antibody
[EPR5367-62] (ab124719)

ELISA analysis of Human Kappa light chain (Free), Human Lambda Light Chains (Free), Human IgA, Human IgM, Human IgG, Rat IgG, Mouse IgG at 1000 ng/mL with ab124719 at 1000~0ng/mL. An Alkaline Phosphatase-conjugated AffiniPure Goat Anti-Rabbit IgG (H+L) at 1/2500 dilution was used as the secondary antibody.



Western blot - Anti-Lambda Light chain antibody
[EPR5367-62] (ab124719)

All lanes : Anti-Lambda Light chain antibody [EPR5367-62] (ab124719) at 1/50000 dilution (unpurified)

Lane 1 : Human tonsil lysate

Lane 2 : Human plasma lysate


Lane 3 : Human spleen lysate

Lysates/proteins at 10 µg per lane.

Secondary

All lanes : HRP-conjugated goat anti-rabbit IgG at 1/2000 dilution

Predicted band size: 25 kDa



Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-Lambda Light chain antibody [EPR5367-62] (ab124719)

Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) analysis of human tonsil tissue labelling Lambda Light chain with unpurified ab124719 at a dilution of 1/250.

Perform heat mediated antigen retrieval before commencing with IHC staining protocol.

Why choose a recombinant antibody?



Research with confidence
Consistent and reproducible results



Long-term and scalable supply
Recombinant technology



Success from the first experiment
Confirmed specificity



Ethical standards compliant
Animal-free production

Anti-Lambda Light chain antibody [EPR5367-62]
(ab124719)

Please note: All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <https://www.abcam.com/abpromise> or contact our technical team.

Terms and conditions

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors