abcam

Product datasheet

Anti-LRRK2 antibody [MJFF4 (c81-8)] - Low endotoxin, Azide free ab205335





2 Images

Overview

Product name Anti-LRRK2 antibody [MJFF4 (c81-8)] - Low endotoxin, Azide free

Rabbit monoclonal [MJFF4 (c81-8)] to LRRK2 - Low endotoxin, Azide free **Description**

Host species Rabbit

Tested applications Suitable for: WB, IHC-P, ICC/IF, IP

Species reactivity Reacts with: Mouse, Human

Predicted to work with: Rat

Immunogen Recombinant fragment corresponding to Human LRRK2 aa 950 to the C-terminus.

General notes ab205335 is the carrier-free version of ab133476.

> Our carrier-free antibodies are typically supplied in a PBS-only formulation, purified and free of BSA, sodium azide and glycerol. The carrier-free buffer and high concentration allow for increased conjugation efficiency.

This conjugation-ready format is designed for use with fluorochromes, metal isotopes, oligonucleotides, and enzymes, which makes them ideal for antibody labelling, functional and cellbased assays, flow-based assays (e.g. mass cytometry) and Multiplex Imaging applications.

Use our conjugation kits for antibody conjugates that are ready-to-use in as little as 20 minutes with <1 minute hands-on-time and 100% antibody recovery: available for fluorescent dyes, HRP, biotin and gold.

This product is compatible with the Maxpar® Antibody Labeling Kit from Fluidigm, without the need for antibody preparation. Maxpar[®] is a trademark of Fluidigm Canada Inc.

This product is a recombinant monoclonal antibody, which offers several advantages including:

- High batch-to-batch consistency and reproducibility
- Improved sensitivity and specificity
- Long-term security of supply
- Animal-free production

For more information see here.

Our RabMAb® technology is a patented hybridoma-based technology for making rabbit monoclonal antibodies. For details on our patents, please refer to **RabMAb patents**.

Our Low endotoxin, azide-free formats have low endotoxin level (≤ 1 EU/ml, determined by the

LAL assay) and are free from azide, to achieve consistent experimental results in functional assays.

This antibody was developed with support from The Michael J. Fox Foundation.



Properties

Form Liquid

Storage instructions Shipped at 4°C. Store at +4°C. Do Not Freeze.

Storage buffer pH: 7.2

Constituent: PBS

Carrier free Yes

Purity Protein A purified

Clonality Monoclonal

Clone number MJFF4 (c81-8)

Isotype IgG

Applications

The Abpromise guarantee

Our **Abpromise guarantee** covers the use of ab205335 in the following tested applications.

The application notes include recommended starting dilutions; optimal dilutions/concentrations should be determined by the end user.

Application	Abreviews	Notes
WB		Use at an assay dependent concentration. Predicted molecular weight: 286 kDa.
IHC-P		Use at an assay dependent concentration.
ICC/IF		Use at an assay dependent concentration.
IP		Use at an assay dependent concentration.

Target

Function

Positively regulates autophagy through a calcium-dependent activation of the CaMKK/AMPK signaling pathway. The process involves activation of nicotinic acid adenine dinucleotide phosphate (NAADP) receptors, increase in lysosomal pH, and calcium release from lysosomes. Together with RAB29, plays a role in the retrograde trafficking pathway for recycling proteins, such as mannose 6 phosphate receptor (M6PR), between lysosomes and the Golgi apparatus in a retromer-dependent manner. Regulates neuronal process morphology in the intact central nervous system (CNS). Plays a role in synaptic vesicle trafficking. Phosphorylates PRDX3. Has GTPase activity. May play a role in the phosphorylation of proteins central to Parkinson disease.

Tissue specificity

Expressed in the brain. Expressed in pyramidal neurons in all cortical laminae of the visual cortex, in neurons of the substantia nigra pars compacta and caudate putamen (at protein level). Expressed throughout the adult brain, but at a lower level than in heart and liver. Also expressed in placenta, lung, skeletal muscle, kidney and pancreas. In the brain, expressed in the cerebellum, cerebral cortex, medulla, spinal cord occipital pole, frontal lobe, temporal lobe and putamen. Expression is particularly high in brain dopaminoceptive areas.

Involvement in disease

Parkinson disease 8

Sequence similarities

Belongs to the protein kinase superfamily. TKL Ser/Thr protein kinase family.

Contains 12 LRR (leucine-rich) repeats. Contains 1 protein kinase domain.

Contains 1 Roc domain. Contains 7 WD repeats.

Domain

The seven-bladed WD repeat region is critical for synaptic vesicle trafficking and mediates interaction with multiple vesicle-associated presynaptic proteins.

The Roc domain mediates homodimerization and regulates kinase activity.


Post-translational modifications

Autophosphorylated.

Cellular localization

Membrane. Cytoplasm. Perikaryon. Mitochondrion. Golgi apparatus. Cell projection, axon. Cell projection, dendrite. Endoplasmic reticulum. Cytoplasmic vesicle, secretory vesicle, synaptic vesicle membrane. Endosome. Lysosome. Mitochondrion outer membrane. Mitochondrion inner membrane. Mitochondrion matrix. Predominantly associated with intracytoplasmic vesicular and membranous structures (By similarity). Localized in the cytoplasm and associated with cellular membrane structures. Predominantly associated with the mitochondrial outer membrane of the mitochondria. Colocalized with RAB29 along tubular structures emerging from Golgi apparatus. Localizes in intracytoplasmic punctate structures of neuronal perikarya and dendritic and axonal processes.

Images



Western blot - Anti-LRRK2 antibody [MJFF4 (c81-8)] - Low endotoxin, Azide free (ab205335)

This data was developed using the same antibody clone in a different buffer formulation (ab133476).

Lane 1: Wild-type HAP1 cell lysate (20 µg)

Lane 2: LRRK2 knockout HAP1 cell lysate (20 µg)

Lane 3: MEF cell lysate (20 µg)

Lane 4: LRRK2 knockout MEF cell lysate (20 µg)

Lane 5: A549 cell lysate (20ug)


Lane 6: LRRK2 knockout A549 cell lysate (20 µg)

Lanes 1 - 6: Merged signal (red and green). Green – <u>ab133476</u> observed at 238 kDa. Red - loading control, <u>ab130007</u>, observed at 124 kDa.

<u>ab133476</u> was shown to specifically react with LRRK2 in wild type A549, MEF and HAP1 cells along with additional cross-reactive bands. No band was observed when LRRK2 knockout samples were used. Wild-type and LRRK2 knockout samples were subjected to SDS-PAGE. <u>ab133476</u> and <u>ab130007</u> (loading

control to Vinculin) were diluted at 1/500 and 1/10000 respectively and incubated overnight at 4°C. Blots were developed with Goat anti-Rabbit lgG H&L (IRDye® 800CW) preadsorbed <u>ab216773</u> and Goat anti-Mouse lgG H&L (IRDye® 680RD) preadsorbed <u>ab216776</u> secondary antibodies at 1/10000 dilution for 1 hour at room temperature before imaging.

Wild-type and LRRK2 knockout MEF and A549 cells were provide as a generous gift from Professor Dario Alessi, MRC Protein Phosphorylation and Ubiquitination Unit (University of Dundee).



Please note: All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- · We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit https://www.abcam.com/abpromise or contact our technical team.

Terms and conditions

• Guarantee only valid for products bought direct from Abcam or one of our authorized distributors