


Anti-Mannose Receptor antibody ab64693

★★★★★ [40 Abreviews](#) [607 References](#) [7 Images](#)

Overview

Product name	Anti-Mannose Receptor antibody
Description	Rabbit polyclonal to Mannose Receptor
Host species	Rabbit
Specificity	From Jan 2024, QC testing of replenishment batches of this polyclonal changed. All tested and expected application and reactive species combinations are still covered by our Abcam product promise. However, we no longer test all applications. For more information on a specific batch, please contact our Scientific Support who will be happy to help.
Tested applications	Suitable for: IHC-P, WB, ICC/IF
Species reactivity	Reacts with: Mouse, Rat, Human Predicted to work with: Chimpanzee, Rhesus monkey 
Immunogen	Synthetic peptide conjugated to KLH derived from within residues 1400 to the C-terminus of Human Mannose Receptor. Read Abcam's proprietary immunogen policy (Peptide available as ab66694 .)
Positive control	WB: Rat liver and lung, human lung, mouse lung tissue lysate. IHC-P: Human, rat and mouse lung tissue. ICC/IF: MOLT-4 whole cells.
General notes	<p>The Life Science industry has been in the grips of a reproducibility crisis for a number of years. Abcam is leading the way in addressing this with our range of recombinant monoclonal antibodies and knockout edited cell lines for gold-standard validation. Please check that this product meets your needs before purchasing.</p> <p>If you have any questions, special requirements or concerns, please send us an inquiry and/or contact our Support team ahead of purchase. Recommended alternatives for this product can be found below, along with publications, customer reviews and Q&As</p>

Properties

Form	Liquid
Storage instructions	Shipped at 4°C. Store at +4°C short term (1-2 weeks). Upon delivery aliquot. Store at -20°C or -80°C. Avoid freeze / thaw cycle.
Storage buffer	pH: 7.40 Preservative: 0.02% Sodium azide Constituent: PBS

Batches of this product that have a concentration < 1mg/ml may have BSA added as a stabilising agent. If you would like information about the formulation of a specific lot, please contact our scientific support team who will be happy to help.

Purity	Immunogen affinity purified
Clonality	Polyclonal
Isotype	IgG

Applications


The Abpromise guarantee Our **Abpromise guarantee** covers the use of ab64693 in the following tested applications. The application notes include recommended starting dilutions; optimal dilutions/concentrations should be determined by the end user.

Application	Abreviews	Notes
IHC-P	★★★★★ (24)	Use a concentration of 0.05 - 0.1 µg/ml. Perform heat mediated antigen retrieval before commencing with IHC staining protocol. We do not recommend methanol fixation with this antibody
WB	★★★★☆ (3)	Use a concentration of 1 µg/ml. Predicted molecular weight: 166 kDa.
ICC/IF	★★★★★ (1)	Use a concentration of 1 µg/ml. Permeabilise with Tween (0.1%)

Target

Function	Mediates the endocytosis of glycoproteins by macrophages. Binds both sulfated and non-sulfated polysaccharide chains. Acts as phagocytic receptor for bacteria, fungi and other pathogens.
Sequence similarities	Contains 8 C-type lectin domains. Contains 1 fibronectin type-II domain. Contains 1 ricin B-type lectin domain.
Cellular localization	Membrane.

Images



All lanes : Anti-Mannose Receptor antibody (ab64693) at 1 µg/ml

Lane 1 : Rat liver tissue lysate

Lane 2 : Human lung tissue lysate

Lane 3 : Rat lung tissue lysate

Lane 4 : Mouse lung normal tissue lysate - total protein ([ab29297](#))

Lysates/proteins at 10 µg per lane.

Secondary

All lanes : Peroxidase AffiniPure Goat Anti-Rabbit IgG (H+L) at 1/5000 dilution


Developed using the ECL technique.

Performed under reducing conditions.

Predicted band size: 166 kDa


Observed band size: 190 kDa

Exposure time: 4 minutes



ab64693 staining Mannose Receptor in MOLT4 cells. The cells were fixed with 4% Formaldehyde (10 min), permeabilized with 0.1% PBS-Tween for 5 minutes and then blocked with 1% BSA/10% normal goat serum/0.3M glycine in 0.1% PBS-Tween for 1h. The cells were then incubated overnight at 4°C with ab64693 at 1µg/ml and [ab7291](#), Mouse monoclonal [DM1A] to alpha Tubulin - Loading Control. Cells were then incubated with [ab150081](#), Goat polyclonal Secondary Antibody to Rabbit IgG - H&L (Alexa Fluor® 488), pre-adsorbed at 1/1000 dilution (shown in green) and [ab150120](#), Goat polyclonal Secondary Antibody to Mouse IgG - H&L (Alexa Fluor® 594), pre-adsorbed at 1/1000 dilution (shown in pseudocolour red). Nuclear DNA was labelled with DAPI (shown in blue).

Image was acquired with a confocal microscope (Leica-Microsystems TCS SP8) and a single confocal section is shown.




Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-Mannose Receptor antibody (ab64693)

IHC image of ab64693 staining in human lung formalin fixed paraffin embedded tissue section*, performed on a Leica Bond™ system using the standard protocol F. The section was pre-treated using heat mediated antigen retrieval with sodium citrate buffer (pH6, epitope retrieval solution 1) for 20 mins. The section was then incubated with ab64693, 0.1 µg/ml, for 15 mins at room temperature and detected using an HRP conjugated compact polymer system. DAB was used as the chromogen. The section was then counterstained with haematoxylin and mounted with DPX.


For other IHC staining systems (automated and non-automated) customers should optimize variable parameters such as antigen retrieval conditions, primary antibody concentration and antibody incubation times.

*Tissue obtained from the Human Research Tissue Bank, supported by the NIHR Cambridge Biomedical Research Centre



Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-Mannose Receptor antibody (ab64693)


Immunohistochemical analysis of formalin fixed paraffin embedded human lung labelling mannose receptor with ab64693 at 0.1 µg/ml. The immunostaining was performed on a Ventana DISCOVERY ULTRA (Roche Tissue Diagnostics) instrument with an OptiView DAB IHC Detection Kit. Heat mediated antigen retrieval was conducted for 32min with ULTRA cell conditioning solution (CC1 pH8.5). ab64693 anti mannose receptor antibody was incubated at 37°C for 16min. Sections were counterstained is with Hematoxylin II. Image inset shows absence of staining in secondary antibody only control.



Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-Mannose Receptor antibody (ab64693)

This image is courtesy of an anonymous Abreview


Paraffin embedded, formalin-fixed mouse lung tissue labelling the mannose receptor with ab64693 at 1/500 in immunohistochemical analysis. An Alexa Fluor® 647 Goat anti-rabbit IgG was used as the secondary antibody, this was artificially colored green by software to enhance stain visibility. Heat mediated antigen retrieval was performed using Tris/EDTA buffer pH 9. 15% serum was used as blocking agent for 1 hour at room temperature.



Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-Mannose Receptor antibody (ab64693)

IHC image of Mannose Receptor staining in a section of formalin-fixed paraffin-embedded normal rat lung performed on a Leica BOND™ system using the standard protocol. The section was pre-treated using heat mediated antigen retrieval with sodium citrate buffer (pH6, epitope retrieval solution 1) for 20mins. The section was then incubated with ab64693, 0.01ug/ml, for 15 mins at room temperature and detected using an HRP conjugated compact polymer system. DAB was used as the chromogen. The section was then counterstained with haematoxylin and mounted with DPX. The inset secondary-only control image is taken from an identical assay without primary antibody.

For other IHC staining systems (automated and non-automated) customers should optimize variable parameters such as antigen retrieval conditions, primary antibody concentration and antibody incubation times.



Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-Mannose Receptor antibody (ab64693)

IHC image of Mannose Receptor staining in a section of formalin-fixed paraffin-embedded normal mouse lung performed on a Leica BOND™ system using the standard protocol. The section was pre-treated using heat mediated antigen retrieval with sodium citrate buffer (pH6, epitope retrieval solution 1) for 20mins. The section was then incubated with ab64693, 0.01ug/ml, for 15 mins at room temperature and detected using an HRP conjugated compact polymer system. DAB was used as the chromogen. The section was then counterstained with haematoxylin and mounted with DPX. The inset secondary-only control image is taken from an identical assay without primary antibody.

For other IHC staining systems (automated and non-automated) customers should optimize variable parameters such as antigen retrieval conditions, primary antibody concentration and antibody incubation times.

Please note: All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <https://www.abcam.com/abpromise> or contact our technical team.

Terms and conditions

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors