

Product datasheet

Anti-MEK1 (phospho S298) antibody [EPR3338] ab96379

Recombinant RabMAb

★★★★★ **6 Abreviews** **57 References** [7 Images](#)

Overview

Product name	Anti-MEK1 (phospho S298) antibody [EPR3338]
Description	Rabbit monoclonal [EPR3338] to MEK1 (phospho S298)
Host species	Rabbit
Specificity	ab96379 detects MEK1 phosphorylated at threonine 298. The mouse and rat recommendation is based on the WB results. We do not guarantee IHC-P for mouse and rat.
Tested applications	Suitable for: IHC-P, Flow Cyt (Intra), ICC/IF, WB Unsuitable for: IP
Species reactivity	Reacts with: Mouse, Rat, Human
Immunogen	Synthetic peptide. This information is proprietary to Abcam and/or its suppliers.
Positive control	Flow Cyt (intra): HeLa cells; ICC/IF: HeLa cells; IHC-P: Human ovarian carcinoma tissue; WB; Rat and mouse skeletal muscle, HeLa cells.
General notes	This product is a recombinant monoclonal antibody, which offers several advantages including: <ul style="list-style-type: none"> - High batch-to-batch consistency and reproducibility - Improved sensitivity and specificity - Long-term security of supply - Animal-free production For more information see here . Our RabMAb [®] technology is a patented hybridoma-based technology for making rabbit monoclonal antibodies. For details on our patents, please refer to RabMAb[®] patents .

Properties

Form	Liquid
Storage instructions	Shipped at 4°C. Store at -20°C. Stable for 12 months at -20°C.
Storage buffer	pH: 7.20 Preservative: 0.01% Sodium azide Constituents: 59% PBS, 40% Glycerol (glycerin, glycerine), 0.05% BSA
Purity	Protein A purified

Clonality	Monoclonal
Clone number	EPR3338
Isotype	IgG

Applications

The Abpromise guarantee Our **Abpromise guarantee** covers the use of ab96379 in the following tested applications. The application notes include recommended starting dilutions; optimal dilutions/concentrations should be determined by the end user.


Application	Abreviews	Notes
IHC-P		1/50. Perform heat mediated antigen retrieval with citrate buffer pH 6 before commencing with IHC staining protocol. The mouse and rat recommendation is based on the WB results. We do not guarantee IHC-P for mouse and rat. For unpurified use at 1/100 - 1/250 dilution.
Flow Cyt (Intra)		1/100.
ICC/IF	★★★★★ (3)	1/100 - 1/250.
WB	★★★★★ (3)	1/1000 - 1/5000. Detects a band of approximately 45 kDa (predicted molecular weight: 43 kDa).

Application notes Is unsuitable for IP.

Target

Function	Catalyzes the concomitant phosphorylation of a threonine and a tyrosine residue in a Thr-Glu-Tyr sequence located in MAP kinases. Activates ERK1 and ERK2 MAP kinases.
Tissue specificity	Widely expressed, with extremely low levels in brain.
Involvement in disease	Defects in MAP2K1 are a cause of cardiofaciocutaneous syndrome (CFC syndrome) [MIM:115150]; also known as cardio-facio-cutaneous syndrome. CFC syndrome is characterized by a distinctive facial appearance, heart defects and mental retardation. Heart defects include pulmonic stenosis, atrial septal defects and hypertrophic cardiomyopathy. Some affected individuals present with ectodermal abnormalities such as sparse, friable hair, hyperkeratotic skin lesions and a generalized ichthyosis-like condition. Typical facial features are similar to Noonan syndrome. They include high forehead with bitemporal constriction, hypoplastic supraorbital ridges, downslanting palpebral fissures, a depressed nasal bridge, and posteriorly angulated ears with prominent helices. The inheritance of CFC syndrome is autosomal dominant.
Sequence similarities	Belongs to the protein kinase superfamily. STE Ser/Thr protein kinase family. MAP kinase kinase subfamily. Contains 1 protein kinase domain.
Post-translational modifications	Phosphorylation on Ser/Thr by MAP kinase kinase kinases (RAF or MEKK1) regulates positively the kinase activity. Acetylation by Yersinia yopJ prevents phosphorylation and activation, thus blocking the MAPK signaling pathway.

Images



Western blot - Anti-MEK1 (phospho S298) antibody [EPR3338] (ab96379)

All lanes : Anti-MEK1 (phospho S298) antibody [EPR3338] (ab96379) at 1/1000 dilution (Purified)


Lane 1 : Rat skeletal muscle lysate

Lane 2 : Rat skeletal muscle lysate, the membrane treated with phosphatase for 1 hour

Secondary


All lanes : Goat Anti-Rabbit IgG H&L (HRP) ([ab97051](#)) at 1/20000 dilution

Predicted band size: 43 kDa



Flow Cytometry (Intracellular) - Anti-MEK1 (phospho S298) antibody [EPR3338] (ab96379)

Intracellular Flow Cytometry analysis of HeLa (Human cervix adenocarcinoma epithelial cell) cells labeling MEK1 with Purified ab96379 at 1/100 dilution (10 µg/mL) (Red). Cells were fixed with 4% Paraformaldehyde and permeabilised with 90% Methanol. A Goat anti rabbit IgG (Alexa Fluor® 488, [ab150077](#)) secondary antibody was used at 1/2000. Isotype control - Rabbit monoclonal IgG (Black). Unlabeled control - Cell without incubation with primary antibody and secondary antibody (Blue).



All lanes : Anti-MEK1 (phospho S298) antibody [EPR3338] (ab96379) at 1/1000 dilution (Purified)


Lane 1 : Mouse skeletal muscle lysate

Lane 2 : Mouse skeletal muscle lysate, the membrane treated with phosphatase for 1 hour


Secondary

All lanes : Goat Anti-Rabbit IgG H&L (HRP) (**ab97051**) at 1/20000 dilution

Predicted band size: 43 kDa



Immunocytochemistry analysis of HeLa (Human cervix adenocarcinoma epithelial cell) treated with lambda phosphatase cells labeling MEK1 with Purified ab96379 at 1/100 dilution (9.53 µg/mL). Cells were fixed in 4% Paraformaldehyde and permeabilized with 0.1% tritonX-100. Cells were counterstained with Ab195889 Anti-alpha Tubulin antibody [DM1A] - Microtubule Marker (Alexa Fluor® 594) 1/200 dilution (2.5 µg/mL). Goat anti rabbit IgG (Alexa Fluor® 488, **ab150077**) was used as the secondary antibody at 1/1000 dilution (2 µg/mL). DAPI (blue) was used as nuclear counterstain. PBS instead of the primary antibody was used as the secondary antibody only control.



All lanes : Anti-MEK1 (phospho S298) antibody [EPR3338] (ab96379) at 1/1000 dilution (Purified)

Lane 1 : HeLa (Human cervix adenocarcinoma epithelial cell) whole cell lysate


Lane 2 : HeLa (Human cervix adenocarcinoma epithelial cell) treated with 100 ng/ml nocodazole for 18 hours whole cell lysate

Lane 3 : HeLa (Human cervix adenocarcinoma epithelial cell) treated with 100ng/ml nocodazole for 18 hours, then the membrane treated with phosphatase for 1 hour

Secondary

All lanes : Goat Anti-Rabbit IgG H&L (HRP) ([ab97051](#)) at 1/20000 dilution

Predicted band size: 43 kDa



Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) analysis of human ovarian carcinoma tissue sections labeling MEK1 with Purified ab96379 at 1/50 dilution (19.06 µg/mL). Perform heat mediated antigen retrieval using [ab93684](#) (Tris/EDTA buffer, pH 9.0). ImmunoHistoProbe one step HRP Polymer (ready to use) was used as the secondary antibody. Negative control: PBS instead of the primary antibody. Hematoxylin was used as a counterstain.

Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-MEK1 (phospho S298) antibody [EPR3338] (ab96379)

Why choose a recombinant antibody?



Research with confidence
Consistent and reproducible results



Long-term and scalable supply
Recombinant technology



Success from the first experiment
Confirmed specificity



Ethical standards compliant
Animal-free production

Anti-MEK1 (phospho S298) antibody [EPR3338]
(ab96379)

Please note: All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <https://www.abcam.com/abpromise> or contact our technical team.

Terms and conditions

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors