

## Product datasheet

### Anti-MMP2 antibody [EPR1184] ab92536

Recombinant **RabMAb**

★★★★★ [7 Abreviews](#) [305 References](#) [10 Images](#)

#### Overview

<b>Product name</b>	Anti-MMP2 antibody [EPR1184]
<b>Description</b>	Rabbit monoclonal [EPR1184] to MMP2
<b>Host species</b>	Rabbit
<b>Specificity</b>	<p>In Western Blot, this product typically gives a weaker signal in some hepatocarcinoma like HepG2, MHCC97L(PMID:33184263 ) et al. Please use recommended positive control when testing these cells.</p> <p>Compared with ab92536, <b>ab181286</b> has higher sensitivity. We recommend <b>ab181286</b> as an alternative for testing MMP2 in western blot.</p>
<b>Tested applications</b>	<p><b>Suitable for:</b> Flow Cyt (Intra), ICC/IF, WB</p> <p><b>Unsuitable for:</b> IHC-P</p>
<b>Species reactivity</b>	<b>Reacts with:</b> Mouse, Rat, Human
<b>Immunogen</b>	<p>Synthetic peptide within Human MMP2 aa 550-650 (C terminal). The exact sequence is proprietary.</p> <p>Database link: <b>P08253</b></p>
<b>Positive control</b>	<p>WB: HepG2, L6, Raw264.7 and NIH/3T3 cell lysates; fetal heart and human skin tissue lysate; Human plasma, brain and breast tissue lysate ICC/IF: PC-3 cells Flow Cyt (intra): HeLa and PC-3 cells</p>
<b>General notes</b>	<p>This product is a recombinant monoclonal antibody, which offers several advantages including:</p> <ul style="list-style-type: none"> <li>- High batch-to-batch consistency and reproducibility</li> <li>- Improved sensitivity and specificity</li> <li>- Long-term security of supply</li> <li>- Animal-free production</li> </ul> <p>For more information <a href="#">see here</a>.</p> <p>Our RabMAb<sup>®</sup> technology is a patented hybridoma-based technology for making rabbit monoclonal antibodies. For details on our patents, please refer to <a href="#">RabMAb<sup>®</sup> patents</a>.</p>

#### Properties

<b>Form</b>	Liquid
<b>Storage instructions</b>	Shipped at 4°C. Store at +4°C short term (1-2 weeks). Upon delivery aliquot. Store at -20°C.

	Stable for 12 months at -20°C.
<b>Storage buffer</b>	pH: 7.20 Preservative: 0.01% Sodium azide Constituents: 59% PBS, 40% Glycerol, 0.05% BSA
<b>Purity</b>	Protein A purified
<b>Clonality</b>	Monoclonal
<b>Clone number</b>	EPR1184
<b>Isotype</b>	IgG

## Applications

**The Abpromise guarantee** Our **Abpromise guarantee** covers the use of ab92536 in the following tested applications. The application notes include recommended starting dilutions; optimal dilutions/concentrations should be determined by the end user.

Application	Abreviews	Notes
Flow Cyt (Intra)		1/400. For unpurified, use 1/70. <b>ab172730</b> - Rabbit monoclonal IgG, is suitable for use as an isotype control with this antibody.
ICC/IF	★★★★★ (1)	1/250.
WB	★★★★★ (4)	1/1000 - 1/5000. Predicted molecular weight: 74 kDa. For Lysate preparation protocol, please refer to the protocol <b>here (downloadable copy)</b> . Compared with ab92536, <b>ab181286</b> has higher sensitivity. We recommend <b>ab181286</b> as an alternative for testing MMP2 in western blot.

**Application notes** Is unsuitable for IHC-P.

## Target

<b>Function</b>	Ubiquitous metalloproteinase that is involved in diverse functions such as remodeling of the vasculature, angiogenesis, tissue repair, tumor invasion, inflammation, and atherosclerotic plaque rupture. As well as degrading extracellular matrix proteins, can also act on several nonmatrix proteins such as big endothelial 1 and beta-type CGRP promoting vasoconstriction. Also cleaves KISS at a Gly-Leu bond. Appears to have a role in myocardial cell death pathways. Contributes to myocardial oxidative stress by regulating the activity of GSK3beta. Cleaves GSK3beta in vitro. PEX, the C-terminal non-catalytic fragment of MMP2, possesses anti-angiogenic and anti-tumor properties and inhibits cell migration and cell adhesion to FGF2 and vitronectin. Ligand for integrin/beta3 on the surface of blood vessels.
<b>Tissue specificity</b>	Produced by normal skin fibroblasts. PEX is expressed in a number of tumors including gliomas, breast and prostate.
<b>Involvement in disease</b>	Defects in MMP2 are the cause of Torg-Winchester syndrome (TWS) [MIM:259600]; also known as multicentric osteolysis nodulosis and arthropathy (MONA). TWS is an autosomal recessive osteolysis syndrome. It is severe with generalized osteolysis and osteopenia. Subcutaneous

nodules are usually absent. Torg-Winchester syndrome has been associated with a number of additional features including coarse face, corneal opacities, patches of thickened, hyperpigmented skin, hypertrichosis and gum hypertrophy. However, these features are not always present and have occasionally been observed in other osteolysis syndromes.

## Sequence similarities

Belongs to the peptidase M10A family.

Contains 3 fibronectin type-II domains.

Contains 4 hemopexin-like domains.

## Domain

The conserved cysteine present in the cysteine-switch motif binds the catalytic zinc ion, thus inhibiting the enzyme. The dissociation of the cysteine from the zinc ion upon the activation-peptide release activates the enzyme.


## Post-translational modifications

Phosphorylation on multiple sites modulates enzymatic activity. Phosphorylated by PKC in vitro. The propeptide is processed by MMP14 (MT-MMP1) and MMP16 (MT-MMP3). Autocatalytic cleavage in the C-terminal produces the anti-angiogenic peptide, PEX. This processing appears to be facilitated by binding integrin $\alpha$ V/ $\beta$ 3.

## Cellular localization

Secreted > extracellular space > extracellular matrix. Membrane. Nucleus. Colocalizes with integrin  $\alpha$ V/ $\beta$ 3 at the membrane surface in angiogenic blood vessels and melanomas. Found in mitochondria, along microfibrils, and in nuclei of cardiomyocytes.

## Images



**All lanes :** Anti-MMP2 antibody [EPR1184] (ab92536) at 1/1000 dilution

**Lane 1 :** Human plasma tissue lysate

**Lane 2 :** Human brain tissue lysate

**Lane 3 :** Human breast tissue lysate

Lysates/proteins at 20  $\mu$ g per lane.

## Secondary

**All lanes :** Goat Anti-Rabbit IgG (HRP) with minimal cross-reactivity with human IgG at 1/2000 dilution

**Predicted band size:** 74 kDa

**Observed band size:** 69,72 kDa


**Exposure time:** 60 seconds

Blocking and diluting buffer and concentration: 5% NFDm/TBST.

**ab181602** was used as a GAPDH loading control.

Compared with ab92536, **ab181286** has higher sensitivity. We recommend **ab181286** as an alternative for testing MMP2 in

western blot.



**All lanes :** Anti-MMP2 antibody [EPR1184] (ab92536) at 1/1000 dilution

**Lane 1 :** HT-1080 (Human fibrosarcoma epithelial cell) whole cell lysate

**Lane 2 :** Untreated HepG2 (Human hepatocellular carcinoma epithelial cell) whole cell lysate

**Lane 3 :** HepG2 (Human hepatocellular carcinoma epithelial cell) treated with 300ng/ml BFA for 24 hours whole cell lysate

**Lane 4 :** Huh7 (Human hepatocellular carcinoma epithelial cell) whole cell lysate

**Lane 5 :** U-87 MG (Human glioblastoma-astrocytoma epithelial cell) whole cell lysate

Lysates/proteins at 20 µg per lane.

### Secondary

**All lanes :** Goat Anti-Rabbit IgG (HRP) with minimal cross-reactivity with human IgG at 1/2000 dilution

**Predicted band size:** 74 kDa


**Observed band size:** 69,72 kDa

**Exposure time:** 60 seconds

Blocking and diluting buffer and concentration: 5% NFDM /TBST.

**ab181602** was used as GAPDH loading control.

Compared with ab92536, **ab181286** has higher sensitivity. We recommend **ab181286** as an alternative for testing MMP2 in western blot.



**All lanes :** Anti-MMP2 antibody [EPR1184] (ab92536) at 1/1000 dilution

**Lane 1 :** L6 (Rat skeletal muscle myoblast) whole cell lysates prepared in 1%SDS Hot lysis method with 5% NFDm/TBST

**Lane 2 :** Mouse liver lysates prepared in RIPA lysis method with 5% NFDm/TBST

**Lane 3 :** Mouse liver lysates prepared in 1%SDS Hot lysis method with 5% NFDm/TBST

**Lane 4 :** Raw264.7 (Mouse Abelson murine leukemia virus-induced tumor macrophage) whole cell lysates prepared in RIPA lysis method with 5% NFDm/TBST

**Lane 5 :** Raw264.7 (Mouse Abelson murine leukemia virus-induced tumor macrophage) whole cell lysates prepared in 1%SDS Hot lysis method with 5% NFDm/TBST

**Lane 6 :** HepG2 (Human hepatocellular carcinoma epithelial cell) whole cell lysates prepared in RIPA lysis method with 5% NFDm/TBST

**Lane 7 :** HepG2 (Human hepatocellular carcinoma epithelial cell) whole cell lysates prepared in 1%SDS Hot lysis method with 5% NFDm/TBST

Lysates/proteins at 20 µg per lane.

### Secondary

**All lanes :** Goat Anti-Rabbit IgG H&L (HRP) (**ab97051**) at 1/20000 dilution (Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated)

**Predicted band size:** 74 kDa


**Observed band size:** 69,72 kDa

**Exposure time:** 10 seconds

The 72 KDa band is pro-MMP2 and the 69 KDa band is active-


MMP2 reported by PMID 11489818 and 22190701.

This antibody shows low affinity in detecting mouse liver and HepG2 lysates which are positive for MMP2 reported by PMID 24096707 and 24297510.




Immunocytochemistry/ Immunofluorescence - Anti-MMP2 antibody [EPR1184] (ab92536)

Immunofluorescence staining of PC-3 cells with purified ab92536 at a working dilution of 1/250, counter-stained with DAPI. The secondary antibody was Alexa Fluor<sup>®</sup> 488 goat anti-rabbit (**ab150077**), used at a dilution of 1/1000. **ab7291**, a mouse anti-tubulin antibody (1/1000), was used to stain tubulin along with **ab150120** (Alexa Fluor<sup>®</sup> 594 goat anti-mouse, 1/1000), shown in the top right hand panel. The cells were fixed in 4% PFA and permeabilized using 0.1% Triton X 100. The negative controls are shown in bottom middle and right hand panels - for negative control 1, purified ab92536 was used at a dilution of 1/500 followed by an Alexa Fluor<sup>®</sup> 594 goat anti-mouse antibody (**ab150120**) at a dilution of 1/500. For negative control 2, **ab7291** (mouse anti-tubulin) was used at a dilution of 1/500 followed by an Alexa Fluor<sup>®</sup> 488 goat anti-rabbit antibody (**ab150077**) at a dilution of 1/400.



Flow Cytometry (Intracellular) - Anti-MMP2 antibody [EPR1184] (ab92536)

Intracellular Flow Cytometry analysis of PC-3 (human prostate adenocarcinoma) cells labeling MMP2 with purified ab92536 at 1/180 dilution (10ug/ml) (red). Cells were fixed with 4% paraformaldehyde and permeabilised with 90% methanol. A Goat anti rabbit IgG (Alexa Fluor<sup>®</sup> 488) (1/2000 dilution) was used as the secondary antibody. Rabbit monoclonal IgG (Black) was used as the isotype control, cells without incubation with primary antibody and secondary antibody (Blue) were used as the unlabeled control.



**All lanes :** Anti-MMP2 antibody [EPR1184] (ab92536) at 1/10000 dilution (purified)

**Lane 1 :** L6 cell lysate

**Lane 2 :** NIH/3T3 cell lysate

Lysates/proteins at 20 µg per lane.

#### Secondary

**All lanes :** HRP goat anti-rabbit IgG (H+L) at 1/1000 dilution


**Predicted band size:** 74 kDa

**Observed band size:** 64,72 kDa

Blocking buffer: 5% NFDM/TBST

Dilution buffer: 5% NFDM/TBST

72kDa: propeptide; 64kDa: active form



**All lanes :** Anti-MMP2 antibody [EPR1184] (ab92536) at 1/1000 dilution (unpurified)

**Lane 1 :** L6 cell lysate

**Lane 2 :** Fetal heart lysate


**Lane 3 :** NIH/3T3 cell lysate

Lysates/proteins at 10 µg per lane.

#### Secondary


**All lanes :** HRP labelled goat anti-rabbit at 1/2000 dilution

**Predicted band size:** 74 kDa



Flow Cytometry (Intracellular) - Anti-MMP2 antibody  
[EPR1184] (ab92536)

Overlay histogram showing HeLa cells fixed in 4% PFA and stained with purified ab92536 at a dilution of 1 in 400 (red line). The secondary antibody used was FITC goat anti-rabbit at a dilution of 1 in 500. Rabbit monoclonal IgG was used as an isotype control (black line) and cells incubated in the absence of both primary and secondary antibody were used as a negative control (blue line).



Western blot - Anti-MMP2 antibody [EPR1184]  
(ab92536)

Anti-MMP2 antibody [EPR1184] (ab92536) at 1/5000 dilution  
(purified) + human skin at 10 µg

#### Secondary

HRP goat anti-rabbit IgG (H+L) at 1/1000 dilution

**Predicted band size:** 74 kDa


**Observed band size:** 64,72 kDa

Blocking buffer: 5% NFDM/TBST

Dilution buffer: 5% NFDM/TBST

72kDa: propeptide; 64kDa: active form

#### Why choose a recombinant antibody?



Anti-MMP2 antibody [EPR1184] (ab92536)



**Please note:** All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

### **Our Abpromise to you: Quality guaranteed and expert technical support**

---

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
  
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <https://www.abcam.com/abpromise> or contact our technical team.

### **Terms and conditions**

---

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors