

# Anti-N myc interactor/NMI antibody [EPR11065(2)] - BSA and Azide free ab250698

**KO VALIDATED** Recombinant RabMAB

7 Images

### Overview

|                            |  |
|----------------------------|--|
| <b>Product name</b>        | Anti-N myc interactor/NMI antibody [EPR11065(2)] - BSA and Azide free                        |
| <b>Description</b>         | Rabbit monoclonal [EPR11065(2)] to N myc interactor/NMI - BSA and Azide free                 |
| <b>Host species</b>        | Rabbit   |
| <b>Tested applications</b> | <b>Suitable for:</b> IHC-P, IP, WB, ICC/IF   |
| <b>Species reactivity</b>  | <b>Reacts with:</b> Human  |
| <b>Immunogen</b>           | Synthetic peptide. This information is proprietary to Abcam and/or its suppliers.            |
| <b>Positive control</b>    | WB: HeLa, A549, K562 and HepG2 cell lysates. IHC-P: Human tonsil tissue. ICC/IF: HeLa cells. |
| <b>General notes</b>       | ab250698 is the carrier-free version of <a href="#">ab183724</a> .                           |

Our **carrier-free** antibodies are typically supplied in a PBS-only formulation, purified and free of BSA, sodium azide and glycerol. The carrier-free buffer and high concentration allow for increased conjugation efficiency.

This conjugation-ready format is designed for use with fluorochromes, metal isotopes, oligonucleotides, and enzymes, which makes them ideal for antibody labelling, functional and cell-based assays, flow-based assays (e.g. mass cytometry) and Multiplex Imaging applications.

Use our **conjugation kits** for antibody conjugates that are ready-to-use in as little as 20 minutes with <1 minute hands-on-time and 100% antibody recovery: available for fluorescent dyes, HRP, biotin and gold.

This product is compatible with the Maxpar<sup>®</sup> Antibody Labeling Kit from Fluidigm, without the need for antibody preparation. Maxpar<sup>®</sup> is a trademark of Fluidigm Canada Inc.

This product is a recombinant monoclonal antibody, which offers several advantages including:

- High batch-to-batch consistency and reproducibility
- Improved sensitivity and specificity
- Long-term security of supply
- Animal-free production

For more information [see here](#).

Our RabMAB<sup>®</sup> technology is a patented hybridoma-based technology for making rabbit monoclonal antibodies. For details on our patents, please refer to [RabMAB<sup>®</sup> patents](#).

## Properties

---

|                             |   |
|-----------------------------|---|
| <b>Form</b>                 | Liquid  |
| <b>Storage instructions</b> | Shipped at 4°C. Store at +4°C. Do Not Freeze. |
| <b>Storage buffer</b>       | pH: 7.2<br>Constituent: PBS                   |
| <b>Carrier free</b>         | Yes   |
| <b>Purity</b>               | Protein A purified                            |
| <b>Clonality</b>            | Monoclonal                                    |
| <b>Clone number</b>         | EPR11065(2)                                   |
| <b>Isotype</b>              | IgG   |

## Applications

---

**The Abpromise guarantee** Our **Abpromise guarantee** covers the use of ab250698 in the following tested applications. The application notes include recommended starting dilutions; optimal dilutions/concentrations should be determined by the end user.

| Application | Abreviews | Notes   |
|-------------|-----------|---|
| IHC-P       |           | Use at an assay dependent concentration. Perform heat mediated antigen retrieval with Tris/EDTA buffer pH 9.0 before commencing with IHC staining protocol. |
| IP          |           | Use at an assay dependent concentration.  |
| WB          |           | Use at an assay dependent concentration. Detects a band of approximately 38 kDa (predicted molecular weight: 35 kDa).                                       |
| ICC/IF      |           | Use at an assay dependent concentration.  |

## Target


---

**Relevance** NMYC interactor (NMI) encodes a protein that interacts with NMYC and CMYC (two members of the oncogene Myc family), and other transcription factors containing a Zip, HLH, or HLH Zip motif. The NMI protein also interacts with all STATs except STAT2 and augments STAT mediated transcription in response to cytokines IL2 and IFN gamma. The NMI mRNA has high expression in myeloid leukemia cell lines.

**Cellular localization** Cytoplasmic

## Images

---



Western blot - Anti-N myc interactor/NMI antibody [EPR11065(2)] - BSA and Azide free (ab250698)

**All lanes** : Anti-N myc interactor/NMI antibody [EPR11065(2)] ([ab183724](#)) at 1/1000 dilution

**Lane 1** : Wild-type A549 (Human lung carcinoma cell line) whole cell lysate

**Lane 2** : NMI knockout A549 (Human lung carcinoma cell line) whole cell lysate

**Lane 3** : HeLa (Human epithelial cell line from cervix adenocarcinoma) whole cell lysate

**Lane 4** : K562 (Human chronic myelogenous leukemia lymphoblast cell line ) whole cell lysate

Lysates/proteins at 20 µg per lane.

### Secondary

**All lanes** : Goat anti-Rabbit IgG H&L (IRDye® 800CW) preadsorbed ([ab216773](#)) at 1/10000 dilution

**Predicted band size:** 35 kDa


**Observed band size:** 39 kDa

This data was developed using [ab183724](#), the same antibody clone in a different buffer formulation.

**Lanes 1-4:** Merged signal (red and green). Green - [ab183724](#) observed at 39 kDa. Red - loading control [ab8245](#) observed at 36 kDa.


[ab183724](#) Anti-N myc interactor/NMI antibody [EPR11065(2)] was shown to specifically react with N myc interactor/NMI in wild-type A549 cells. Loss of signal was observed when knockout cell line [ab267013](#) (knockout cell lysate [ab258077](#)) was used. Wild-type and N myc interactor/NMI knockout samples were subjected to SDS-PAGE. [ab183724](#) and Anti-GAPDH antibody [6C5] - Loading Control ([ab8245](#)) were incubated overnight at 4°C at 1 in 1000 dilution and 1 in 20000 dilution respectively. Blots were developed with Goat anti-Rabbit IgG H&L (IRDye® 800CW) preadsorbed ([ab216773](#)) and Goat anti-Mouse IgG H&L (IRDye® 680RD) preadsorbed ([ab216776](#)) secondary antibodies at 1 in 20000

dilution for 1 hour at room temperature before imaging.



Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-N myc interactor/NMI antibody [EPR11065(2)] - BSA and Azide free (ab250698)

This data was developed using **ab183724**, the same antibody clone in a different buffer formulation. Immunohistochemical analysis of paraffin-embedded Human tonsil tissue labeling N myc interactor/NMI with **ab183724** at 1/50 dilution. The slide is counterstained with Hematoxylin. Perform heat mediated antigen retrieval with EDTA buffer pH 9 before commencing with IHC staining protocol.



Western blot - Anti-N myc interactor/NMI antibody [EPR11065(2)] - BSA and Azide free (ab250698)

This data was developed using **ab183724**, the same antibody clone in a different buffer formulation.

**Lane 1:** Wild-type HAP1 cell lysate (20 µg)


**Lane 2:** N myc interactor/NMI knockout HAP1 cell lysate (20 µg)

**Lane 3:** HeLa cell lysate (20 µg)

**Lane 4:** K562 cell lysate (20 µg)

**Lanes 1 - 4:** Merged signal (red and green). Green - **ab183724** observed at 39 kDa. Red - loading control, **ab18058**, observed at 124 kDa.

**ab183724** was shown to specifically react with N myc interactor/NMI when N myc interactor/NMI knockout samples were used. Wild-type and N myc interactor/NMI knockout samples were subjected to SDS-PAGE. **ab183724** and **ab18058** (loading control to Vinculin) were diluted at 1/2000 and 1/10 000 respectively and incubated overnight at 4°C. Blots were developed with Goat anti-Rabbit IgG H&L (IRDye® 800CW) preadsorbed (**ab216773**) and Goat anti-Mouse IgG H&L (IRDye® 680RD) preadsorbed (**ab216776**) secondary antibodies at 1/10000 dilution for 1 hour at room temperature before imaging.



Western blot - Anti-N myc interactor/NMI antibody [EPR11065(2)] - BSA and Azide free (ab250698)

**All lanes** : Anti-N myc interactor/NMI antibody [EPR11065(2)] ([ab183724](#)) at 1/20000 dilution

**Lane 1** : K562 cell lysate

**Lane 2** : HeLa cell lysate

**Lane 3** : HepG2 cell lysate


Lysates/proteins at 20 µg per lane.

**Secondary**

**All lanes** : Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/1000 dilution


**Predicted band size:** 35 kDa

This data was developed using [ab183724](#), the same antibody clone in a different buffer formulation.



Immunocytochemistry/ Immunofluorescence - Anti-N myc interactor/NMI antibody [EPR11065(2)] - BSA and Azide free (ab250698)

This data was developed using [ab183724](#), the same antibody clone in a different buffer formulation. Immunofluorescence analysis of acetone-fixed HeLa cells labeling N myc interactor/NMI with [ab183724](#) at 1/100 dilution. Goat anti-rabbit IgG (Alexa Fluor® 488) at 1/200 dilution was used as the secondary antibody (green). The slide on the right is stained with Dapi (blue).



This data was developed using [ab183724](#), the same antibody clone in a different buffer formulation. Western blot analysis of K562 cell lysate precipitated with [ab183724](#) at 1/50 dilution. A Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated secondary antibody at 1/1000 dilution was then used. The blocking buffer and dilution buffer was 5% NFD/MTBST.

Immunoprecipitation - Anti-N myc interactor/NMI antibody [EPR11065(2)] - BSA and Azide free (ab250698)

### Why choose a recombinant antibody?

- 

**Research with confidence**  
Consistent and reproducible results
- 

**Long-term and scalable supply**  
Recombinant technology
- 

**Success from the first experiment**  
Confirmed specificity
- 

**Ethical standards compliant**  
Animal-free production

Anti-N myc interactor/NMI antibody [EPR11065(2)] - BSA and Azide free (ab250698)

**Please note:** All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

## **Our Abpromise to you: Quality guaranteed and expert technical support**

---

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
  
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <https://www.abcam.com/abpromise> or contact our technical team.

## **Terms and conditions**

---

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors