

Anti-NMDAR2B antibody [NR2B] ab28373

★★★★★ [3 Abreviews](#) [34 References](#) [5 Images](#)

Overview

Product name	Anti-NMDAR2B antibody [NR2B]
Description	Mouse monoclonal [NR2B] to NMDAR2B
Host species	Mouse
Specificity	Detects the N-methyl-D-aspartate (NMDA) receptor type 2B. This antibody has been tested in mouse and rat brain in western blotting. It has not been tested in human for Western blotting - only for ICC on transfected HEK cells.
Tested applications	Suitable for: WB, IHC-P, ICC, Flow Cyt
Species reactivity	Reacts with: Mouse, Rat, Human
Immunogen	Recombinant fragment corresponding to Rat NMDAR2B aa 934-1457.
Positive control	WB: Mouse brain tissue ICC: transfected HEK cells
General notes	<p>The Life Science industry has been in the grips of a reproducibility crisis for a number of years. Abcam is leading the way in addressing this with our range of recombinant monoclonal antibodies and knockout edited cell lines for gold-standard validation. Please check that this product meets your needs before purchasing.</p> <p>If you have any questions, special requirements or concerns, please send us an inquiry and/or contact our Support team ahead of purchase. Recommended alternatives for this product can be found below, along with publications, customer reviews and Q&As</p>

Properties

Form	Liquid
Storage instructions	Shipped at 4°C. Upon delivery aliquot and store at -20°C. Avoid freeze / thaw cycles.
Storage buffer	Preservative: 0.05% Sodium azide Constituents: PBS, 0.1% BSA
Purity	Protein A purified
Purification notes	Protein A Chromatography
Clonality	Monoclonal
Clone number	NR2B
Isotype	IgG1

Applications

The Abpromise guarantee

Our **Abpromise guarantee** covers the use of ab28373 in the following tested applications.

The application notes include recommended starting dilutions; optimal dilutions/concentrations should be determined by the end user.

Application	Abreviews	Notes
WB	★★★★☆ (2)	Use a concentration of 2 µg/ml. Predicted molecular weight: 166 kDa.
IHC-P		1/10 - 1/100.
ICC	★★★★☆ (1)	Use at an assay dependent concentration.
Flow Cyt		Use 1-2µg for 10 ⁶ cells. ab170190 - Mouse monoclonal IgG1, is suitable for use as an isotype control with this antibody.

Target

Function

NMDA receptor subtype of glutamate-gated ion channels with high calcium permeability and voltage-dependent sensitivity to magnesium. Mediated by glycine.

Tissue specificity

Primarily found in the fronto-parieto-temporal cortex and hippocampus pyramidal cells, lower expression in the basal ganglia.

Sequence similarities

Belongs to the glutamate-gated ion channel (TC 1.A.10.1) family. NR2B/GRIN2B subfamily.

Cellular localization


Cell membrane. Cell junction > synapse > postsynaptic cell membrane.

Images




Anti-NMDAR2B antibody [NR2B] (ab28373) at 2 µg/ml + mouse brain tissue

Predicted band size: 166 kDa



Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-NMDAR2B antibody [NR2B] (ab28373)

ab28373 staining NMDAR2B in Human breast tissue sections (right) compared with negative control (left) by Immunohistochemistry (IHC-P - paraformaldehyde-fixed, paraffin-embedded sections). Tissue was fixed with paraformaldehyde and blocked with 3% H₂O₂-methanol for 15 minutes at room temperature; antigen retrieval was by heat mediation in a sodium citrate buffer. Samples were incubated with primary antibody (1/20) overnight at 4°C. A HRP-conjugated secondary antibody was used for detection.



Western blot - Anti-NMDAR2B antibody [NR2B] (ab28373)


Lanes 2-4 : Anti-NMDAR2B antibody [NR2B] (ab28373) at 1/500 dilution

Lanes 5-7 : Anti-NMDAR2B antibody [NR2B] (ab28373) at 1/1000 dilution

Lanes 3 & 6 : Mouse brain (50ug)

Lanes 4 & 7 : Rat brain (50ug)

Predicted band size: 166 kDa



Western blot - Anti-NMDAR2B antibody [NR2B]
(ab28373)

This image is courtesy of an anonymous Abreview

Anti-NMDAR2B antibody [NR2B] (ab28373) at 1/500 dilution + Rat P14 Cortex Lysate, vesicular fraction at 5 µg

Secondary

HRP conjugated Donkey anti-mouse IgG at 1/15000 dilution


Developed using the ECL technique.

Performed under reducing conditions.

Predicted band size: 166 kDa

Samples were incubated with primary antibody for 12 hours at 4°C.

Blocked with 5% milk for 1 hour at 22°C.



Flow Cytometry - Anti-NMDAR2B antibody [NR2B]
(ab28373)

Overlay histogram showing SH-SY5Y cells stained with ab28373 (red line). The cells were fixed with 4% paraformaldehyde (10 min) and then permeabilized with 0.1% PBS-Tween for 20 min. The cells were then incubated in 1x PBS / 10% normal goat serum / 0.3M glycine to block non-specific protein-protein interactions followed by the antibody (ab28373, 1 µg/1x10⁶ cells) for 30 min at 22°C. The secondary antibody used was DyLight® 488 goat anti-mouse IgG (H+L) ([ab96879](#)) at 1/500 dilution for 30 min at 22°C. Isotype control antibody (black line) was mouse IgG1 [ICIGG1] ([ab91353](#), 2 µg/1x10⁶ cells) used under the same conditions. Acquisition of >5,000 events was performed. This antibody gave a positive signal in SH-SY5Y cells fixed with 80% methanol (5 min)/permeabilized with 0.1% PBS-Tween for 20 min used under the same conditions.

Please note: All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <https://www.abcam.com/abpromise> or contact our technical team.

Terms and conditions

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors