


## Product datasheet

# Anti-NQO1 antibody [A180] - BSA and Azide free ab264434

KO VALIDATED

7 Images

### Overview

<b>Product name</b>	Anti-NQO1 antibody [A180] - BSA and Azide free
<b>Description</b>	Mouse monoclonal [A180] to NQO1 - BSA and Azide free
<b>Host species</b>	Mouse
<b>Tested applications</b>	<b>Suitable for:</b> Flow Cyt (Intra), ICC/IF, Sandwich ELISA, WB, IHC-P
<b>Species reactivity</b>	<b>Reacts with:</b> Human <b>Predicted to work with:</b> Mouse, Rat, Dog, Monkey 
<b>Immunogen</b>	Recombinant full length protein corresponding to Human NQO1.
<b>Positive control</b>	WB: HAP1 and HepG2 whole cell lysates; human kidney tissue lysate. ICC/IF: HepG2 cells. Flow Cyt (Intra): HeLa cells. IHC-P: FFPE human breast adenocarcinoma and pancreas adenocarcinoma tissue sections.
<b>General notes</b>	<p>ab264434 is the carrier-free version of <a href="#">ab28947</a>.</p> <p>This antibody clone is manufactured by Abcam. If you require a custom buffer formulation or conjugation for your experiments, please contact <a href="mailto:orders@abcam.com">orders@abcam.com</a>.</p> <p>Our <b>carrier-free</b> antibodies are typically supplied in a PBS-only formulation, purified and free of BSA, sodium azide and glycerol. The carrier-free buffer and high concentration allow for increased conjugation efficiency.</p> <p>This conjugation-ready format is designed for use with fluorochromes, metal isotopes, oligonucleotides, and enzymes, which makes them ideal for antibody labelling, functional and cell-based assays, flow-based assays (e.g. mass cytometry) and Multiplex Imaging applications.</p> <p>Use our <b>conjugation kits</b> for antibody conjugates that are ready-to-use in as little as 20 minutes with &lt;1 minute hands-on-time and 100% antibody recovery: available for fluorescent dyes, HRP, biotin and gold.</p> <p>This product is compatible with the Maxpar<sup>®</sup> Antibody Labeling Kit from Fluidigm, without the need for antibody preparation. Maxpar<sup>®</sup> is a trademark of Fluidigm Canada Inc.</p> <p>The Life Science industry has been in the grips of a reproducibility crisis for a number of years. Abcam is leading the way in addressing this with our range of recombinant monoclonal antibodies and knockout edited cell lines for gold-standard validation. Please check that this product meets your needs before purchasing.</p>

If you have any questions, special requirements or concerns, please send us an inquiry and/or contact our Support team ahead of purchase. Recommended alternatives for this product can be found below, along with publications, customer reviews and Q&As

## Properties

<b>Form</b>	Liquid
<b>Storage instructions</b>	Shipped at 4°C. Store at +4°C. Do Not Freeze.
<b>Storage buffer</b>	Constituent: PBS
<b>Carrier free</b>	Yes
<b>Purity</b>	IgG fraction
<b>Clonality</b>	Monoclonal
<b>Clone number</b>	A180
<b>Isotype</b>	IgG1

## Applications


**The Abpromise guarantee** Our **Abpromise guarantee** covers the use of ab264434 in the following tested applications. The application notes include recommended starting dilutions; optimal dilutions/concentrations should be determined by the end user.

Application	Abreviews	Notes
Flow Cyt (Intra)		Use 1µg for 10 <sup>6</sup> cells. <b>ab170190</b> - Mouse monoclonal IgG1, is suitable for use as an isotype control with this antibody.
ICC/IF		Use at an assay dependent concentration.
Sandwich ELISA		1/500. Can be used as capture antibody in conjunction with <b>ab34173</b> as detection antibody.
WB		Use a concentration of 1 µg/ml. Predicted molecular weight: 30 kDa.
IHC-P		Use at an assay dependent concentration.

## Target

<b>Function</b>	The enzyme apparently serves as a quinone reductase in connection with conjugation reactions of hydroquinons involved in detoxification pathways as well as in biosynthetic processes such as the vitamin K-dependent gamma-carboxylation of glutamate residues in prothrombin synthesis.
<b>Sequence similarities</b>	Belongs to the NAD(P)H dehydrogenase (quinone) family.
<b>Cellular localization</b>	Cytoplasm.

## Images



Western blot - Anti-NQO1 antibody [A180] - BSA and Azide free (ab264434)

**Lane 1:** Wild-type HAP1 whole cell lysate (20 µg)

**Lane 2:** NQO1 knockout HAP1 whole cell lysate (20 µg)


**Lane 3:** HepG2 whole cell lysate (20 µg)

**Lanes 1 - 3:** Merged signal (red and green). Green - **ab28947** observed at 31 kDa. Red - loading control, **ab176560**, observed at 50 kDa.

**ab28947** was shown to specifically react with NQO1 in wild-type HAP1 cells as signal was lost in NQO1 knockout cells. Wild-type and NQO1 knockout samples were subjected to SDS-PAGE.

**ab28947** and **ab176560** (Rabbit anti-alpha Tubulin loading control) were incubated overnight at 4°C at 1 µg/ml and 1/20000 dilution respectively. Blots were developed with Goat anti-Mouse IgG H&L (IRDye® 800CW) preabsorbed **ab216772** and Goat anti-Rabbit IgG H&L (IRDye® 680RD) preabsorbed **ab216777** secondary antibodies at 1/20000 dilution for 1 hour at room temperature before imaging.

This data was developed using the same antibody clone in a different buffer formulation containing PBS and Azide (**ab28947**).




Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-NQO1 antibody [A180] - BSA and Azide free (ab264434)

IHC image of NQO1 staining in sections of formalin fixed paraffin embedded normal human pancreas\* (left) and human pancreas adenocarcinoma\* (right), performed on a Leica BOND™ system using the standard protocol F. The section was pre-treated using heat mediated antigen retrieval with sodium citrate buffer (pH6, epitope retrieval solution 1) for 20 mins. The section was then incubated with **ab28947**, 0.1 µg/ml, for 15 mins at room temperature and detected using an HRP conjugated compact polymer system. DAB was used as the chromogen. The section was then counterstained with haematoxylin and mounted with DPX.

For other IHC staining systems (automated and non-automated) customers should optimize variable parameters such as antigen retrieval conditions, primary antibody concentration and antibody incubation times.

*\*Tissue obtained from the Human Research Tissue Bank, supported by the NIHR Cambridge Biomedical Research Centre*

This data was developed using the same antibody clone in a different buffer formulation containing PBS and Azide (**ab28947**).




Immunocytochemistry/ Immunofluorescence - Anti-NQO1 antibody [A180] - BSA and Azide free (ab264434)

This data was developed using the same antibody clone in a different buffer formulation containing PBS and sodium azide (**ab28947**)

**ab28947** staining NQO1 in HepG2 cells. The cells were fixed with 100% methanol (5 min), permeabilized with 0.1% PBS-Tween for 5 minutes and then blocked with 1% BSA/10% normal goat serum/0.3M glycine in 0.1% PBS-Tween for 1h. The cells were then incubated overnight at 4°C with **ab28947** at 5 µg/ml and **ab6046**, Rabbit polyclonal to beta Tubulin - Loading Control. Cells were then incubated with **ab150117**, Goat polyclonal Secondary Antibody to Mouse IgG H&L (Alexa Fluor® 488) preadsorbed at 1/1000 dilution (shown in green) and **ab150080**, Goat polyclonal Secondary Antibody to Rabbit IgG - H&L (Alexa Fluor® 594) at 1/1000 dilution (shown in pseudocolour red). Nuclear DNA was labelled with DAPI (shown in blue).

Image was acquired with a high-content analyser (Operetta CLS, Perkin Elmer) and a maximum intensity projection of confocal sections is shown.




Flow Cytometry (Intracellular) - Anti-NQO1 antibody [A180] - BSA and Azide free (ab264434)

Overlay histogram showing HeLa cells stained with **ab28947** (red line). The cells were fixed with 80% methanol (5 min) and then permeabilized with 0.1% PBS-Tween for 20 min. The cells were then incubated in 1x PBS / 10% normal goat serum / 0.3M glycine to block non-specific protein-protein interactions followed by the antibody (**ab28947**, 1µg/1x10<sup>6</sup> cells) for 30 min at 22°C. The secondary antibody used was goat **anti-mouse DyLight® 488** (IgG H+L) (**ab96879**) at 1/500 dilution for 30 min at 22°C. Isotype control antibody (black line) was mouse IgG1 [ICIGG1] (**ab91353**, 2µg/1x10<sup>6</sup> cells) used under the same conditions.

Acquisition of >5,000 events was performed. This antibody gave a positive signal in HeLa cells fixed with 4% paraformaldehyde (10 min)/permeabilized with 0.1% PBS-Tween for 20 min used under the same conditions.

This data was developed using the same antibody clone in a different buffer formulation containing PBS and Azide (**ab28947**).




Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-NQO1 antibody [A180] - BSA and Azide free (ab264434)

IHC image of NQO1 staining in human breast adenocarcinoma formalin fixed paraffin embedded tissue section, performed on a Leica Bond system using the standard protocol F. The section was pre-treated using heat mediated antigen retrieval with sodium citrate buffer (pH6, epitope retrieval solution 1) for 20 mins. The section was then incubated with **ab28947**, 0.1µg/ml, for 15 mins at room temperature and detected using an HRP conjugated compact polymer system. DAB was used as the chromogen. The section was then counterstained with haematoxylin and mounted with DPX.

For other IHC staining systems (automated and non-automated) customers should optimize variable parameters such as antigen retrieval conditions, primary antibody concentration and antibody incubation times.


This data was developed using the same antibody clone in a different buffer formulation containing PBS and Azide (**ab28947**).



Sandwich ELISA - Anti-NQO1 antibody [A180] - BSA and Azide free (ab264434)

Sandwich ELISA for the detection of NQO1, using **ab28947** (1/500) as the capture antibody and **ab34173** (1/1000) for the detection

This data was developed using the same antibody clone in a different buffer formulation containing PBS and Azide (**ab28947**).



2006 Abcam

Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-NQO1 antibody [A180] - BSA and Azide free (ab264434)

Human breast cancer tissue stained with **ab28947** NQO1 antibody.

This data was developed using the same antibody clone in a different buffer formulation containing PBS and Azide (**ab28947**).

**Please note:** All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

### Our Abpromise to you: Quality guaranteed and expert technical support

---

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <https://www.abcam.com/abpromise> or contact our technical team.

### Terms and conditions

---

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors