

## Product datasheet

# Anti-PARK7/DJ1 antibody [EP2815Y] - BSA and Azide free ab218373

**KO VALIDATED** Recombinant RabMAB

[6 References](#) [10 Images](#)

### Overview

<b>Product name</b>	Anti-PARK7/DJ1 antibody [EP2815Y] - BSA and Azide free
<b>Description</b>	Rabbit monoclonal [EP2815Y] to PARK7/DJ1 - BSA and Azide free
<b>Host species</b>	Rabbit
<b>Specificity</b>	The mouse and rat recommendation is based on the WB results. We do not guarantee IHC-P for mouse and rat.
<b>Tested applications</b>	<b>Suitable for:</b> IP, ICC/IF, WB, IHC-P, Flow Cyt (Intra)
<b>Species reactivity</b>	<b>Reacts with:</b> Mouse, Rat, Human
<b>Immunogen</b>	Synthetic peptide. This information is proprietary to Abcam and/or its suppliers.
<b>Positive control</b>	WB: Jurkat, HeLa, NIH3T3 or 293T cell lysate. Human fetal brain; Human brain nuclear fraction tissue lysate; Mouse brain and Rat brain tissue lysates. IHC-P: Human Lung and Brain tissue. ICC/IF: PANC-1 and Jurkat cell lines. Flow Cyt (intra): HepG2 cells. IP: Mouse brain lysate.
<b>General notes</b>	<p>ab218373 is the carrier-free version of <a href="#">ab76008</a>.</p> <p>Our <b>carrier-free</b> antibodies are typically supplied in a PBS-only formulation, purified and free of BSA, sodium azide and glycerol. The carrier-free buffer and high concentration allow for increased conjugation efficiency.</p> <p>This conjugation-ready format is designed for use with fluorochromes, metal isotopes, oligonucleotides, and enzymes, which makes them ideal for antibody labelling, functional and cell-based assays, flow-based assays (e.g. mass cytometry) and Multiplex Imaging applications.</p> <p>Use our <b>conjugation kits</b> for antibody conjugates that are ready-to-use in as little as 20 minutes with &lt;1 minute hands-on-time and 100% antibody recovery: available for fluorescent dyes, HRP, biotin and gold.</p> <p>This product is compatible with the Maxpar<sup>®</sup> Antibody Labeling Kit from Fluidigm, without the need for antibody preparation. Maxpar<sup>®</sup> is a trademark of Fluidigm Canada Inc.</p> <p>Our RabMAB<sup>®</sup> technology is a patented hybridoma-based technology for making rabbit monoclonal antibodies. For details on our patents, please refer to <a href="#">RabMAB<sup>®</sup> patents</a>.</p>

### Properties

<b>Form</b>	Liquid
<b>Storage instructions</b>	Shipped at 4°C. Store at +4°C. Do Not Freeze.
<b>Storage buffer</b>	pH: 7.20 Constituent: PBS
<b>Carrier free</b>	Yes
<b>Purity</b>	Protein A purified
<b>Clonality</b>	Monoclonal
<b>Clone number</b>	EP2815Y
<b>Isotype</b>	IgG

## Applications

**The Abpromise guarantee** Our **Abpromise guarantee** covers the use of ab218373 in the following tested applications. The application notes include recommended starting dilutions; optimal dilutions/concentrations should be determined by the end user.

Application	Abreviews	Notes
<b>IP</b>		Use at an assay dependent concentration.
<b>ICC/IF</b>		Use at an assay dependent concentration.
<b>WB</b>		Use at an assay dependent concentration. Predicted molecular weight: 20 kDa.
<b>IHC-P</b>		Use at an assay dependent concentration. Perform heat mediated antigen retrieval with citrate buffer pH 6 before commencing with IHC staining protocol. Perform heat mediated antigen retrieval using 0.01M Sodium Citrate Buffer, pH 6.0 before commencing with IHC staining protocol.  The mouse and rat recommendation is based on the WB results
<b>Flow Cyt (Intra)</b>		Use at an assay dependent concentration. <b>ab199376</b> - Rabbit monoclonal IgG (Low endotoxin, Azide free), is suitable for use as an isotype control with this antibody.

## Target

**Function** Protects cells against oxidative stress and cell death. Plays a role in regulating expression or stability of the mitochondrial uncoupling proteins SLC25A14 and SLC25A27 in dopaminergic neurons of the substantia nigra pars compacta and attenuates the oxidative stress induced by calcium entry into the neurons via L-type channels during pacemaking. Eliminates hydrogen peroxide and protects cells against hydrogen peroxide-induced cell death. May act as an atypical peroxiredoxin-like peroxidase that scavenges hydrogen peroxide. Following removal of a C-terminal peptide, displays protease activity and enhanced cytoprotective action against oxidative stress-induced apoptosis. Stabilizes NFE2L2 by preventing its association with KEAP1 and its subsequent ubiquitination. Binds to OTUD7B and inhibits its deubiquitinating activity. Enhances RELA nuclear translocation. Binds to a number of mRNAs containing multiple copies of GG or CC

motifs and partially inhibits their translation but dissociates following oxidative stress. Required for correct mitochondrial morphology and function and for autophagy of dysfunctional mitochondria. Regulates astrocyte inflammatory responses. Acts as a positive regulator of androgen receptor-dependent transcription. Prevents aggregation of SNCA. Plays a role in fertilization. Has no proteolytic activity. Has cell-growth promoting activity and transforming activity. May function as a redox-sensitive chaperone.

### Tissue specificity

Highly expressed in pancreas, kidney, skeletal muscle, liver, testis and heart. Detected at slightly lower levels in placenta and brain. Detected in astrocytes, Sertoli cells, spermatogonia, spermatids and spermatozoa.

### Involvement in disease

Defects in PARK7 are the cause of Parkinson disease type 7 (PARK7) [MIM:606324]. A neurodegenerative disorder characterized by resting tremor, postural tremor, bradykinesia, muscular rigidity, anxiety and psychotic episodes. PARK7 has onset before 40 years, slow progression and initial good response to levodopa. Some patients may show traits reminiscent of amyotrophic lateral sclerosis-parkinsonism/dementia complex (Guam disease).

### Sequence similarities

Belongs to the peptidase C56 family.

### Post-translational modifications

Sumoylated on Lys-130 by PIAS2 or PIAS4; which is enhanced after ultraviolet irradiation and essential for cell-growth promoting activity and transforming activity.


Cys-106 is easily oxidized to sulfinic acid.

Undergoes cleavage of a C-terminal peptide and subsequent activation of protease activity in response to oxidative stress.

### Cellular localization

Cytoplasm. Nucleus. Mitochondrion. Under normal conditions, located predominantly in the cytoplasm and, to a lesser extent, in the nucleus and mitochondrion. Translocates to the mitochondrion and subsequently to the nucleus in response to oxidative stress and exerts an increased cytoprotective effect against oxidative damage. Detected in tau inclusions in brains from neurodegenerative disease patients.

## Images



**All lanes :** Anti-PARK7/DJ1 antibody [EP2815Y] ([ab76008](#)) at 1/1000 dilution

**Lane 1 :** Wild-type HEK-293T (Human epithelial cell line from embryonic kidney transformed with large T antigen) whole cell lysate

**Lane 2 :** PARK7 knockout HEK-293T (Human epithelial cell line from embryonic kidney transformed with large T antigen) whole cell lysate

**Lane 3 :** HeLa (Human epithelial cell line from cervix adenocarcinoma) whole cell lysate

**Lane 4 :** Human brain nuclear fraction tissue lysate

Lysates/proteins at 20 µg per lane.

### Secondary

**All lanes :** Goat anti-Rabbit IgG H&L (IRDye® 800CW) preadsorbed ([ab216773](#)) at 1/10000 dilution


**Predicted band size:** 20 kDa

**Observed band size:** 24 kDa

This data was developed using [ab76008](#), the same antibody clone in a different buffer formulation.

**Lanes 1-4:** Merged signal (red and green). Green - [ab76008](#) observed at 24 kDa. Red - loading control [ab8245](#) observed at 36 kDa.


[ab76008](#) Anti-PARK7/DJ1 antibody [EP2815Y] was shown to specifically react with PARK7/DJ1 in wild-type HEK-293T cells. Loss of signal was observed when knockout cell line [ab266338](#) (knockout cell lysate [ab257016](#)) was used. Wild-type and PARK7/DJ1 knockout samples were subjected to SDS-PAGE. [ab76008](#) and Anti-GAPDH antibody [6C5] - Loading Control ([ab8245](#)) were incubated overnight at 4°C at 1 in 1000 dilution and 1 in 20000 dilution respectively. Blots were developed with Goat anti-Rabbit IgG H&L (IRDye® 800CW) preadsorbed ([ab216773](#)) and Goat anti-Mouse IgG H&L (IRDye® 680RD) preadsorbed ([ab216776](#)) secondary antibodies at 1 in 20000 dilution for 1 hour at room temperature before imaging.



Immunocytochemistry/ Immunofluorescence - Anti-PARK7/DJ1 antibody [EP2815Y] - BSA and Azide free (ab218373)

This IHC data was generated using the same anti-PARK7 antibody clone, EP2815Y, in a different buffer formulation (cat# [ab76008](#)).


Immunocytochemistry/ Immunofluorescence analysis of Jurkat (Human T cell leukemia T lymphocyte) cells labeling PARK7/DJ1 with Purified [ab76008](#) at 1:500 dilution (0.2 µg/ml). Cells were fixed in 4% Paraformaldehyde and permeabilized with 0.1% tritonX-100. Cells were counterstained with Ab195889 Anti-alpha Tubulin antibody [DM1A] - Microtubule Marker (Alexa Fluor® 594) 1:200 (2.5 µg/ml). Goat anti rabbit IgG (Alexa Fluor® 488, [ab150077](#)) was used as the secondary antibody at 1:1000 (2 µg/ml) dilution. DAPI nuclear counterstain. PBS instead of the primary antibody was used as the secondary antibody only control.



Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-PARK7/DJ1 antibody [EP2815Y] - BSA and Azide free (ab218373)

This IHC data was generated using the same anti-PARK7 antibody clone, EP2815Y, in a different buffer formulation (cat# **ab76008**).


Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) analysis of Human lung cancer tissue sections labeling PARK7/DJ1 with Purified **ab76008** at 1:1000 dilution (0.11 µg/ml). Heat mediated antigen retrieval was performed using **ab93684** (Tris/EDTA buffer, pH 9.0). ImmunoHistoProbe one step HRP Polymer (ready to use) was used as the secondary antibody. Negative control: PBS instead of the primary antibody. Hematoxylin was used as a counterstain.



Flow Cytometry (Intracellular) - Anti-PARK7/DJ1 antibody [EP2815Y] - BSA and Azide free (ab218373)

This IHC data was generated using the same anti-PARK7 antibody clone, EP2815Y, in a different buffer formulation (cat **ab76008**).

Intracellular Flow Cytometry analysis of HepG2 (Human hepatocellular carcinoma epithelial cell) cells labeling PARK7/DJ1 with Purified **ab76008** at 1/20 dilution (10µg/ml) (red). Cells were fixed with 4% Paraformaldehyde. A Goat anti rabbit IgG (Alexa Fluor® 488, **ab150077**) secondary antibody was used at 1/2000. Isotype control - Rabbit monoclonal IgG (Black). Unlabeled control - Cell without incubation with primary antibody and secondary antibody (Blue).



Western blot - Anti-PARK7/DJ1 antibody [EP2815Y]  
- BSA and Azide free (ab218373)

This WB data was generated using the same anti-PARK7 antibody clone, EP2815Y, in a different buffer formulation (cat# [ab76008](#)).

**Lane 1:** Wild-type HAP1 cell lysate (20 µg)

**Lane 2:** PARK7/DJ1 knockout HAP1 cell lysate (20 µg)

**Lane 3:** HeLa cell lysate (20 µg)

**Lane 4:** Human brain tissue lysate (20 µg)

**Lanes 1 - 4:** Merged signal (red and green). Green - [ab76008](#) observed at 24 kDa. Red - loading control, [ab8245](#), observed at 37 kDa.

[ab76008](#) was shown to specifically react with PARK/DJ1 when PARK/DJ1 knockout samples were used. Wild-type and PARK/DJ1 knockout samples were subjected to SDS-PAGE. [ab76008](#) and [ab8245](#) (loading control to GAPDH) were both diluted 1/10 000 and incubated overnight at 4°C. Blots were developed with Goat anti-Rabbit IgG H&L (IRDye® 800CW) preadsorbed ([ab216773](#)) and Goat anti-Mouse IgG H&L (IRDye® 680RD) preadsorbed ([ab216776](#)) secondary antibodies at 1/10 000 dilution for 1 h at room temperature before imaging.



Immunoprecipitation - Anti-PARK7/DJ1 antibody  
[EP2815Y] - BSA and Azide free (ab218373)

This IHC data was generated using the same anti-PARK7 antibody clone, EP2815Y, in a different buffer formulation (cat# [ab76008](#)).

[ab76008](#) (purified) at 1:20 dilution (0.5µg) immunoprecipitating PARK7/DJ1 in Mouse brain lysate.


Lane 1 (input): Mouse brain lysate 10µg

Lane 2 (+): [ab76008](#) & Mouse brain lysate

Lane 3 (-): Rabbit monoclonal IgG ([ab172730](#)) instead of [ab76008](#) in Mouse brain lysate

For western blotting, VeriBlot for IP Detection Reagent (HRP) ([ab131366](#)) was used for detection at 1:1000 dilution.

Blocking and diluting buffer: 5% NFD/MTBST.




Immunocytochemistry/ Immunofluorescence - Anti-PARK7/DJ1 antibody [EP2815Y] - BSA and Azide free (ab218373)

Immunocytochemistry/Immunofluorescence analysis of MCF-7 (human breast carcinoma) labelling PARK7/DJ1 with purified **ab76008** at 1/500. Cells were fixed with 4% PFA and permeabilized with 0.1% Triton X-100. An Alexa Fluor® 488-conjugated goat anti-rabbit IgG (1/1000) was used as the secondary antibody (Ab150077). Nuclei counterstained with DAPI (blue).

Control: PBS only

This data was developed using the same antibody clone in a different buffer formulation containing PBS, BSA, glycerol, and sodium azide (**ab76008**).

**This image was generated using the unpurified version of the product.**




Flow Cytometry (Intracellular) - Anti-PARK7/DJ1 antibody [EP2815Y] - BSA and Azide free (ab218373)

Overlay histogram showing HepG2 cells stained with **ab76008** (red line). The cells were fixed with 4% paraformaldehyde (10 min) and then permeabilized with 0.1% PBS-Tween for 20 min. The cells were then incubated in 1x PBS / 10% normal goat serum / 0.3M glycine to block non-specific protein-protein interactions followed by the antibody (**ab76008**, 1/100 dilution) for 30 min at 22°C. The secondary antibody used was Alexa Fluor® 488 goat anti-rabbit IgG (H+L) (**ab150077**) at 1/2000 dilution for 30 min at 22°C. Isotype control antibody (black line) was rabbit IgG (monoclonal) (1µg/1x10<sup>6</sup> cells) used under the same conditions. Unlabelled sample (blue line) was also used as a control. Acquisition of >5,000 events were collected using a 20mW Argon ion laser (488nm) and 525/30 bandpass filter. This antibody gave a positive signal in HepG2 cells fixed with 80% methanol (5 min)/permeabilized with 0.1% PBS-Tween for 20 min used under the same conditions.

This data was developed using the same antibody clone in a different buffer formulation containing PBS, BSA, glycerol, and sodium azide (**ab76008**).

**This image was generated using the unpurified version of the product.**



Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-PARK7/DJ1 antibody [EP2815Y] - BSA and Azide free (ab218373)

This IHC data was generated using the same anti-PARK7 antibody clone, EP2815Y, in a different buffer formulation (cat# **ab76008**).

**ab76008**, at 1/250 dilution, staining PARK7/DJ1 in human brain by immunohistochemistry using paraffin-embedded tissue.

**This image was generated using the unpurified version of the product.**

Heat mediated antigen retrieval was performed with citrate buffer pH 6 before commencing with IHC staining protocol.

### Why choose a recombinant antibody?



**Research with confidence**  
Consistent and reproducible results



**Long-term and scalable supply**  
Recombinant technology



**Success from the first experiment**  
Confirmed specificity



**Ethical standards compliant**  
Animal-free production

Anti-PARK7/DJ1 antibody [EP2815Y] - BSA and Azide free (ab218373)

**Please note:** All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

### Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <https://www.abcam.com/abpromise> or contact our technical team.

### Terms and conditions



- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors