

Anti-Parvalbumin antibody ab11427

★★★★★ [30 Abreviews](#) [253 References](#) [8 Images](#)

Overview

Product name	Anti-Parvalbumin antibody
Description	Rabbit polyclonal to Parvalbumin
Host species	Rabbit
Tested applications	Suitable for: ICC/IF, IHC-P
Species reactivity	Reacts with: Rat, Human
Immunogen	Full length native protein (purified) corresponding to Rat Parvalbumin. Purified parvalbumin from rat skeletal muscle.
Positive control	ICC/IF: U251, HeLa, C6, and rat cordical cells; IHC-P: Human tonsil, cerebellum and skeletal muscle tissue sections.
General notes	<p>The Life Science industry has been in the grips of a reproducibility crisis for a number of years. Abcam is leading the way in addressing this with our range of recombinant monoclonal antibodies and knockout edited cell lines for gold-standard validation. Please check that this product meets your needs before purchasing.</p> <p>If you have any questions, special requirements or concerns, please send us an inquiry and/or contact our Support team ahead of purchase. Recommended alternatives for this product can be found below, along with publications, customer reviews and Q&As</p>

Properties

Form	Liquid
Storage instructions	Shipped at 4°C. Store at +4°C short term (1-2 weeks). Upon delivery aliquot. Store at -20°C. Avoid freeze / thaw cycle.
Storage buffer	<p>pH: 6.50</p> <p>Preservative: 0.1% Sodium azide</p> <p>Constituents: 2% BSA, 1.62% Sodium phosphate</p>
Purity	Immunogen affinity purified
Clonality	Polyclonal
Isotype	IgG

Applications

The Abpromise guarantee

Our **Abpromise guarantee** covers the use of ab11427 in the following tested applications.

The application notes include recommended starting dilutions; optimal dilutions/concentrations should be determined by the end user.

Application	Abreviews	Notes
ICC/IF	★★★★★ (5)	1/100 - 1/200.
IHC-P	★★★★★ (5)	Use a concentration of 1 µg/ml.

Target


Function

In muscle, parvalbumin is thought to be involved in relaxation after contraction. It binds two calcium ions.

Sequence similarities


Belongs to the parvalbumin family.
Contains 2 EF-hand domains.

Images



Immunocytochemistry/ Immunofluorescence - Anti-Parvalbumin antibody (ab11427)


Immunocytochemistry/Immunofluorescence analysis of U251 cells labeling Parvalbumin (green) with ab11427 at 1/200. F-Actin staining with Phalloidin (red) and nuclei with DAPI (blue). Cells were fixed with formaldehyde and incubated with the primary antibody overnight at 4°C. A DyLight 488-conjugated secondary antibody was used. 60X magnification. Right - negative control.



Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-Parvalbumin antibody (ab11427)


Immunohistochemistry was performed on normal biopsies of deparaffinized Human tonsil tissue. To expose target proteins heat induced antigen retrieval was performed using 10mM sodium citrate (pH6.0) buffer microwaved for 8-15 minutes. Following antigen retrieval tissues were blocked in 3% BSA-PBS for 30 minutes at room temperature. Tissues were then probed at a dilution of 1:100 with a rabbit polyclonal antibody recognizing Parvalbumin ab11427 or without primary antibody (negative control) overnight at 4°C in a humidified chamber. Tissues were washed extensively with PBST and endogenous peroxidase activity was quenched with a peroxidase suppressor. Detection was performed using a biotin-conjugated secondary antibody and SA-HRP followed by colorimetric detection using DAB. Tissues were

counterstained with hematoxylin and prepped for mounting.




Immunocytochemistry/ Immunofluorescence - Anti-Parvalbumin antibody (ab11427)

Immunocytochemistry/Immunofluorescence analysis of HeLa cells labeling Parvalbumin (green) with ab11427 at 1/100. F-Actin staining with Phalloidin (red) and nuclei with DAPI (blue). Cells were fixed with formaldehyde and incubated with the primary antibody overnight at 4°C. A DyLight 488-conjugated secondary antibody was used. 60X magnification. Right - negative control.



Immunocytochemistry/ Immunofluorescence - Anti-Parvalbumin antibody (ab11427)


Immunocytochemistry/Immunofluorescence analysis of C6 (rat glial tumor cell line) cells labeling Parvalbumin (green) with ab11427 at 1/100. F-Actin staining with Phalloidin (red) and nuclei with DAPI (blue). Cells were fixed with formaldehyde and incubated with the primary antibody overnight at 4°C. A DyLight 488-conjugated secondary antibody was used. 60X magnification. Right - negative control.



Immunocytochemistry/ Immunofluorescence - Anti-Parvalbumin antibody (ab11427)

Image is courtesy of an AbReview submitted by Ms Babben Tinner.

Immunocytochemical immunofluorescence analysis of 4% PFA & 0.2% Picric acid fixed rat cordical cells in culture, labelling parvalbumin with ab11427 at a dilution of 1/500 incubated for 12 hours at 4°C in 10mM PBS & 0.03% Triton X diluent blend. The secondary was a Donkey anti-Rabbit polyclonal Alexa Fluor® 488 conjugate at 1/200.




Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-Parvalbumin antibody (ab11427)

Image courtesy of an anonymous Abreview.


ab11427 staining Parvalbumin in human brain tissue sections by Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections).

Tissue was fixed in formaldehyde and a heat mediated antigen retrieval step was performed using EDTA pH 8.0 for 20 minutes at 100°C. Samples were then incubated with ab11427 at a 1/1000 dilution for 20 minutes at 25°C. The secondary used was an undiluted HRP conjugated goat anti-mouse/ rabbit IgG.



Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-Parvalbumin antibody (ab11427)

Immunohistochemistry was performed on normal biopsies of deparaffinized Human cerebellum tissue. To expose target proteins heat induced antigen retrieval was performed using 10mM sodium citrate (pH6.0) buffer microwaved for 8-15 minutes. Following antigen retrieval tissues were blocked in 3% BSA-PBS for 30 minutes at room temperature. Tissues were then probed at a dilution of 1:100 with a rabbit polyclonal antibody recognizing Parvalbumin [ab114227](#) or without primary antibody (negative control) overnight at 4°C in a humidified chamber. Tissues were washed extensively with PBST and endogenous peroxidase activity was quenched with a peroxidase suppressor. Detection was performed using a biotin-conjugated secondary antibody and SA-HRP followed by colorimetric detection using DAB. Tissues were counterstained with hematoxylin and prepped for mounting.



Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-Parvalbumin antibody (ab11427)

Immunohistochemistry was performed on normal biopsies of deparaffinized Human skeletal muscle tissue. To expose target proteins heat induced antigen retrieval was performed using 10mM sodium citrate (pH6.0) buffer microwaved for 8-15 minutes. Following antigen retrieval tissues were blocked in 3% BSA-PBS for 30 minutes at room temperature. Tissues were then probed at a dilution of 1:20 with a rabbit polyclonal antibody recognizing Parvalbumin ab11427 or without primary antibody (negative control) overnight at 4°C in a humidified chamber. Tissues were washed extensively with PBST and endogenous peroxidase activity was quenched with a peroxidase suppressor. Detection was performed using a biotin-conjugated secondary antibody and SA-HRP

followed by colorimetric detection using DAB. Tissues were counterstained with hematoxylin and prepped for mounting.

Please note: All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <https://www.abcam.com/abpromise> or contact our technical team.

Terms and conditions

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors