

Product datasheet

Anti-PD-L1 antibody [CAL10] - Mouse IgG1 (Chimeric) - BSA and Azide free ab279304

KO VALIDATED Recombinant

7 Images

Overview

Product name	Anti-PD-L1 antibody [CAL10] - Mouse IgG1 (Chimeric) - BSA and Azide free
Description	Mouse monoclonal [CAL10] to PD-L1 - Chimeric – BSA and Azide free
Host species	Mouse
Tested applications	Suitable for: WB, Flow Cyt (Intra), ICC/IF, IHC-P
Species reactivity	Reacts with: Human
Immunogen	Synthetic peptide. This information is proprietary to Abcam and/or its suppliers.
Positive control	WB: PD-L1 stably expressed CHO whole cell lysate. Human placenta tissue lysate. NCI-H1299 whole cell lysate. ICC: PD-L1 stably expressed CHO cells. Flow Cyt (intra): PD-L1 stably expressed CHO cells. IHC-P: Human tonsil tissue.
General notes	<p>ab279304 is the carrier free version of ab279292.</p> <p>This mouse monoclonal chimeric antibody has been engineered from a RabMAb parent antibody (ab237726). By necessity, some rabbit sequence is retained as part of the variable domain. When multiplexing with other rabbit-derived antibodies, using cross absorbed Fc-reactive secondary antibodies are recommended.</p> <p>Our carrier-free antibodies are typically supplied in a PBS-only formulation, purified and free of BSA, sodium azide and glycerol. The carrier-free buffer and high concentration allow for increased conjugation efficiency.</p> <p>This conjugation-ready format is designed for use with fluorochromes, metal isotopes, oligonucleotides, and enzymes, which makes them ideal for antibody labelling, functional and cell-based assays, flow-based assays (e.g. mass cytometry) and Multiplex Imaging applications.</p> <p>Use our conjugation kits for antibody conjugates that are ready-to-use in as little as 20 minutes with <1 minute hands-on-time and 100% antibody recovery: available for fluorescent dyes, HRP, biotin and gold.</p> <p>This product is compatible with the Maxpar[®] Antibody Labeling Kit from Fluidigm, without the need for antibody preparation. Maxpar[®] is a trademark of Fluidigm Canada Inc.</p>

Properties

Form	Liquid
Storage instructions	Shipped at 4°C. Store at +4°C. Do Not Freeze.
Storage buffer	Constituent: 100% PBS
Carrier free	Yes
Purity	Protein A purified
Clonality	Monoclonal
Clone number	CAL10
Isotype	IgG1

Applications


The Abpromise guarantee Our **Abpromise guarantee** covers the use of ab279304 in the following tested applications. The application notes include recommended starting dilutions; optimal dilutions/concentrations should be determined by the end user.

Application	Abreviews	Notes
WB		Use at an assay dependent concentration.
Flow Cyt (Intra)		Use at an assay dependent concentration.
ICC/IF		Use at an assay dependent concentration.
IHC-P		Use at an assay dependent concentration. Perform heat mediated antigen retrieval with citrate buffer pH 6 before commencing with IHC staining protocol.

Target

Function	Involved in the costimulatory signal, essential for T-cell proliferation and production of IL10 and IFNG, in an IL2-dependent and a PDCD1-independent manner. Interaction with PDCD1 inhibits T-cell proliferation and cytokine production.
Tissue specificity	Highly expressed in the heart, skeletal muscle, placenta and lung. Weakly expressed in the thymus, spleen, kidney and liver. Expressed on activated T- and B-cells, dendritic cells, keratinocytes and monocytes.
Sequence similarities	Belongs to the immunoglobulin superfamily. BTN/MOG family. Contains 1 Ig-like C2-type (immunoglobulin-like) domain. Contains 1 Ig-like V-type (immunoglobulin-like) domain.
Cellular localization	Cell membrane and Endomembrane system.

Images



Western blot - Anti-PD-L1 antibody [CAL10] - Mouse IgG1 (Chimeric) - BSA and Azide free (ab279304)

All lanes : Anti-PD-L1 antibody [CAL10] - Mouse IgG1 (Chimeric) ([ab279292](#)) at 1/1000 dilution

Lane 1 : Wild-type A549 Control IFN-gamma (0 ng/mL, 48 h), [ab255450](#)

Lane 2 : Wild-type A549 Treated IFN-gamma (100 ng/mL, 48 h), [ab255450](#)

Lane 3 : CD274 knockout A549 Control IFN-gamma (0 ng/mL, 48 h), [ab267055](#)

Lane 4 : CD274 knockout A549 Treated IFN-gamma (100 ng/mL, 48 h), [ab267055](#)


Lysates/proteins at 20 µg per lane.

Performed under reducing conditions.

Observed band size: 45 kDa

This data was produced using [ab279292](#), the same clone in a different formulation.

False colour image of Western blot: Anti-PD-L1 antibody [CAL10] - Mouse IgG1 staining at 1/1000 dilution, shown in green; Rabbit anti-Vinculin antibody ([ab219649](#)) loading control staining at 1/1000 dilution, shown in red. In Western blot, [ab279292](#) was shown to bind specifically to PD-L1. A band was observed at 45 kDa in wild-type A549 cell lysates with no signal observed at this size in Cd274 knockout cell line. To generate this image, wild-type and Cd274 knockout A549 cell lysates were analysed. First, samples were run on an SDS-PAGE gel then transferred onto a nitrocellulose membrane. Membranes were blocked in 3 % milk in TBS-0.1 % Tween® 20 (TBS-T) before incubation with primary antibodies overnight at 4 °C. Blots were washed four times in TBS-T, incubated with secondary antibodies for 1 h at room temperature, washed again four times then imaged. Secondary antibodies used were Goat anti-Mouse IgG H&L 800CW and Goat anti-Rabbit IgG H&L 680RD at 1/20000 dilution.



All lanes : Anti-PD-L1 antibody [CAL10] - Mouse IgG2a (Chimeric) ([ab279293](#)) at 1/1000 dilution

Lane 1 : Wild-type A549 Treated IFN-gamma (100 ng/mL, 48 h) cell lysate

Lane 2 : Wild-type A549 Vehicle Control IFN-gamma (0 ng/mL, 48 h) cell lysate

Lane 3 : CD274 knockout A549 Treated IFN-gamma (100 ng/mL, 48 h) cell lysate

Lane 4 : CD274 knockout A549 Vehicle Control IFN-gamma (0 ng/mL, 48 h) cell lysate


Lane 5 : Human Placenta cell lysate

Lysates/proteins at 20 µg per lane.

Performed under reducing conditions.

Observed band size: 45-65 kDa

False colour image of Western blot: Anti-PD-L1 antibody [CAL10] - Mouse IgG2a staining at 1/1000 dilution, shown in green; Rabbit Anti-GAPDH antibody [EPR16891] ([ab181602](#)) loading control staining at 1/20000 dilution, shown in red. In Western blot, [ab279293](#) was shown to bind specifically to PD-L1. A band was observed at 45-65 kDa in treated wild-type A549 cell lysates with no signal observed at this size in Cd274 knockout cell line [ab267054](#) (knockout cell lysate [ab256831](#)). To generate this image, wild-type and Cd274 knockout A549 cell lysates were analysed. First, samples were run on an SDS-PAGE gel then transferred onto a nitrocellulose membrane. Membranes were blocked in fluorescent western blot (TBS-based) blocking solution before incubation with primary antibodies overnight at 4 °C. Blots were washed four times in TBS-T, incubated with secondary antibodies for 1 h at room temperature, washed again four times then imaged. Secondary antibodies used were Goat anti-Mouse IgG H&L 800CW and Goat anti-Rabbit IgG H&L 680RD at 1/20000 dilution.



Western blot - Anti-PD-L1 antibody [CAL10] - Mouse IgG1 (Chimeric) - BSA and Azide free (ab279304)

All lanes : Anti-PD-L1 antibody [CAL10] - Mouse IgG1 (Chimeric) ([ab279292](#)) at 1/1000 dilution

Lane 1 : Wild-type A549 Treated IFN-gamma (100 ng/ml) for 48 hours cell lysate


Lane 2 : CD274 knockout A549 Treated IFN-gamma (100 ng/ml) for 48 hours cell lysate

Lysates/proteins at 20 µg per lane.

Performed under reducing conditions.

Observed band size: 48 kDa

False colour image of Western blot: Anti-PD-L1 antibody [CAL10] - Mouse IgG1 staining at 1/1000 dilution, shown in green; Rabbit anti-alpha Tubulin antibody [EP1332Y] ([ab52866](#)) loading control staining at 1/20000 dilution, shown in red. In Western blot, [ab279292](#) was shown to bind specifically to PD-L1. A band was observed at 48 kDa in treated wild-type A549 cell lysates with no signal observed at this size in Cd274 knockout cell line [ab267054](#) (knockout cell lysate [ab256831](#)). To generate this image, wild-type and Cd274 knockout A549 cell lysates were analysed. First, samples were run on an SDS-PAGE gel then transferred onto a nitrocellulose membrane. Membranes were blocked in 3 % milk in TBS-0.1 % Tween[®] 20 (TBS-T) before incubation with primary antibodies overnight at 4°C. Blots were washed four times in TBS-T, incubated with secondary antibodies for 1 h at room temperature, washed again four times then imaged. Secondary antibodies used were Goat anti-Mouse IgG H&L (IRDye[®] 800CW) preabsorbed ([ab216772](#)) and Goat anti-Rabbit IgG H&L (IRDye[®] 680RD) preabsorbed ([ab216777](#)) at 1/20000 dilution.



All lanes : Anti-PD-L1 antibody [CAL10] - Mouse IgG1 (Chimeric) ([ab279292](#)) at 1/1000 dilution

Lane 1 : CHO-S (Chinese hamster ovary epithelial cell) whole cell lysate

Lane 2 : CHO-PD-L1 (PD-L1 stably expressed Chinese hamster ovary epithelial cell) whole cell lysate

Lane 3 : Human placenta tissue lysate

Lane 4 : NCI-H1299 (human lung carcinoma epithelial cell), whole cell lysate

Lysates/proteins at 20 µg per lane.

Secondary


All lanes : Peroxidase-Conjugated Goat anti-Mouse IgG (H+L) at 1/5000 dilution

Observed band size: 40-60 kDa

Exposure time: 15 seconds

This data was produced using [ab279292](#), the same clone in a different formulation.

Blocking/Dilution buffer: 5% NFDM/TBST.




This data was produced using **ab279292**, the same clone in a different formulation.

IHC image of PD-L1 staining in a section of formalin-fixed paraffin-embedded normal human tonsil* performed on a Leica BOND™ system using the standard protocol F.

The section was pre-treated using heat mediated antigen retrieval with sodium citrate buffer (pH6, epitope retrieval solution 1) for 20mins. The section was then incubated with **ab279292**, 1ug/ml, for 15 mins at room temperature. A rabbit anti-mouse IgG1, **ab125913**, was added for 8 mins at room temperature and detected using an HRP conjugated goat anti-rabbit compact polymer system. DAB was used as the chromogen. The section was then counterstained with haematoxylin and mounted with DPX. The inset secondary-only control image is taken from an identical assay without primary antibody.

For other IHC staining systems (automated and non-automated) customers should optimize variable parameters such as antigen retrieval conditions, primary antibody concentration and antibody incubation times.

*Tissue obtained from the Human Research Tissue Bank, supported by the NIHR Cambridge Biomedical Research Centre




Immunocytochemistry/ Immunofluorescence - Anti-PD-L1 antibody [CAL10] - Mouse IgG1 (Chimeric) - BSA and Azide free (ab279304)

This data was produced using [ab279292](#), the same clone in a different formulation.

Immunocytochemical analysis of 4% paraformaldehyde-fixed, 0.1% Triton X-100-fixed permeabilized CHO-PD-L1 cells labeling PD-L1 with [ab279292](#) at 1/100 dilution, followed by [ab150113](#) Goat Anti-Mouse IgG H&L (Alexa Fluor® 488) antibody at 1/1000 dilution (Green). [ab179513](#) Anti-beta Tubulin rabbit monoclonal antibody was used to counterstain tubulin at 1/200 dilution, followed by [ab150080](#) Goat Anti-Rabbit IgG H&L (Alexa Fluor® 594) at a 1/500 dilution (Red). The nuclear counterstain was DAPI (Blue). Confocal image showing membranous and cytoplasmic staining in CHO-PD-L1 cells.

Negative control 1: [ab279292](#) at a 1/100 dilution followed by [ab150080](#) at a 1/200 dilution.

Negative control 2: [ab179513](#) at a 1/200 dilution followed by [ab150157](#) at a 1/1000 dilution.



Flow Cytometry (Intracellular) - Anti-PD-L1 antibody [CAL10] - Mouse IgG1 (Chimeric) - BSA and Azide free (ab279304)

This data was produced using [ab279292](#), the same clone in a different formulation.

Flow cytometric analysis of 4% paraformaldehyde-fixed, 90% methanol-permeabilized CHO-s (Chinese hamster ovary epithelial cell, Blue) / CHO-PDL1 (PD-L1 stably expressed Chinese hamster ovary epithelial cell, Red) labelling PD-L1 with [ab279292](#) at 1/50 dilution (0.1 µg).

Goat Anti-Mouse IgG (Alexa Fluor® 488, [ab150113](#)) at 1/2000 dilution was used as the secondary antibody.

Please note: All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you

- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <https://www.abcam.com/abpromise> or contact our technical team.

Terms and conditions

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors