

Product datasheet

PE Anti-CD38 antibody [90] ab23308

1 References

Overview

Product name	PE Anti-CD38 antibody [90]
Description	PE Rat monoclonal [90] to CD38
Host species	Rat
Conjugation	PE. Ex: 488nm, Em: 575nm
Specificity	This antibody recognises the murine CD38 cell surface antigen, a 42-46kD glycoprotein expressed by B lymphocytes, a proportion of peripheral T lymphocytes and by peritoneal macrophages. In the mouse CD38 is expressed by follicular B cells, is down regulated on germinal centre B cells, and not expressed by mature plasma cells. This pattern of expression contrasts strikingly with that seen in the human.
Tested applications	Suitable for: Flow Cyt
Species reactivity	Reacts with: Mouse
Immunogen	Tissue, cells or virus corresponding to Mouse CD38. Pre-B cells derived from IL-7 dependent bone marrow cultures. (Mouse).
General notes	<p>Purified IgG conjugated to R. Phycoerythrin (RPE).</p> <p>The Life Science industry has been in the grips of a reproducibility crisis for a number of years. Abcam is leading the way in addressing this with our range of recombinant monoclonal antibodies and knockout edited cell lines for gold-standard validation. Please check that this product meets your needs before purchasing.</p> <p>If you have any questions, special requirements or concerns, please send us an inquiry and/or contact our Support team ahead of purchase. Recommended alternatives for this product can be found below, along with publications, customer reviews and Q&As</p>

Properties

Form	Liquid
Storage instructions	Shipped at 4°C. Store at +4°C.
Storage buffer	Preservative: 0.09% Sodium azide Constituents: PBS, 0.5% BSA
Purity	Protein A purified
Clonality	Monoclonal

Clone number	90
Myeloma	P3x63-Ag8.653
Isotype	IgG2a

Applications

The Abpromise guarantee Our **Abpromise guarantee** covers the use of ab23308 in the following tested applications. The application notes include recommended starting dilutions; optimal dilutions/concentrations should be determined by the end user.

Application	Abreviews	Notes
Flow Cyt		Use at an assay dependent concentration.

Target

Function	Synthesizes cyclic ADP-ribose, a second messenger for glucose-induced insulin secretion. Also has cADPr hydrolase activity. Also moonlights as a receptor in cells of the immune system.
Tissue specificity	Expressed at high levels in pancreas, liver, kidney, brain, testis, ovary, placenta, malignant lymphoma and neuroblastoma.
Sequence similarities	Belongs to the ADP-ribosyl cyclase family.
Developmental stage	Preferentially expressed at both early and late stages of the B and T-cell maturation. It is also detected on erythroid and myeloid progenitors in bone marrow, where the level of surface expression was shown to decrease during differentiation of blast-forming unit E to colony-forming unit E.
Cellular localization	Membrane.
Form	There are 2 isoforms produced by alternative splicing.

Please note: All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <https://www.abcam.com/abpromise> or contact our technical team.

Terms and conditions

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors