abcam

Product datasheet

PE/DyLight™ 594 Anti-CD10 antibody [MEM-78] ab239327

1 Image

Overview

Product name PE/DyLight™ 594 Anti-CD10 antibody [MEM-78]

Description PE/DyLight[™] 594 Mouse monoclonal [MEM-78] to CD10

Host species Mouse

Conjugation PE/DyLight™ 594

Tested applications Suitable for: Flow Cyt
Species reactivity Reacts with: Human

Immunogen Tissue, cells or virus corresponding to Human CD10. NALM-6 human pre-B cell line.

Positive control Flow Cyt: Human peripheral blood.

General notes

The Life Science industry has been in the grips of a reproducibility crisis for a number of years.

Abcam is leading the way in addressing this with our range of recombinant monoclonal antibodies and knockout edited cell lines for gold-standard validation. Please check that this product meets

your needs before purchasing.

If you have any questions, special requirements or concerns, please send us an inquiry and/or contact our Support team ahead of purchase. Recommended alternatives for this product can be

found below, along with publications, customer reviews and Q&As

Properties

Form Liquid

Storage instructions Shipped at 4°C. Store at +4°C. Store In the Dark.

Storage buffer pH: 7.4

Preservative: 0.0975% Sodium azide

Constituent: PBS

Purity Size exclusion

Purification notes The purified antibody is conjugated with tandem dye PE-DyLight™ 594 (PE-DL594) under

optimum conditions. The conjugate is purified by size-exclusion chromatography and adjusted for

direct use. No reconstitution is necessary.

Clonality Monoclonal
Clone number MEM-78

1

Applications

The Abpromise guarantee Our Abpromise guarantee cover

Our **Abpromise guarantee** covers the use of ab239327 in the following tested applications.

The application notes include recommended starting dilutions; optimal dilutions/concentrations should be determined by the end user.

Application	Abreviews	Notes
Flow Cyt		Use 4µl for 10 ⁶ cells. (or 100 µl of whole blood)

Target

Function Thermolysin-like specificity, but is almost confined on acting on polypeptides of up to 30 amino


acids. Biologically important in the destruction of opioid peptides such as Met- and Leuenkephalins by cleavage of a Gly-Phe bond. Able to cleave angiotensin-1, angiotensin-2 and angiotensin 1-9. Involved in the degradation of atrial natriuretic factor (ANF). Displays UV-

inducible elastase activity toward skin preelastic and elastic fibers.

Sequence similarities Belongs to the peptidase M13 family.

Cellular localization Cell membrane.

Images



Flow cytometric analysis of human peripheral blood lableing CD10 with ab239327. Surface staining. Gated on leukocytes.

Please note: All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

Our Abpromise to you: Quality guaranteed and expert technical support

• Replacement or refund for products not performing as stated on the datasheet

- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit https://www.abcam.com/abpromise or contact our technical team.

Terms and conditions

• Guarantee only valid for products bought direct from Abcam or one of our authorized distributors