

Product datasheet

Anti-Pellino 1 antibody [EPR19302] ab199336

Recombinant **RabMAb**

[7 References](#) [9 Images](#)

Overview

Product name	Anti-Pellino 1 antibody [EPR19302]
Description	Rabbit monoclonal [EPR19302] to Pellino 1
Host species	Rabbit
Tested applications	Suitable for: WB, IP, ICC/IF
Species reactivity	Reacts with: Mouse, Rat, Human, Recombinant fragment
Immunogen	Recombinant full length protein. This information is proprietary to Abcam and/or its suppliers.
Positive control	WB: Human PELI1 full length recombinant protein; Human fetal liver, fetal heart and fetal kidney lysates; PC-12, NIH/3T3, Ramos, HeLa, RAW 264.7 (treated with 1 µg/ml LPS for 6 hours) and PC-12 (treated with 1 µg/ml LPS for 6 hours) whole cell lysates. ICC/IF: HeLa and Ramos cells. IP: Ramos whole cell lysate.
General notes	<p>This product is a recombinant monoclonal antibody, which offers several advantages including:</p> <ul style="list-style-type: none"> - High batch-to-batch consistency and reproducibility - Improved sensitivity and specificity - Long-term security of supply - Animal-free production <p>For more information see here.</p> <p>Our RabMAb[®] technology is a patented hybridoma-based technology for making rabbit monoclonal antibodies. For details on our patents, please refer to RabMAb[®] patents.</p>

Properties

Storage instructions	Shipped at 4°C. Store at +4°C short term (1-2 weeks). Upon delivery aliquot. Store at -20°C long term. Avoid freeze / thaw cycle.
Storage buffer	<p>pH: 7.2</p> <p>Preservative: 0.01% Sodium azide</p> <p>Constituents: 59% PBS, 0.05% BSA, 40% Glycerol</p>
Purity	Protein A purified
Clonality	Monoclonal
Clone number	EPR19302
Isotype	IgG

Applications

The Abpromise guarantee

Our **Abpromise guarantee** covers the use of ab199336 in the following tested applications.

The application notes include recommended starting dilutions; optimal dilutions/concentrations should be determined by the end user.

Application	Abreviews	Notes
WB		1/1000. Detects a band of approximately 46 kDa (predicted molecular weight: 46 kDa).
IP		1/30.
ICC/IF		1/100.

Target


Function

Scaffold protein involved in the IL-1 signaling pathway via its interaction with the complex containing IRAK kinases and TRAF6. Required for NF-kappa-B activation and IL-8 gene expression in response to IL-1.

Sequence similarities

Belongs to the pellino family.

Images



All lanes : Anti-Pellino 1 antibody [EPR19302] (ab199336) at 1/5000 dilution

Lane 1 : Human PELI1 full length recombinant protein

Lane 2 : Human PELI2 full length recombinant protein

Lane 3 : Human PELI3 full length recombinant protein

Lysates/proteins at 0.01 µg per lane.

Secondary

All lanes : Goat Anti-Rabbit IgG H&L (HRP) (**ab97051**) at 1/100000 dilution


Predicted band size: 46 kDa

Observed band size: 46 kDa

Exposure time: 3 minutes

Blocking/Dilution buffer: 5% NFDM/TBST.

Human PELI 1 full length recombinant protein contain aa1-418 with a His-Tag®. Human PELI 2 full length recombinant protein contain aa1-420 with a His-Tag®. Human PELI3 full length recombinant protein contain aa1-469 with a His-Tag®. All three recombinant human fragment proteins were made in-house.



Immunocytochemistry/ Immunofluorescence - Anti-Pellino 1 antibody [EPR19302] (ab199336)


Immunofluorescent analysis of 4% paraformaldehyde-fixed, 0.1% Triton X-100 permeabilized HeLa (Human epithelial cell line from cervix adenocarcinoma) cells labeling Pellino 1 with ab199336 at 1/100 dilution, followed by Goat Anti-Rabbit IgG (Alexa Fluor® 488) (**ab150077**) secondary antibody at 1/1000 dilution (green). Confocal image showing cytoplasmic and nuclear staining on HeLa cell line. The nuclear counter stain is DAPI (blue).

Tubulin is detected with Anti-alpha Tubulin antibody [EPR19302] - Loading Control (**ab7291**) at 1/1000 dilution and Goat Anti-Mouse IgG H&L (Alexa Fluor® 594) preadsorbed (**ab150120**) at 1/1000 dilution (red).

The negative controls are as follows:

-ve control 1: ab199336 at 1/100 dilution followed by **ab150120** at 1/1000 dilution.

-ve control 2: **ab7291** at 1/1000 dilution followed by **ab150077** at 1/1000 dilution.



Immunoprecipitation - Anti-Pellino 1 antibody [EPR19302] (ab199336)


Lane 1 (input): Mouse brain lysate, 10µg

Lane 2 (+): Mouse brain lysate

Lane 3 (-): Rabbit monoclonal IgG (**ab172730**) instead of ab199336 in Mouse brain lysate

Ab199336 immunoprecipitating Pellino in Mouse brain lysate. For western blotting, primary antibody used was ab199336 at 1:500 dilution. VeriBlot for IP Detection Reagent (HRP) (**ab131366**), was used for detection at 1/1000 dilution.

Blocking and diluting buffer: 5% NFDM/TBST



Western blot - Anti-Pellino 1 antibody [EPR19302] (ab199336)

All lanes : Anti-Pellino 1 antibody [EPR19302] (ab199336) at 1/1000 dilution

Lane 1 : Human fetal liver lysate

Lane 2 : Human fetal heart lysate

Lane 3 : Human fetal kidney lysate

Lysates/proteins at 10 µg per lane.

Secondary


All lanes : Goat Anti-Rabbit IgG Peroxidase Conjugate, specific to the non-reduced form of IgG at 1/10000 dilution

Predicted band size: 46 kDa

Observed band size: 46 kDa

Exposure time: 3 minutes

Blocking/Dilution buffer: 5% NFDM/TBST.



Western blot - Anti-Pellino 1 antibody [EPR19302] (ab199336)

All lanes : Anti-Pellino 1 antibody [EPR19302] (ab199336) at 1/1000 dilution

Lane 1 : PC-12 (Rat adrenal gland pheochromocytoma cell line) whole cell lysate

Lane 2 : NIH/3T3 (Mouse embryonic fibroblast cell line) whole cell lysate

Lysates/proteins at 10 µg per lane.

Secondary


All lanes : Goat Anti-Rabbit IgG H&L (HRP) ([ab97051](#)) at 1/10000 dilution

Predicted band size: 46 kDa

Observed band size: 46 kDa

Exposure time: 1 minute

Blocking/Dilution buffer: 5% NFDM/TBST.



Western blot - Anti-Pellino 1 antibody [EPR19302]
(ab199336)

All lanes : Anti-Pellino 1 antibody [EPR19302] (ab199336) at 1/1000 dilution

Lane 1 : Ramos (Human Burkitt's lymphoma cell line) whole cell lysate

Lane 2 : HeLa (Human epithelial cell line from cervix adenocarcinoma) whole cell lysate

Lane 3 : Untreated RAW 264.7 (Mouse macrophage cell line transformed with Abelson murine leukemia virus) whole cell lysate

Lane 4 : RAW 264.7 (Mouse macrophage cell line transformed with Abelson murine leukemia virus) treated with 1 µg/ml LPS for 6 hours whole cell lysate

Lane 5 : Untreated PC-12 (Rat adrenal gland pheochromocytoma cell line) whole cell lysate

Lane 6 : PC-12 (Rat adrenal gland pheochromocytoma cell line) treated with 1 µg/ml LPS for 6 hours whole cell lysate

Lysates/proteins at 20 µg per lane.

Secondary

All lanes : Goat Anti-Rabbit IgG H&L (HRP) ([ab97051](#)) at 1/100000 dilution


Predicted band size: 46 kDa

Observed band size: 46 kDa

Exposure time: 3 minutes

Blocking/Dilution buffer: 5% NFDM/TBST.

The expression of Pellino 1 can be induced by LPS. (PMID: 21204785).



Immunocytochemistry/ Immunofluorescence - Anti-Pellino 1 antibody [EPR19302] (ab199336)


Immunofluorescent analysis of 4% paraformaldehyde-fixed, 0.1% Triton X-100 permeabilized Ramos (Human Burkitt's lymphoma cell line) cells labeling Pellino 1 with ab199336 at 1/100 dilution, followed by Goat Anti-Rabbit IgG (Alexa Fluor® 488) (**ab150077**) secondary antibody at 1/1000 dilution (green). Confocal image showing cytoplasmic and nuclear staining on Ramos cell line. The nuclear counter stain is DAPI (blue).

Tubulin is detected with Anti-alpha Tubulin antibody [EPR19302] - Loading Control (**ab7291**) at 1/1000 dilution and Goat Anti-Mouse IgG H&L (Alexa Fluor® 594) preadsorbed (**ab150120**) at 1/1000 dilution (red).

The negative controls are as follows:

-ve control 1: ab199336 at 1/100 dilution followed by **ab150120** at 1/1000 dilution.

-ve control 2: **ab7291** at 1/1000 dilution followed by **ab150077** at 1/1000 dilution.



Immunoprecipitation - Anti-Pellino 1 antibody [EPR19302] (ab199336)

Pellino 1 was immunoprecipitated from 1mg of Ramos (Human Burkitt's lymphoma cell line) whole cell lysate with ab199336 at 1/30 dilution. Western blot was performed from the immunoprecipitate using ab199336 at 1/1000 dilution. VeriBlot for IP Detection Reagent (HRP) (**ab131366**), was used for detection at 1/10000 dilution.

Lane 1: Ramos whole cell lysate 10µg (Input).

Lane 2: ab199336 IP in Ramos whole cell lysate.

Lane 3: Rabbit IgG, monoclonal [EPR19302] - Isotype Control (**ab172730**) instead of ab199336 in Ramos whole cell lysate.

Blocking and dilution buffer and concentration: 5% NFDN/TBST.

Exposure time: (A): 30 seconds; (B): 3 minutes.

Why choose a recombinant antibody?

 <p>Research with confidence Consistent and reproducible results</p>	 <p>Long-term and scalable supply Recombinant technology</p>
 <p>Success from the first experiment Confirmed specificity</p>	 <p>Ethical standards compliant Animal-free production</p>

Anti-Pellino 1 antibody [EPR19302] (ab199336)

Please note: All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <https://www.abcam.com/abpromise> or contact our technical team.

Terms and conditions

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors