abcam

Product datasheet

Anti-Phospholipase C gamma 1/PLC-gamma-1 (phospho Y783) antibody [EPR2611Y] - BSA and Azide free ab248714



2 Images

Overview

Anti-Phospholipase C gamma 1/PLC-gamma-1 (phospho Y783) antibody [EPR2611Y] - BSA **Product name**

and Azide free

Rabbit monoclonal [EPR2611Y] to Phospholipase C gamma 1/PLC-gamma-1 (phospho Y783) -**Description**

BSA and Azide free

Host species Rabbit

Suitable for: WB **Tested applications**

Unsuitable for: Flow Cyt,ICC/IF,IHC-P or IP

Species reactivity Reacts with: Human

Immunogen Synthetic peptide. This information is proprietary to Abcam and/or its suppliers.

General notes ab248714 is the carrier-free version of ab134186.

> Our carrier-free antibodies are typically supplied in a PBS-only formulation, purified and free of BSA, sodium azide and glycerol. The carrier-free buffer and high concentration allow for increased conjugation efficiency.

This conjugation-ready format is designed for use with fluorochromes, metal isotopes, oligonucleotides, and enzymes, which makes them ideal for antibody labelling, functional and cellbased assays, flow-based assays (e.g. mass cytometry) and Multiplex Imaging applications.

Use our conjugation kits for antibody conjugates that are ready-to-use in as little as 20 minutes with <1 minute hands-on-time and 100% antibody recovery: available for fluorescent dyes, HRP, biotin and gold.

This product is compatible with the Maxpar® Antibody Labeling Kit from Fluidigm, without the need for antibody preparation. Maxpar® is a trademark of Fluidigm Canada Inc.

This product is a recombinant monoclonal antibody, which offers several advantages including:

- High batch-to-batch consistency and reproducibility
- Improved sensitivity and specificity
- Long-term security of supply
- Animal-free production

For more information see here.

Our RabMAb[®] technology is a patented hybridoma-based technology for making rabbit monoclonal antibodies. For details on our patents, please refer to **RabMAb**[®] **patents**.

Mouse, Rat: We have preliminary internal testing data to indicate this antibody may not react with these species. Please contact us for more information.

Properties

Form Liquid

Storage instructions Shipped at 4°C. Store at +4°C. Do Not Freeze.

Storage buffer pH: 7.2

Constituent: PBS

Carrier free Yes

Purity Protein A purified

Clonality Monoclonal
Clone number EPR2611Y

Isotype IgG

Applications

The Abpromise guarantee Our Abpromise guarantee covers the use of ab248714 in the following tested applications.

The application notes include recommended starting dilutions; optimal dilutions/concentrations should be determined by the end user.

Application	Abreviews	Notes
WB		Use at an assay dependent concentration. Detects a band of approximately 150 kDa (predicted molecular weight: 148 kDa).

Application notes Is unsuitable for Flow Cyt,ICC/IF,IHC-P or IP.

Target

Function Plays a role in actin reorganization and cell migration. The production of the second messenger

molecules diacylglycerol (DAG) and inositol 1,4,5-trisphosphate (IP3) is mediated by activated phosphatidylinositol-specific phospholipase C enzymes. Major substrate for heparin-binding

growth factor 1 (acidic fibroblast growth factor)-activated tyrosine kinase.

Sequence similarities Contains 1 C2 domain.

Contains 1 EF-hand domain. Contains 2 PH domains.

Contains 1 PI-PLC X-box domain.
Contains 1 PI-PLC Y-box domain.

Contains 2 SH2 domains. Contains 1 SH3 domain.

Domain The SH3 domain mediates interaction with CLNK (By similarity). The SH3 domain also mediates

interaction with RALGPS1.

Post-translational modifications

The receptor-mediated activation of PLC-gamma-1 and PLC-gamma-2 involves their

phosphorylation by tyrosine kinases in response to ligation of a variety of growth factor receptors

and immune system receptors. May be dephosphorylated by PTPRJ.


Ubiquitinated by CBLB in activated T-cells.

Cellular localization

Cell projection > lamellipodium. Cell projection > ruffle. Rapidly redistributed to ruffles and

lamellipodia structures in response to epidermal growth factor (EGF) treatment.

Images



Western blot - Anti-Phospholipase C gamma 1/PLCgamma-1 (phospho Y783) antibody [EPR2611Y] -BSA and Azide free (ab248714) **All lanes :** Anti-Phospholipase C gamma 1/PLC-gamma-1 (phospho Y783) antibody [EPR2611Y] (ab134186) at 1/10000 dilution

Lane 1: Jurkat cell lysate untreated

Lane 2: Jurkat cell lysate treated with Pervanadate

Lysates/proteins at 10 µg per lane.

Secondary

All lanes: HRP labelled goat anti-rabbit at 1/2000 dilution

Predicted band size: 148 kDa Observed band size: 150 kDa

This data was developed using <u>ab134186</u>, the same antibody clone in a different buffer formulation.

Why choose a recombinant antibody?







scalable supply Recombinant technology





Animal-free

production

Confirmed specificity

Azide free (ab248714)

Anti-Phospholipase C gamma 1/PLC-gamma-1 (phospho Y783) antibody [EPR2611Y] - BSA and

Please note: All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit https://www.abcam.com/abpromise or contact our technical team.

Terms and conditions

· Guarantee only valid for products bought direct from Abcam or one of our authorized distributors