

Anti-PI 3 Kinase p85 alpha antibody [EPR18702] - BSA and Azide free ab223792

KO VALIDATED Recombinant RabMAB

8 Images

Overview

Product name	Anti-PI 3 Kinase p85 alpha antibody [EPR18702] - BSA and Azide free
Description	Rabbit monoclonal [EPR18702] to PI 3 Kinase p85 alpha - BSA and Azide free
Host species	Rabbit
Tested applications	Suitable for: Flow Cyt (Intra), ICC/IF, IP, WB
Species reactivity	Reacts with: Mouse, Rat, Human, African green monkey
Immunogen	Synthetic peptide. This information is proprietary to Abcam and/or its suppliers.
Positive control	WB: Human PI3K p85 alpha full length recombinant protein; Human fetal liver, fetal heart and fetal kidney lysates; HeLa, HepG2, MCF7, Raji, Jurkat, C6, RAW 264.7, PC-12 and NIH/3T3 whole cell lysates; Mouse brain, heart, kidney and spleen lysates; Rat brain, heart, kidney and spleen lysates. ICC/IF: HepG2 and NIH/3T3 cells. Flow Cyt (intra): NIH/3T3 cells; IP: MCF7 whole cell lysate.
General notes	<p>ab223792 is the carrier-free version of ab191606.</p> <p>Our carrier-free antibodies are typically supplied in a PBS-only formulation, purified and free of BSA, sodium azide and glycerol. The carrier-free buffer and high concentration allow for increased conjugation efficiency.</p> <p>This conjugation-ready format is designed for use with fluorochromes, metal isotopes, oligonucleotides, and enzymes, which makes them ideal for antibody labelling, functional and cell-based assays, flow-based assays (e.g. mass cytometry) and Multiplex Imaging applications.</p> <p>Use our conjugation kits for antibody conjugates that are ready-to-use in as little as 20 minutes with <1 minute hands-on-time and 100% antibody recovery: available for fluorescent dyes, HRP, biotin and gold.</p> <p>This product is compatible with the Maxpar[®] Antibody Labeling Kit from Fluidigm, without the need for antibody preparation. Maxpar[®] is a trademark of Fluidigm Canada Inc.</p> <p>This product is a recombinant monoclonal antibody, which offers several advantages including:</p> <ul style="list-style-type: none">- High batch-to-batch consistency and reproducibility- Improved sensitivity and specificity- Long-term security of supply- Animal-free production <p>For more information see here.</p>

Our RabMAb[®] technology is a patented hybridoma-based technology for making rabbit monoclonal antibodies. For details on our patents, please refer to [RabMAb[®] patents](#).

Properties

Form	Liquid
Storage instructions	Shipped at 4°C. Store at +4°C. Do Not Freeze.
Storage buffer	pH: 7.2 Constituent: PBS
Carrier free	Yes
Purity	Protein A purified
Clonality	Monoclonal
Clone number	EPR18702
Isotype	IgG

Applications

The Abpromise guarantee Our [Abpromise guarantee](#) covers the use of ab223792 in the following tested applications. The application notes include recommended starting dilutions; optimal dilutions/concentrations should be determined by the end user.

Application	Abreviews	Notes
Flow Cyt (Intra)		Use at an assay dependent concentration. ab199376 - Rabbit monoclonal IgG, is suitable for use as an isotype control with this antibody.
ICC/IF		Use at an assay dependent concentration.
IP		Use at an assay dependent concentration.
WB		Use at an assay dependent concentration. Detects a band of approximately 85,46 kDa (predicted molecular weight: 84 kDa).

Target

Function	Binds to activated (phosphorylated) protein-Tyr kinases, through its SH2 domain, and acts as an adapter, mediating the association of the p110 catalytic unit to the plasma membrane. Necessary for the insulin-stimulated increase in glucose uptake and glycogen synthesis in insulin-sensitive tissues.
Tissue specificity	Isoform 2 is expressed in skeletal muscle and brain, and at lower levels in kidney and cardiac muscle. Isoform 2 and isoform 4 are present in skeletal muscle (at protein level).
Sequence similarities	Belongs to the PI3K p85 subunit family. Contains 1 Rho-GAP domain. Contains 2 SH2 domains. Contains 1 SH3 domain.


Domain

The SH3 domain mediates the binding to CBLB, and to HIV-1 Nef.

Post-translational modifications

Polyubiquitinated in T-cells by CBLB; which does not promote proteasomal degradation but impairs association with CD28 and CD3Z upon T-cell activation.
Phosphorylated. Dephosphorylated by PTPRJ.

Images



Western blot - Anti-PI 3 Kinase p85 alpha antibody [EPR18702] - BSA and Azide free (ab223792)

This WB data was generated using the same anti-PI 3 Kinase p85 alpha antibody clone [EPR18702] in a different buffer formulation containing PBS, BSA, glycerol, and sodium azide (cat# **ab191606**).


Lane 1: Wild type HAP1 whole cell lysate (20 µg)

Lane 2: PIK3R1 knockout HAP1 whole cell lysate (20 µg)

Lane 3: Jurkat whole cell lysate (20 µg)

Lanes 1 - 4: Merged signal (red and green). Green - **ab191606** observed at 90 kDa. Red - loading control, **ab9484**, observed at 37 kDa.


ab191606 was shown to specifically react with PIK3R1 when PIK3R1 knockout samples were used. Wild-type and PIK3R1 knockout samples were subjected to SDS-PAGE. Ab191606 and **ab9484** (Mouse anti GAPDH loading control) were incubated overnight at 4°C at 1000 dilution and 1/20000 dilution respectively. Blots were developed with Goat anti-Rabbit IgG H&L (IRDye® 800CW) preabsorbed **ab216773** and Goat anti-Mouse IgG H&L (IRDye® 680RD) preabsorbed **ab216776** secondary antibodies at 1/20000 dilution for 1 hour at room temperature before imaging.



Flow Cytometry (Intracellular) - Anti-PI 3 Kinase p85 alpha antibody [EPR18702] - BSA and Azide free (ab223792)

Overlay histogram showing HepG2 cells fixed in 4% PFA and stained with **ab191606** at a dilution of 1/80 (red line). The secondary antibody used was Alexa Fluor® 488 goat anti-rabbit at a dilution of 1/500. Rabbit monoclonal IgG (**ab172730**) was used as an isotype control (black line) and cells incubated in the absence of both primary and secondary antibody were used as a negative control (blue line).

This data was developed using the same antibody clone in a different buffer formulation containing PBS, BSA, glycerol, and sodium azide (**ab191606**).



Immunocytochemistry/ Immunofluorescence - Anti-PI 3 Kinase p85 alpha antibody [EPR18702] - BSA and Azide free (ab223792)


Immunofluorescent analysis of 4% paraformaldehyde-fixed, 0.1% Triton X-100 permeabilized HepG2 (Human liver hepatocellular carcinoma cell line) cells labeling PI3K p85 with **ab191606** at 1/250 dilution, followed by Goat Anti-Rabbit IgG (Alexa Fluor® 488) (**ab150077**) secondary antibody at 1/1000 dilution (green). Confocal image showing cytoplasmic staining on HepG2 cell line. The nuclear counter stain is DAPI (blue). Tubulin is detected with Anti-alpha Tubulin antibody - Loading Control (**ab7291**) at 1/1000 dilution and Goat Anti-Mouse IgG (AlexaFluor®594) preadsorbed (**ab150120**) at 1/1000 dilution (red).

The negative controls are as follows:

-ve control 1: **ab191606** at 1/500 dilution followed by **ab150120** at 1/1000 dilution.

-ve control 2: **ab7291** at 1/1000 dilution followed by **ab150077** at 1/1000 dilution.

This data was developed using the same antibody clone in a different buffer formulation containing PBS, BSA, glycerol, and sodium azide (**ab191606**).



Western blot - Anti-PI 3 Kinase p85 alpha antibody [EPR18702] - BSA and Azide free (ab223792)

All lanes : Anti-PI 3 Kinase p85 alpha antibody [EPR18702] (**ab191606**) at 1/1000 dilution

Lane 1 : Wild-type HeLa cell lysate

Lane 2 : PIK3R1 knockout HeLa cell lysate

Lysates/proteins at 20 µg per lane.

Performed under reducing conditions.

Predicted band size: 84 kDa

Observed band size: 90 kDa


This data was developed using the same antibody clone in a different buffer formulation (**ab191606**).

Lanes 1- 2: Merged signal (red and green). Green - **ab191606** observed at 90 kDa. Red - Anti-GAPDH antibody [6C5] - Loading Control (**ab8245**) observed at 37 kDa.

ab191606 was shown to react with PI 3 Kinase p85 alpha in wild-type HeLa cells in western blot. Loss of signal was observed when knockout cell line **ab265116** (knockout cell lysate **ab257029**) was

used. Wild-type HeLa and PIK3R1 knockout HeLa cell lysates were subjected to SDS-PAGE. Membrane was blocked for 1 hour at room temperature in 0.1% TBST with 3% non-fat dried milk.

ab191606 and Anti-GAPDH antibody [6C5] - Loading Control (**ab8245**) overnight at 4°C at a 1 in 1000 dilution and a 1 in 20000 dilution respectively. Blots were developed with Goat anti-Rabbit IgG H&L (IRDye®800CW) preadsorbed (**ab216773**) and Goat anti-Mouse IgG H&L (IRDye®680RD) preadsorbed (**ab216776**) secondary antibodies at 1 in 20000 dilution for 1 hour at room temperature before imaging.



Immunocytochemistry/ Immunofluorescence - Anti-PI 3 Kinase p85 alpha antibody [EPR18702] - BSA and Azide free (ab223792)


Immunofluorescent analysis of 4% paraformaldehyde-fixed, 100% Methanol permeabilized NIH/3T3 (Mouse embryonic fibroblast cell line) cells labeling PI3K p85 with **ab191606** at 1/250 dilution, followed by Goat anti-rabbit IgG (Alexa Fluor® 488) (**ab150077**) secondary antibody at 1/1000 dilution (green). Confocal image showing cytoplasmic staining on NIH/3T3 cell line. The nuclear counter stain is DAPI (blue). Tubulin is detected with Anti-alpha Tubulin antibody - Loading Control (**ab7291**) at 1/1000 dilution and Goat Anti-Mouse IgG (AlexaFluor®594) preadsorbed (**ab150120**) at 1/1000 dilution (red).

The negative controls are as follows:

-ve control 1: **ab191606** at 1/500 dilution followed by **ab150120** at 1/1000 dilution.

-ve control 2: **ab7291** at 1/1000 dilution followed by **ab150077** at 1/1000 dilution.


This data was developed using the same antibody clone in a different buffer formulation containing PBS, BSA, glycerol, and sodium azide (**ab191606**).



Flow Cytometry (Intracellular) - Anti-PI 3 Kinase p85 alpha antibody [EPR18702] - BSA and Azide free (ab223792)

Intracellular flow cytometric analysis of 4% paraformaldehyde-fixed NIH/3T3 (Mouse embryonic fibroblast cell line) cells labeling PI3K p85 with **ab191606** at 1/150 dilution (red) compared with a Rabbit IgG, monoclonal - Isotype control (**ab172730**) (black) and an unlabelled control (cells without incubation with primary antibody and secondary antibody) (blue). Goat anti Rabbit IgG (Alexa Fluor® 488) at 1/500 dilution was used as the secondary antibody.

This data was developed using the same antibody clone in a different buffer formulation containing PBS, BSA, glycerol, and sodium azide (**ab191606**).



Immunoprecipitation - Anti-PI 3 Kinase p85 alpha antibody [EPR18702] - BSA and Azide free (ab223792)

PI3K p85 was immunoprecipitated from 1mg of MCF7 (Human breast adenocarcinoma cell line) whole cell lysate with **ab191606** at 1/50 dilution. Western blot was performed from the immunoprecipitate using **ab191606** at 1/1000 dilution. VeriBlot for IP Detection Reagent (HRP) (**ab131366**), was used for detection at 1/10000 dilution.

Lane 1: MCF7 whole cell lysate, 10µg (Input).

Lane 2: **ab191606** IP in MCF7 whole cell lysate.

Lane 3: Rabbit IgG, monoclonal - Isotype Control (**ab172730**) instead of **ab191606** in MCF7 whole cell lysate.

Blocking and dilution buffer and concentration: 5% NFDm/TBST.

Exposure time: 5 seconds.

This data was developed using the same antibody clone in a different buffer formulation containing PBS, BSA, glycerol, and sodium azide (**ab191606**).

Why choose a recombinant antibody?



Research with confidence
Consistent and reproducible results



Long-term and scalable supply
Recombinant technology



Success from the first experiment
Confirmed specificity



Ethical standards compliant
Animal-free production

Anti-PI 3 Kinase p85 alpha antibody [EPR18702] -
BSA and Azide free (ab223792)

Please note: All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <https://www.abcam.com/abpromise> or contact our technical team.

Terms and conditions

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors