

Product datasheet

Anti-PKN1 antibody [EPR18808] - BSA and Azide free ab251202

KO VALIDATED Recombinant RabMAb

6 Images

Overview

Product name	Anti-PKN1 antibody [EPR18808] - BSA and Azide free
Description	Rabbit monoclonal [EPR18808] to PKN1 - BSA and Azide free
Host species	Rabbit
Tested applications	Suitable for: WB, IP
Species reactivity	Reacts with: Mouse, Rat, Human
Immunogen	Recombinant fragment. This information is proprietary to Abcam and/or its suppliers.
General notes	<p>ab251202 is the carrier-free version of ab195264.</p> <p>Our carrier-free antibodies are typically supplied in a PBS-only formulation, purified and free of BSA, sodium azide and glycerol. The carrier-free buffer and high concentration allow for increased conjugation efficiency.</p> <p>This conjugation-ready format is designed for use with fluorochromes, metal isotopes, oligonucleotides, and enzymes, which makes them ideal for antibody labelling, functional and cell-based assays, flow-based assays (e.g. mass cytometry) and Multiplex Imaging applications.</p> <p>Use our conjugation kits for antibody conjugates that are ready-to-use in as little as 20 minutes with <1 minute hands-on-time and 100% antibody recovery: available for fluorescent dyes, HRP, biotin and gold.</p> <p>This product is compatible with the Maxpar[®] Antibody Labeling Kit from Fluidigm, without the need for antibody preparation. Maxpar[®] is a trademark of Fluidigm Canada Inc.</p> <p>This product is a recombinant monoclonal antibody, which offers several advantages including:</p> <ul style="list-style-type: none"> - High batch-to-batch consistency and reproducibility - Improved sensitivity and specificity - Long-term security of supply - Animal-free production <p>For more information see here.</p> <p>Our RabMAb[®] technology is a patented hybridoma-based technology for making rabbit monoclonal antibodies. For details on our patents, please refer to RabMAb[®] patents.</p>

Properties

Form	Liquid
Storage instructions	Shipped at 4°C. Store at +4°C. Do Not Freeze.
Storage buffer	pH: 7.2 Constituent: PBS
Carrier free	Yes
Purity	Protein A purified
Clonality	Monoclonal
Clone number	EPR18808
Isotype	IgG

Applications

The Abpromise guarantee Our **Abpromise guarantee** covers the use of ab251202 in the following tested applications.

The application notes include recommended starting dilutions; optimal dilutions/concentrations should be determined by the end user.

Application	Abreviews	Notes
WB		Use at an assay dependent concentration. Detects a band of approximately 120 kDa (predicted molecular weight: 104 kDa).
IP		Use at an assay dependent concentration.

Target

Function	PKC-related serine/threonine-protein kinase involved in various processes such as regulation of the intermediate filaments of the actin cytoskeleton, cell migration, tumor cell invasion and transcription regulation. Regulates the cytoskeletal network by phosphorylating proteins such as VIM and neurofilament proteins NEFH, NEFL and NEFM, leading to inhibit their polymerization. Phosphorylates 'Ser-575', 'Ser-637' and 'Ser-669' of MAPT/Tau, lowering its ability to bind to microtubules, resulting in disruption of tubulin assembly. Acts as a key coactivator of androgen receptor (ANDR)-dependent transcription, by being recruited to ANDR target genes and specifically mediating phosphorylation of 'Thr-11' of histone H3 (H3T11ph), a specific tag for epigenetic transcriptional activation that promotes demethylation of histone H3 'Lys-9' (H3K9me) by KDM4C/JMJD2C. Phosphorylates HDAC5, HDAC7 and HDAC9, leading to impair their import in the nucleus. Phosphorylates 'Thr-38' of PPP1R14A, 'Ser-159', 'Ser-163' and 'Ser-170' of MARCKS, and GFAP. Able to phosphorylate RPS6 in vitro.
Tissue specificity	Found ubiquitously. Expressed in heart, brain, placenta, lung, skeletal muscle, kidney and pancreas. Expressed in numerous tumor cell lines, especially in breast tumor cells.
Sequence similarities	Belongs to the protein kinase superfamily. AGC Ser/Thr protein kinase family. PKC subfamily. Contains 1 AGC-kinase C-terminal domain. Contains 1 C2 domain. Contains 1 protein kinase domain. Contains 3 REM (Hr1) repeats.
Domain	The C1 domain does not bind the diacylglycerol (DAG).
Post-translational	Autophosphorylated; preferably on serine. Phosphorylated during mitosis.


modifications

Cellular localization

Activated by limited proteolysis with trypsin.

Cytoplasm. Nucleus. Endosome. Cell membrane. Cleavage furrow. Midbody. Associates with chromatin in a ligand-dependent manner. Localization to endosomes is mediated via its interaction with RHOB. Association to the cell membrane is dependent on Ser-374 phosphorylation. Accumulates during telophase at the cleavage furrow and finally concentrates around the midbody in cytokinesis.

Images



This data was developed using [ab195264](#), the same antibody clone in a different buffer formulation.

Lane 1: Wild-type HAP1 cell lysate (40 µg)


Lane 2: PKN1 knockout HAP1 cell lysate (40 µg)

Lane 3: MCF7 cell lysate (20 µg)

Lane 4: K562 cell lysate (20 µg)

Lanes 1 - 4: Merged signal (red and green). Green - [ab195264](#) observed at 125 kDa. Red - loading control, [ab8245](#), observed at 37 kDa.

[ab195264](#) was shown to specifically react with PKN1 when PKN1 knockout samples were used. Wild-type and PKN1 knockout samples were subjected to SDS-PAGE. [ab195264](#) and [ab8245](#) (loading control to GAPDH) were diluted at 1/1000 and 1/10000 dilution respectively and incubated overnight at 4°C. Blots were developed with Goat anti-Rabbit IgG H&L (IRDye® 800CW) preadsorbed [ab216773](#) and Goat anti-Mouse IgG H&L (IRDye® 680RD) preadsorbed [ab216776](#) secondary antibodies at 1/10000 dilution for 1 hour at room temperature before imaging.



All lanes : Anti-PKN1 antibody [EPR18808] ([ab195264](#)) at 1/1000 dilution

Lane 1 : Jurkat (Human T cell leukemia cell line from peripheral blood) whole cell lysate

Lane 2 : LNCaP (Human prostate cancer cell line) whole cell lysate

Lane 3 : MDA-MB-231 (Human breast adenocarcinoma cell line) whole cell lysate

Lane 4 : K562 (Human chronic myelogenous leukemia cell line from bone marrow) whole cell lysate

Lane 5 : HEK-293 (Human epithelial cell line from embryonic kidney) whole cell lysate

Lane 6 : MCF-7 (Human breast adenocarcinoma cell line) whole cell lysate

Lysates/proteins at 20 µg per lane.

Secondary

All lanes : Goat Anti-Rabbit IgG H&L (HRP) ([ab97051](#)) at 1/100000 dilution


Predicted band size: 104 kDa

Observed band size: 120 kDa


Exposure time: 3 minutes

This data was developed using [ab195264](#), the same antibody clone in a different buffer formulation.

Blocking and dilution buffer: 5% NFDM/TBST.



This data was developed using [ab195264](#), the same antibody clone in a different buffer formulation. PKN1 was immunoprecipitated from 1mg of Jurkat (Human T cell leukemia cell line from peripheral blood) whole cell lysate with [ab195264](#) at 1/40 dilution. Western blot was performed from the immunoprecipitate using [ab195264](#) at 1/1000 dilution. VeriBlot for IP Detection Reagent (HRP) ([ab131366](#)), was used for detection at 1/10000 dilution. Lane 1: Jurkat whole cell lysate, 10µg (Input). Lane 2: [ab195264](#) IP in Jurkat whole cell lysate. Lane 3: Rabbit IgG, monoclonal-Isotype Control ([ab172730](#)) instead of [ab195264](#) in Jurkat whole cell lysate. Blocking and dilution buffer and concentration: 5% NFDM/TBST. Exposure time: 3 minutes.



Western blot - Anti-PKN1 antibody [EPR18808] - BSA and Azide free (ab251202)

All lanes : Anti-PKN1 antibody [EPR18808] ([ab195264](#)) at 1/1000 dilution

Lane 1 : C6 (Rat glial tumor cell line) whole cell lysate

Lane 2 : RAW 264.7 (Mouse macrophage cell line transformed with Abelson murine leukemia virus) whole cell lysate

Lane 3 : PC-12 (Rat adrenal gland pheochromocytoma cell line) whole cell lysate

Lane 4 : NIH/3T3 (Mouse embryonic fibroblast cell line) whole cell lysate

Lysates/proteins at 10 µg per lane.

Secondary

All lanes : Goat Anti-Rabbit IgG H&L (HRP) ([ab97051](#)) at 1/100000 dilution


Predicted band size: 104 kDa

Observed band size: 120 kDa

Exposure time: 3 minutes

This data was developed using [ab195264](#), the same antibody clone in a different buffer formulation.

Blocking and dilution buffer: 5% NFDM/TBST.



Western blot - Anti-PKN1 antibody [EPR18808] - BSA and Azide free (ab251202)

All lanes : Anti-PKN1 antibody [EPR18808] ([ab195264](#)) at 1/1000 dilution

Lane 1 : Rat brain lysate

Lane 2 : Rat spleen lysate

Lysates/proteins at 10 µg per lane.

Secondary

All lanes : Goat Anti-Rabbit IgG H&L (HRP) ([ab97051](#)) at 1/100000 dilution


Predicted band size: 104 kDa

Observed band size: 120 kDa

Exposure time: 3 minutes

This data was developed using **ab195264**, the same antibody clone in a different buffer formulation.

Why choose a recombinant antibody?



- Research with confidence**
Consistent and reproducible results
- Long-term and scalable supply**
Recombinant technology
- Success from the first experiment**
Confirmed specificity
- Ethical standards compliant**
Animal-free production

Anti-PKN1 antibody [EPR18808] - BSA and Azide free (ab251202)

Please note: All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <https://www.abcam.com/abpromise> or contact our technical team.

Terms and conditions

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors