

## Product datasheet

# Anti-PUS1 antibody [EPR20181] ab203010

**KO VALIDATED** Recombinant RabMAB

[1 References](#) [10 Images](#)

### Overview

<b>Product name</b>	Anti-PUS1 antibody [EPR20181]
<b>Description</b>	Rabbit monoclonal [EPR20181] to PUS1
<b>Host species</b>	Rabbit
<b>Tested applications</b>	<b>Suitable for:</b> WB, ICC/IF, IP, Flow Cyt (Intra)
<b>Species reactivity</b>	<b>Reacts with:</b> Mouse, Rat, Human
<b>Immunogen</b>	Recombinant fragment. This information is proprietary to Abcam and/or its suppliers.
<b>Positive control</b>	WB: Human skeletal muscle lysate; HeLa mitochondrial and cytoplasm fraction lysates; A431, Daudi, HEK-293, C6, RAW 264.7, PC-12 and NIH/3T3 whole cell lysates; Mouse heart and spleen lysates; Rat spleen lysate. ICC/IF: HeLa cells. Flow Cyt (intra): HeLa cells. IP: HEK-293 whole cell lysate.
<b>General notes</b>	<p>This product is a recombinant monoclonal antibody, which offers several advantages including:</p> <ul style="list-style-type: none"><li>- High batch-to-batch consistency and reproducibility</li><li>- Improved sensitivity and specificity</li><li>- Long-term security of supply</li><li>- Animal-free production</li></ul> <p>For more information <a href="#">see here</a>.</p> <p>Our RabMAB<sup>®</sup> technology is a patented hybridoma-based technology for making rabbit monoclonal antibodies. For details on our patents, please refer to <a href="#">RabMAB<sup>®</sup> patents</a>.</p>

### Properties

<b>Form</b>	Liquid
<b>Storage instructions</b>	Shipped at 4°C. Store at +4°C short term (1-2 weeks). Upon delivery aliquot. Store at -20°C long term. Avoid freeze / thaw cycle.
<b>Storage buffer</b>	pH: 7.2 Preservative: 0.01% Sodium azide Constituents: 59% PBS, 40% Glycerol (glycerin, glycerine), 0.05% BSA
<b>Purity</b>	Protein A purified
<b>Clonality</b>	Monoclonal
<b>Clone number</b>	EPR20181

Isotype

IgG

## Applications


**The Abpromise guarantee** Our **Abpromise guarantee** covers the use of ab203010 in the following tested applications. The application notes include recommended starting dilutions; optimal dilutions/concentrations should be determined by the end user.

Application	Abreviews	Notes
WB		1/1000. Detects a band of approximately 47, 44 kDa (predicted molecular weight: 47 kDa).
ICC/IF		1/500. This antibody was not successful when we used it on RAW 264.7 (Mouse macrophage cell line transformed with Abelson murine leukemia virus) cells in ICC application. This antibody was not tested on rat cells in ICC.
IP		1/30.
Flow Cyt (Intra)		1/400.

## Target

<b>Function</b>	Converts specific uridines to PSI in a number of tRNA substrates. Acts on positions 27/28 in the anticodon stem and also positions 34 and 36 in the anticodon of an intron containing tRNA. Involved in regulation of nuclear receptor activity possibly through pseudouridylation of SRA1 RNA.
<b>Tissue specificity</b>	Widely expressed. High levels of expression found in brain and skeletal muscle.
<b>Involvement in disease</b>	Defects in PUS1 are a cause of myopathy with lactic acidosis and sideroblastic anemia type 1 (MLASA1) [MIM:600462]; also known as mitochondrial myopathy and sideroblastic anemia. MLASA is a rare autosomal recessive oxidative phosphorylation disorder specific to skeletal muscle and bone marrow.
<b>Sequence similarities</b>	Belongs to the tRNA pseudouridine synthase TruA family.
<b>Cellular localization</b>	Mitochondrion and Nucleus.

## Images



Western blot - Anti-PUS1 antibody [EPR20181]  
(ab203010)

**All lanes** : Anti-PUS1 antibody [EPR20181] (ab203010) at 1/1000 dilution

**Lane 1** : Wild-type HEK-293T cell lysate

**Lane 2** : PUS1 knockout HEK-293T cell lysate

**Lane 3** : HeLa cell lysate

**Lane 4** : Daudi cell lysate

Lysates/proteins at 20 µg per lane.


Performed under reducing conditions.

**Predicted band size:** 47 kDa

**Observed band size:** 45 kDa

**Lanes 1- 4:** Merged signal (red and green). Green - ab203010 observed at 45 kDa. Red - Anti-alpha Tubulin antibody [DM1A] - Loading Control ([ab7291](#)) observed at 50 kDa.

ab203010 was shown to react with PUS1 in wild-type HEK-293T cells in western blot. Loss of signal was observed when knockout cell line [ab266091](#) (knockout cell lysate [ab258158](#)) was used. Wild-type HEK-293T and PUS1 knockout HEK-293T cell lysates were subjected to SDS-PAGE. Membrane was blocked for 1 hour at room temperature in 0.1% TBST with 3% non-fat dried milk. ab203010 and Anti-alpha Tubulin antibody [DM1A] - Loading Control ([ab7291](#)) overnight at 4°C at a 1 in 1000 dilution and a 1 in 20000 dilution respectively. Blots were developed with Goat anti-Rabbit IgG H&L (IRDye®800CW) preadsorbed ([ab216773](#)) and Goat anti-Mouse IgG H&L (IRDye®680RD) preadsorbed ([ab216776](#)) secondary antibodies at 1 in 20000 dilution for 1 hour at room temperature before imaging.




Immunocytochemistry/ Immunofluorescence - Anti-PUS1 antibody [EPR20181] (ab203010)

Immunofluorescent analysis of 4% paraformaldehyde-fixed, 0.1% Triton X-100 permeabilized HeLa (Human epithelial cell line from cervix adenocarcinoma) cells labeling PUS1 with ab203010 at 1/500 dilution, followed by Goat anti-rabbit IgG (Alexa Fluor® 488) (**ab150077**) secondary antibody at 1/1000 dilution (green). Confocal image showing cytoplasmic staining on HeLa cell line.

The nuclear counterstain is DAPI (blue). Tubulin is detected with **ab195889** (Anti-alpha Tubulin antibody [DM1A] - Microtubule Marker (Alexa Fluor® 594)) at 1/200 dilution (red).

Secondary antibody only control: Used PBS instead of primary antibody, secondary antibody is Goat anti-rabbit IgG (Alexa Fluor® 488) (**ab150077**) at 1/1000 dilution.

This antibody was not successful when we used it on RAW 264.7 (Mouse macrophage cell line transformed with Abelson murine leukemia virus) cells in ICC application. This antibody was not tested on rat cells in ICC.



Immunocytochemistry/ Immunofluorescence - Anti-PUS1 antibody [EPR20181] (ab203010)


Immunofluorescent analysis of 4% paraformaldehyde-fixed, 0.1% Triton X-100 permeabilized HeLa (Human epithelial cell line from cervix adenocarcinoma) cells labeling PUS1 with ab203010 at 1/500 dilution, followed by Goat anti-rabbit IgG (Alexa Fluor® 488) (**ab150077**) secondary antibody at 1/1000 dilution (green). Confocal image showing mitochondrial staining on HeLa cell line.

The nuclear counterstain is DAPI (blue). COXIV is detected with **ab33985** (Anti-COX IV(mouse mAb)) at 1/1000 dilution followed by Goat anti-mouse IgG (Alexa Fluor® 594) (**ab150120**) secondary antibody at 1/1000 dilution (red).

The negative controls are as follows:-

-ve control 1: ab203010 at 1/500 dilution followed by **ab150120** (Alexa Fluor® 594 Goat anti-Mouse secondary) at 1/1000 dilution.

-ve control 2: **ab33985** (anti-COX IV(mouse mAb)) at 1/1000 dilution followed by **ab150077** (Alexa Fluor® 488 Goat Anti-Rabbit IgG H&L) at 1/1000 dilution.



Western blot - Anti-PUS1 antibody [EPR20181]  
(ab203010)

Anti-PUS1 antibody [EPR20181] (ab203010) at 1/1000 dilution +  
Human skeletal muscle lysate at 10 µg

**Secondary**


Goat Anti-Rabbit IgG Peroxidase Conjugate, specific to the non-reduced form of IgG at 1/1000 dilution

**Predicted band size:** 47 kDa

**Observed band size:** 47 kDa

**Exposure time:** 3 minutes

Blocking/Dilution buffer: 5% NFDm/TBST.



Western blot - Anti-PUS1 antibody [EPR20181]  
(ab203010)

**All lanes :** Anti-PUS1 antibody [EPR20181] (ab203010) at 1/5000 dilution

**Lane 1 :** HeLa (Human epithelial cell line from cervix adenocarcinoma) mitochondria lysate

**Lane 2 :** HeLa cytoplasm fraction lysate

**Lane 3 :** A431 (Human epidermoid carcinoma cell line) whole cell lysate

Lysates/proteins at 20 µg per lane.

**Secondary**

**All lanes :** Goat Anti-Rabbit IgG H&L (HRP) (**ab97051**) at 1/100000 dilution

**Predicted band size:** 47 kDa

**Observed band size:** 47 kDa

**Exposure time:** 8 seconds

Blocking/Dilution buffer: 5% NFDm/TBST.

The expression profile observed is consistent with what has been described in UniProt.

Anti-PUS1 antibody [EPR20181] (ab203010) at 1/1000 dilution + HEK-293 (Human epithelial cell line from embryonic kidney) whole cell lysate at 10 µg

### Secondary

Goat Anti-Rabbit IgG H&L (HRP) ([ab97051](#)) at 1/100000 dilution

**Predicted band size:** 47 kDa

**Observed band size:** 44,47 kDa

**Exposure time:** 3 seconds

Blocking/Dilution buffer: 5% NFDm/TBST.

Based on sequence alignment, the antibody can recognize 2 isoforms, the predicted MW are 47kDa and 44kDa, respectively [PMID: 17056637].

**All lanes :** Anti-PUS1 antibody [EPR20181] (ab203010) at 1/2000 dilution

**Lane 1 :** Mouse heart tissue lysate

**Lane 2 :** Mouse spleen tissue lysate

**Lane 3 :** Rat spleen tissue lysate

**Lane 4 :** C6 (Rat glial tumor cell line) whole cell lysate

**Lane 5 :** RAW 264.7 (Mouse macrophage cell line transformed with Abelson murine leukemia virus) whole cell lysate

**Lane 6 :** PC-12 (Rat adrenal gland pheochromocytoma cell line) whole cell lysate


**Lane 7 :** NIH/3T3 (Mouse embryonic fibroblast cell line) whole cell lysate

Lysates/proteins at 10 µg per lane.


### Secondary

**All lanes :** Goat Anti-Rabbit IgG H&L (HRP) ([ab97051](#)) at 1/100000 dilution

**Predicted band size:** 47 kDa



Western blot - Anti-PUS1 antibody [EPR20181] (ab203010)




Western blot - Anti-PUS1 antibody [EPR20181] (ab203010)

**Observed band size:** 47 kDa


Blocking/Dilution buffer: 5% NFD/MTBST.

Exposure time: Lane 1 and 2: 10 seconds; Lane 3: 3minutes; Lane 4,5,6 and 7: 10 seconds.



Flow Cytometry (Intracellular) - Anti-PUS1 antibody [EPR20181] (ab203010)

Intracellular flow cytometric analysis of 4% paraformaldehyde-fixed HeLa (Human epithelial cell line from cervix adenocarcinoma) cells labeling PUS1 with ab203010 at 1/400 dilution (red) compared with a rabbit monoclonal IgG isotype control (**ab172730**; black) and an unlabeled control (cells without incubation with primary antibody and secondary antibody; blue). Goat anti rabbit IgG (Alexa Fluor® 488) at 1/2000 dilution was used as the secondary antibody.



Immunoprecipitation - Anti-PUS1 antibody [EPR20181] (ab203010)

PUS1 was immunoprecipitated from 0.35 mg of HEK-293 (Human epithelial cell line from embryonic kidney) whole cell lysate with ab203010 at 1/30 dilution. Western blot was performed from the immunoprecipitate using ab203010 at 1/1000 dilution. VeriBlot for IP Detection Reaction (HRP) (**ab131366**), was used for detection at 1/10000 dilution.

Lane 1: HEK-293 whole cell lysate, 10µg (Input).


Lane 2: ab203010 IP in HEK-293 whole cell lysate.

Lane 3: Rabbit monoclonal IgG (**ab172730**) instead of ab203010 in HEK-293 whole cell lysate.

Blocking and dilution buffer and concentration: 5% NFD/MTBST.

Exposure time: 5 seconds.

Why choose a recombinant antibody?



**Research with confidence**  
Consistent and reproducible results

**Long-term and scalable supply**  
Recombinant technology

**Success from the first experiment**  
Confirmed specificity

**Ethical standards compliant**  
Animal-free production

Anti-PUS1 antibody [EPR20181] (ab203010)

**Please note:** All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

### Our Abpromise to you: Quality guaranteed and expert technical support

---

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <https://www.abcam.com/abpromise> or contact our technical team.

### Terms and conditions

---

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors