# abcam

## Product datasheet

## Anti-Quiescin Q6 antibody [EPR21866] - BSA and Azide free ab237600



## 5 Images

#### Overview

**Product name** Anti-Quiescin Q6 antibody [EPR21866] - BSA and Azide free

**Description** Rabbit monoclonal [EPR21866] to Quiescin Q6 - BSA and Azide free

**Host species** Rabbit

**Tested applications** Suitable for: Flow Cyt (Intra), IHC-P, WB, IP

Species reactivity Reacts with: Human

**Immunogen** Synthetic peptide. This information is proprietary to Abcam and/or its suppliers.

Positive control IHC-P: Human pancreatic cancer tissue.

**General notes** ab237600 is the carrier-free version of ab235444.

> Our carrier-free antibodies are typically supplied in a PBS-only formulation, purified and free of BSA, sodium azide and glycerol. The carrier-free buffer and high concentration allow for increased conjugation efficiency.

This conjugation-ready format is designed for use with fluorochromes, metal isotopes, oligonucleotides, and enzymes, which makes them ideal for antibody labelling, functional and cellbased assays, flow-based assays (e.g. mass cytometry) and Multiplex Imaging applications.

Use our conjugation kits for antibody conjugates that are ready-to-use in as little as 20 minutes with <1 minute hands-on-time and 100% antibody recovery: available for fluorescent dyes, HRP, biotin and gold.

This product is compatible with the Maxpar® Antibody Labeling Kit from Fluidigm, without the need for antibody preparation. Maxpar<sup>®</sup> is a trademark of Fluidigm Canada Inc.

This product is a recombinant monoclonal antibody, which offers several advantages including:

- High batch-to-batch consistency and reproducibility
- Improved sensitivity and specificity
- Long-term security of supply
- Animal-free production

For more information see here.

Our RabMAb® technology is a patented hybridoma-based technology for making rabbit monoclonal antibodies. For details on our patents, please refer to **RabMAb**® **patents**.

#### **Properties**

Form Liquid

**Storage instructions** Shipped at 4°C. Store at +4°C. Do Not Freeze.

Storage buffer pH: 7.2

Constituent: PBS

Carrier free Yes

Purity Protein A purified

Clonality Monoclonal
Clone number EPR21866

**Isotype** IgG

#### **Applications**

The Abpromise guarantee Our Abpromise guarantee covers the use of ab237600 in the following tested applications.

The application notes include recommended starting dilutions; optimal dilutions/concentrations should be determined by the end user.

Application	Abreviews	Notes
Flow Cyt (Intra)		Use at an assay dependent concentration.
IHC-P		Use at an assay dependent concentration. Perform heat mediated antigen retrieval with Tris/EDTA buffer pH 9.0 before commencing with IHC staining protocol.
WB		Use at an assay dependent concentration. Detects a band of approximately 83, 67 kDa (predicted molecular weight: 83 kDa).
IP		Use at an assay dependent concentration.

## **Target**

**Function**Catalyzes the oxidation of sulfhydryl groups in peptide and protein thiols to disulfides with the reduction of oxygen to hydrogen peroxide. May contribute to disulfide bond formation in a variety

of secreted proteins. In fibroblasts, it may have tumor-suppressing capabilities being involved in

growth regulation.

**Tissue specificity** Expressed in heart, placenta, lung, liver, skeletal muscle, pancreas and very weakly in brain and

kidney.

Sequence similarities Belongs to the quiescin-sulfhydryl oxidase (QSOX) family.


Contains 1 ERV/ALR sulfhydryl oxidase domain.

Contains 1 thioredoxin domain.


**Cellular localization** Golgi apparatus membrane and Secreted > extracellular space. Found in the extracellular

medium of guiescent cells but is not found in proliferating cells.

#### **Images**



Immunoprecipitation - Anti-Quiescin Q6 antibody [EPR21866] - BSA and Azide free (ab237600)



Lane 1: MDA-MB-231 whole cell lysate 10 µg (Input).

Lane 2: ab235444 IP in MDA-MD-231 whole cell lysate.


 $\label{lambda} \mbox{Lane 3: Rabbit monoclonal lgG } (\mbox{$ab172730$}) \mbox{ instead of } \mbox{$ab235444$} \mbox{ in MDA-MD-231 whole cell lysate.}$ 

**Blocking/Dilution buffer:** 5% NFDM/TBST.

Exposure time: 30 seconds.


Quiescin Q6 has two isoforms due to expression from a long (Quiescin Q6-L) and a short (Quiescin Q6-S) transcript.

This data was developed using the same antibody clone in a different buffer formulation containing PBS, BSA, glycerol, and sodium azide (ab235444).



Flow Cytometry (Intracellular) - Anti-Quiescin Q6 antibody [EPR21866] - BSA and Azide free (ab237600) Intracellular flow cytometric analysis of 4% paraformaldehyde-fixed, 90% methanol-permeabilized MDA-MB-231 (Human breast adenocarcinoma cell line) cells labeling Quiescin Q6 with <a href="mailto:ab235444">ab235444</a> at 1/60 (red) compared with a Rabbit monoclonal IgG (<a href="mailto:ab172730">ab172730</a>) (black) and an unlabeled control (cells without incubation with primary antibody and secondary antibody) (blue). Goat Anti-Rabbit IgG H&L (Alexa Fluor<sup>®</sup> 488) (<a href="mailto:ab150077">ab150077</a>), at 1/2000 dilution was used as the secondary antibody.

This data was developed using the same antibody clone in a different buffer formulation containing PBS, BSA, glycerol, and sodium azide (ab235444).




Immunohistochemistry (Formalin/PFA-fixed paraffinembedded sections) - Anti-Quiescin Q6 antibody
[EPR21866] - BSA and Azide free (ab237600)

Immunohistochemical analysis of paraffin-embedded human breast ductal cancer tissue labeling Quiescin Q6 with <u>ab235444</u> at 1/200 dilution, followed by Goat Anti-Rabbit IgG H&L (HRP) ready to use. Cytoplasmic staining in human breast ductal cancer (PMID: 23536962) is observed. Counter stained with hematoxylin.

Secondary antibody only control: Used PBS instead of primary antibody, secondary antibody is Goat Anti-Rabbit lgG H&L (HRP) ready to use.

Heat-mediated antigen retrieval using <u>ab93684</u> (Tris/EDTA buffer, pH 9.0).

This data was developed using the same antibody clone in a different buffer formulation containing PBS, BSA, glycerol, and sodium azide (ab235444).




Immunohistochemistry (Formalin/PFA-fixed paraffinembedded sections) - Anti-Quiescin Q6 antibody
[EPR21866] - BSA and Azide free (ab237600)

Immunohistochemical analysis of paraffin-embedded human pancreatic cancer tissue labeling Quiescin Q6 with **ab235444** at 1/200 dilution, followed by Goat Anti-Rabbit IgG H&L (HRP) ready to use. Cytoplasmic staining in human pancreatic cancer (PMID: 19795908, 21989104) is observed. Counter stained with hematoxylin.

Secondary antibody only control: Used PBS instead of primary antibody, secondary antibody is Goat Anti-Rabbit lgG H&L (HRP) ready to use.

Heat-mediated antigen retrieval using <u>ab93684</u> (Tris/EDTA buffer, pH 9.0).

This data was developed using the same antibody clone in a different buffer formulation containing PBS, BSA, glycerol, and sodium azide (ab235444).



Please note: All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

#### Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit **https://www.abcam.com/abpromise** or contact our technical team.

#### Terms and conditions

• Guarantee only valid for products bought direct from Abcam or one of our authorized distributors