# abcam

## Product datasheet

# Anti-RHAG antibody [EPR10011] - BSA and Azide free ab249187



#### 7 Images

#### Overview

**Product name** Anti-RHAG antibody [EPR10011] - BSA and Azide free

**Description** Rabbit monoclonal [EPR10011] to RHAG - BSA and Azide free

**Host species** Rabbit

**Tested applications** Suitable for: IHC-P, WB

Unsuitable for: ICC/IF or IP

Species reactivity Reacts with: Human

Predicted to work with: Mouse, Rat

**Immunogen** Synthetic peptide. This information is proprietary to Abcam and/or its suppliers.

**General notes** ab249187 is the carrier-free version of ab155094.

> Our carrier-free antibodies are typically supplied in a PBS-only formulation, purified and free of BSA, sodium azide and glycerol. The carrier-free buffer and high concentration allow for increased conjugation efficiency.

This conjugation-ready format is designed for use with fluorochromes, metal isotopes, oligonucleotides, and enzymes, which makes them ideal for antibody labelling, functional and cellbased assays, flow-based assays (e.g. mass cytometry) and Multiplex Imaging applications.

Use our conjugation kits for antibody conjugates that are ready-to-use in as little as 20 minutes with <1 minute hands-on-time and 100% antibody recovery: available for fluorescent dyes, HRP, biotin and gold.

This product is compatible with the Maxpar<sup>®</sup> Antibody Labeling Kit from Fluidigm, without the need for antibody preparation. Maxpar<sup>®</sup> is a trademark of Fluidigm Canada Inc.

This product is a recombinant monoclonal antibody, which offers several advantages including:

- High batch-to-batch consistency and reproducibility
- Improved sensitivity and specificity
- Long-term security of supply
- Animal-free production

For more information see here.

Our RabMAb® technology is a patented hybridoma-based technology for making rabbit monoclonal antibodies. For details on our patents, please refer to **RabMAb**® **patents**.

1

#### **Properties**

Form Liquid

**Storage instructions** Shipped at 4°C. Store at +4°C. Do Not Freeze.

Storage buffer pH: 7.2

Constituent: PBS

Carrier free Yes

Purity Protein A purified

Clonality Monoclonal
Clone number EPR10011

**Isotype** IgG

#### **Applications**

The Abpromise guarantee Our Abpromise guarantee covers the use of ab249187 in the following tested applications.

The application notes include recommended starting dilutions; optimal dilutions/concentrations should be determined by the end user.

Application	Abreviews	Notes
IHC-P		Use at an assay dependent concentration. Perform heat mediated antigen retrieval before commencing with IHC staining protocol.
WB		Use at an assay dependent concentration. Predicted molecular weight: 44 kDa.


**Application notes** Is unsuitable for ICC/IF or IP.

## **Target**

RHAG (Rhesus blood group-associated glycoprotein) is associated with rhesus blood group antigen expression. It may be part of an oligomeric complex which is likely to have a transport or channel function in the erythrocyte membrane. The Rh blood group antigens are associated with human erythrocyte membrane proteins of approximately 30 kD, the so-called Rh30 polypeptides. Heterogeneously glycosylated membrane proteins of 50 and 45 kD, the Rh50 glycoproteins, are coprecipitated with the Rh30 polypeptides on immunoprecipitation with anti-Rh-specific monoand polyclonal antibodies. The Rh antigens appear to exist as a multisubunit complex of CD47, LW, glycophorin B and play a critical role in the Rh50 glycoprotein.

Cellular localization Cell Membrane; multi-pass membrane protein


#### **Images**



Immunohistochemistry (Formalin/PFA-fixed paraffinembedded sections) - Anti-RHAG antibody
[EPR10011] - BSA and Azide free (ab249187)


This data was developed using <u>ab155094</u>, the same antibody clone in a different buffer formulation.

<u>ab155094</u> showing +ve staining in Human normal tonsil. Perform heat mediated antigen retrieval before commencing with IHC staining protocol.




Immunohistochemistry (Formalin/PFA-fixed paraffinembedded sections) - Anti-RHAG antibody
[EPR10011] - BSA and Azide free (ab249187)

This data was developed using <u>ab155094</u>, the same antibody clone in a different buffer formulation.<u>ab155094</u> showing +ve staining in Human normal heart. Perform heat mediated antigen retrieval before commencing with IHC staining protocol.



Immunohistochemistry (Formalin/PFA-fixed paraffinembedded sections) - Anti-RHAG antibody
[EPR10011] - BSA and Azide free (ab249187)

This data was developed using <u>ab155094</u>, the same antibody clone in a different buffer formulation.<u>ab155094</u> showing +ve staining in Human normal colon. Perform heat mediated antigen retrieval before commencing with IHC staining protocol.



Western blot - Anti-RHAG antibody [EPR10011] - BSA and Azide free (ab249187)

**All lanes :** Anti-RHAG antibody [EPR10011] (<u>ab155094</u>) at 1/1000 dilution

Lane 1 : Human fetal kidney cell lysate

Lane 2 : Human fetal liver cell lysate

Lane 3: Human bone marrow cell lysate


Lysates/proteins at 10 µg per lane.

#### Secondary

All lanes: Goat anti-rabbit HRP at 1/2000 dilution


Predicted band size: 44 kDa

This data was developed using <u>ab155094</u>, the same antibody clone in a different buffer formulation.




Immunohistochemistry (Formalin/PFA-fixed paraffinembedded sections) - Anti-RHAG antibody [EPR10011] - BSA and Azide free (ab249187)

This data was developed using ab155094, the same antibody clone in a different buffer formulation.lmmunohistochemical analysis of paraffin-embedded Human lung tissue labeling RHAG with ab155094 at 1/250 dilution. Perform heat mediated antigen retrieval before commencing with IHC staining protocol.



Immunohistochemistry (Formalin/PFA-fixed paraffinembedded sections) - Anti-RHAG antibody [EPR10011] - BSA and Azide free (ab249187)

This data was developed using ab155094, the same antibody clone in a different buffer formulation.lmmunohistochemical analysis of paraffin-embedded Human spleen tissue labeling RHAG with ab155094 at 1/250 dilution. Perform heat mediated antigen retrieval before commencing with IHC staining protocol.





Research with confidence Consistent and reproducible results



Long-term and scalable supply Recombinant technology



specificity





compliant Animal-free production

Anti-RHAG antibody [EPR10011] - BSA and Azide free (ab249187)

Please note: All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <a href="https://www.abcam.com/abpromise">https://www.abcam.com/abpromise</a> or contact our technical team.

#### Terms and conditions

• Guarantee only valid for products bought direct from Abcam or one of our authorized distributors