# abcam

# Product datasheet

# Anti-SCML4 antibody ab154011

# 3 Images

#### Overview

Product name Anti-SCML4 antibody

**Description** Rabbit polyclonal to SCML4

Host species Rabbit

Tested applications Suitable for: WB, IHC-P

Species reactivity Reacts with: Mouse, Human

Immunogen Recombinant fragment, corresponding to a region within amino acids 1-237 of Human SCML4

(UniProt ID: Q8N228).

**Positive control** SK N SH whole cell lysate; BT483 xenograft tissue; C2C12 xenograft tissue.

**General notes**The Life Science industry has been in the grips of a reproducibility crisis for a number of years.

Abcam is leading the way in addressing this with our range of recombinant monoclonal antibodies and knockout edited cell lines for gold-standard validation. Please check that this product meets

your needs before purchasing.

If you have any questions, special requirements or concerns, please send us an inquiry and/or

contact our Support team ahead of purchase. Recommended alternatives for this product can be

found below, along with publications, customer reviews and Q&As

**Properties** 

Form Liquid

**Storage instructions** Shipped at 4°C. Upon delivery aliquot. Store at -20°C or -80°C. Avoid freeze / thaw cycle.

**Storage buffer** pH: 7.00

Preservative: 0.01% Thimerosal (merthiolate)

Constituents: 78.99% PBS, 1% BSA, 20% Glycerol (glycerin, glycerine)

Purity Immunogen affinity purified

**Clonality** Polyclonal

**Isotype** IgG

**Applications** 

The Abpromise guarantee Our Abpromise guarantee covers the use of ab154011 in the following tested applications.

1

The application notes include recommended starting dilutions; optimal dilutions/concentrations should be determined by the end user.

Application	Abreviews	Notes
WB		1/500 - 1/3000. Predicted molecular weight: 45 kDa.
IHC-P		1/100 - 1/1000. Perform heat mediated antigen retrieval with citrate buffer pH 6 before commencing with IHC staining protocol.

# **Target**


Relevance SCML4 is a putative Polycomb group (PcG) protein. PcG proteins act by forming multiprotein

complexes, which are required to maintain the transcriptionally repressive state of homeotic

genes throughout development.

Cellular localization Nuclear


# **Images**



Anti-SCML4 antibody (ab154011) at 1/1000 dilution + SK-N-SH whole cell lysate at 30  $\mu g$ 


Predicted band size: 45 kDa

12% SDS PAGE



Immunohistochemistry (Formalin/PFA-fixed paraffinembedded sections) - Anti-SCML4 antibody (ab154011)

Immunohistochemical analysis of paraffin embedded BT483 xenograft tissue labeling SCML4 with ab154011 at 1/750 dilution.



Immunohistochemistry (Formalin/PFA-fixed paraffinembedded sections) - Anti-SCML4 antibody (ab154011)

Immunohistochemical analysis of paraffin embedded C2C12 xenograft tissue, labeling SCML4 with ab154011 at 1/750 dilution.

Please note: All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

### Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise,

please visit <a href="https://www.abcam.com/abpromise">https://www.abcam.com/abpromise</a> or contact our technical team.

# Terms and conditions

• Guarantee only valid for products bought direct from Abcam or one of our authorized distributors