

# Anti-SESN2/Sestrin-2 antibody [EPR18907] - BSA and Azide free ab236025

KO VALIDATED

Recombinant

RabMAb

[1 References](#) [8 Images](#)

## Overview

<b>Product name</b>	Anti-SESN2/Sestrin-2 antibody [EPR18907] - BSA and Azide free
<b>Description</b>	Rabbit monoclonal [EPR18907] to SESN2/Sestrin-2 - BSA and Azide free
<b>Host species</b>	Rabbit
<b>Tested applications</b>	<b>Suitable for:</b> Flow Cyt (Intra), IP, ICC/IF, WB
<b>Species reactivity</b>	<b>Reacts with:</b> Mouse, Rat, Human
<b>Immunogen</b>	Recombinant fragment. This information is proprietary to Abcam and/or its suppliers.
<b>Positive control</b>	ICC/IF: HCT 116 cells. WB: HeLa, LoVo and HEK-293T, HEK-293 cell lysate. Flow Cyt (intra): NIH/3T3 and HCT 116 cells. IP: HeLa cell lysate.
<b>General notes</b>	<p>ab236025 is the carrier-free version of <a href="#">ab178518</a>.</p> <p>Our <b>carrier-free</b> antibodies are typically supplied in a PBS-only formulation, purified and free of BSA, sodium azide and glycerol. The carrier-free buffer and high concentration allow for increased conjugation efficiency.</p> <p>This conjugation-ready format is designed for use with fluorochromes, metal isotopes, oligonucleotides, and enzymes, which makes them ideal for antibody labelling, functional and cell-based assays, flow-based assays (e.g. mass cytometry) and Multiplex Imaging applications.</p> <p>Use our <b>conjugation kits</b> for antibody conjugates that are ready-to-use in as little as 20 minutes with &lt;1 minute hands-on-time and 100% antibody recovery: available for fluorescent dyes, HRP, biotin and gold.</p> <p>This product is compatible with the Maxpar<sup>®</sup> Antibody Labeling Kit from Fluidigm, without the need for antibody preparation. Maxpar<sup>®</sup> is a trademark of Fluidigm Canada Inc.</p> <p>This product is a recombinant monoclonal antibody, which offers several advantages including:</p> <ul style="list-style-type: none"> <li>- High batch-to-batch consistency and reproducibility</li> <li>- Improved sensitivity and specificity</li> <li>- Long-term security of supply</li> <li>- Animal-free production</li> </ul> <p>For more information <a href="#">see here</a>.</p> <p>Our RabMAb<sup>®</sup> technology is a patented hybridoma-based technology for making rabbit monoclonal antibodies. For details on our patents, please refer to <a href="#">RabMAb<sup>®</sup> patents</a>.</p>

## Properties

Form	Liquid
Storage instructions	Shipped at 4°C. Store at +4°C. Do Not Freeze.
Storage buffer	pH: 7.2 Constituent: PBS
Carrier free	Yes
Purity	Protein A purified
Clonality	Monoclonal
Clone number	EPR18907
Isotype	IgG


## Applications

**The Abpromise guarantee** Our **Abpromise guarantee** covers the use of ab236025 in the following tested applications. The application notes include recommended starting dilutions; optimal dilutions/concentrations should be determined by the end user.

Application	Abreviews	Notes
Flow Cyt (Intra)		Use at an assay dependent concentration.
IP		Use at an assay dependent concentration.
ICC/IF		Use at an assay dependent concentration. ICC/IF is recommended for human and rat only.
WB		Use at an assay dependent concentration. Detects a band of approximately 54 kDa (predicted molecular weight: 54 kDa).

## Target

## Images



Western blot - Anti-SESN2/Sestrin-2 antibody [EPR18907] - BSA and Azide free (ab236025)

**All lanes :** Anti-SESN2/Sestrin-2 antibody [EPR18907] ([ab178518](#)) at 1/1000 dilution

**Lane 1 :** Wild-type HEK-293 cell lysate


**Lane 2 :** SESN2 knockout HEK-293 cell lysate

Performed under reducing conditions.

**Predicted band size:** 54 kDa

**Observed band size:** 54 kDa

False colour image of Western blot: Anti-SESN2/Sestrin-2 antibody [EPR18907] staining at 1/1000 dilution, shown in green; Mouse anti-GAPDH antibody [6C5] ([ab8245](#)) loading control staining at 1/20000 dilution, shown in red. In Western blot, [ab178518](#) was shown to bind specifically to SESN2/Sestrin-2. A band was observed at 54 kDa in wild-type HEK-293 cell lysates with no signal observed at this size in SESN2 knockout cell line [ab269486](#) (knockout cell lysate [ab269650](#)). To generate this image, wild-type and SESN2 knockout HEK-293 cell lysates were analysed. First, samples were run on an SDS-PAGE gel then transferred onto a nitrocellulose membrane. Membranes were blocked in 3 % milk in TBS-0.1 % Tween<sup>®</sup> 20 (TBS-T) before incubation with primary antibodies overnight at 4 °C. Blots were washed four times in TBS-T, incubated with secondary antibodies for 1 h at room temperature, washed again four times then imaged. Secondary antibodies used were Goat anti-Rabbit IgG H&L (IRDye<sup>®</sup> 800CW) preabsorbed ([ab216773](#)) and Goat anti-Mouse IgG H&L (IRDye<sup>®</sup> 680RD) preabsorbed ([ab216776](#)) at 1/20000 dilution.



Western blot - Anti-SESN2/Sestrin-2 antibody [EPR18907] - BSA and Azide free (ab236025)

**All lanes :** Anti-SESN2/Sestrin-2 antibody [EPR18907] ([ab178518](#)) at 1/1000 dilution

**Lane 1 :** Wild-type HeLa cell lysate

**Lane 2 :** SESN2 knockout HeLa cell lysate

**Lane 3 :** LoVo cell lysate

**Lane 4 :** HEK-293T cell lysate

Lysates/proteins at 20 µg per lane.

### Secondary

**All lanes :** Goat anti-Rabbit IgG H&L (IRDye® 800CW) preadsorbed ([ab216773](#)) at 1/10000 dilution

**Predicted band size:** 54 kDa


**Observed band size:** 54 kDa

This data was developed using the same antibody clone in a different buffer formulation ([ab178518](#)).

**Lanes 1-4:** Merged signal (red and green). Green - [ab178518](#) observed at 54 kDa. Red - loading control [ab8245](#) observed at 36 kDa.

[ab178518](#) Anti-SESN2/Sestrin-2 antibody [EPR18907] was shown to specifically react with 2 in wild-type HeLa cells. Loss of signal was observed when knockout sample [ab257665](#) was used. Wild-type and 2 knockout samples were subjected to SDS-PAGE.

[ab178518](#) and Anti-GAPDH antibody [6C5] - Loading Control ([ab8245](#)) were incubated at room temperature for 2.5 hours at 1 in 1000 dilution and 1 in 20000 dilution respectively. Blots were developed with Goat anti-Rabbit IgG H&L (IRDye® 800CW) preadsorbed ([ab216773](#)) and Goat anti-Mouse IgG H&L (IRDye® 680RD) preadsorbed ([ab216776](#)) secondary antibodies at 1 in 20000 dilution for 1 hour at room temperature before imaging.




Immunocytochemistry/ Immunofluorescence - Anti-  
SESN2/Sestrin-2 antibody [EPR18907] - BSA and  
Azide free (ab236025)

Immunofluorescent analysis of 4% paraformaldehyde-fixed, 0.1% Triton X-100 permeabilized HCT 116 (Human colorectal carcinoma cell line) cells labeling SESN2/Sestrin-2 with **ab178518** at 1/100 dilution, followed by Goat anti-rabbit IgG (Alexa Fluor® 488) (**ab150077**) secondary antibody at 1/1000 dilution (green). Confocal image showing cytoplasmic staining on HCT116 cells. The nuclear counter stain is DAPI (blue). Tubulin is detected with **ab195889** Anti-alpha Tubulin antibody [DM1A] - Microtubule Marker (Alexa Fluor® 594) at 1/250 dilution (red).

Secondary antibody only control: Used PBS instead of primary antibody, secondary antibody is Goat anti-rabbit IgG (Alexa Fluor® 488) (**ab150077**) secondary antibody at 1/1000 dilution.

This data was developed using the same antibody clone in a different buffer formulation containing PBS, BSA, glycerol, and sodium azide (**ab178518**).



Western blot - Anti-SESN2/Sestrin-2 antibody  
[EPR18907] - BSA and Azide free (ab236025)

**All lanes** : Anti-SESN2/Sestrin-2 antibody [EPR18907]  
(**ab178518**) at 1/1000 dilution

**Lane 1** : Wild-type HeLa cell lysate

**Lane 2** : SESN2 knockout HeLa cell lysate

**Lane 3** : HeLa cell lysate

Lysates/proteins at 20 µg per lane.

### Secondary

**All lanes** : Goat anti-Rabbit IgG H&L (IRDye® 800CW)  
preadsorbed (**ab216773**) at 1/10000 dilution


**Predicted band size:** 54 kDa

**Observed band size:** 54 kDa

This data was developed using the same antibody clone in a different buffer formulation (**ab178518**).

**Lanes 1-3:** Merged signal (red and green). Green - **ab178518** observed at 54 kDa. Red - loading control **ab8245** observed at 36 kDa.

**ab178518** Anti-SESN2/Sestrin-2 antibody [EPR18907] was shown to specifically react with SESN2/Sestrin-2 in wild-type HeLa cells. Loss of signal was observed when knockout cell line **ab265669** (knockout cell lysate **ab257665**) was used. Wild-type and SESN2/Sestrin-2 knockout samples were subjected to SDS-PAGE. **ab178518** and Anti-GAPDH antibody [6C5] - Loading Control (**ab8245**) were incubated overnight at 4°C at 1 in 1000 dilution and 1 in 20000 dilution respectively. Blots were developed with Goat anti-Rabbit IgG H&L (IRDye® 800CW) preadsorbed (**ab216773**) and Goat anti-Mouse IgG H&L (IRDye® 680RD) preadsorbed (**ab216776**) secondary antibodies at 1 in 20000 dilution for 1 hour at room temperature before imaging.



SESN2/Sestrin-2 was immunoprecipitated from 0.35 mg of HeLa (Human epithelial cell line from cervix adenocarcinoma) whole cell lysate with **ab178518** at 1/30 dilution. Western blot was performed from the immunoprecipitate using **ab178518** at 1/1000 dilution. VeriBlot for IP Detection Reagent (HRP) (**ab131366**), was used for detection at 1/10000 dilution.

Lane 1: HeLa whole cell lysate, 10µg (Input).

Lane 2: **ab178518** IP in HeLa whole cell lysate.


Lane 3: Rabbit IgG, monoclonal [EPR25A] - Isotype Control (**ab172730**) instead of **ab178518** in HeLa whole cell lysate.

Blocking and dilution buffer and concentration: 5% NFDm/TBST.

Exposure time: 3 minutes.

SESN2/Sestrin-2 expression is low in HeLa cells and can be enriched through immunoprecipitation.


This data was developed using the same antibody clone in a different buffer formulation containing PBS, BSA, glycerol, and sodium azide (**ab178518**).



Flow Cytometry (Intracellular) - Anti-SESN2/Sestrin-2 antibody [EPR18907] - BSA and Azide free (ab236025)

Intracellular flow cytometric analysis of 4% paraformaldehyde-fixed NIH/3T3 (Mouse embryonic fibroblast cell line) cells labeling SESN2/Sestrin-2 with **ab178518** at 1/60 dilution (red) compared with a Rabbit IgG,monoclonal[EPR25A]-Isotype control (**ab172730**) (black) and an unlabelled control (cells without incubation with primary antibody and secondary antibody) (blue). Goat anti Rabbit IgG (Alexa Fluor® 488) at 1/2000 dilution was used as the secondary antibody.

This data was developed using the same antibody clone in a different buffer formulation containing PBS, BSA, glycerol, and sodium azide (**ab178518**).



Flow Cytometry (Intracellular) - Anti-SESN2/Sestrin-2 antibody [EPR18907] - BSA and Azide free (ab236025)

Intracellular flow cytometric analysis of 4% paraformaldehyde-fixed HCT 116 (Human colorectal carcinoma cell line) cells labeling SESN2/Sestrin-2 with **ab178518** at 1/60 dilution (red) compared with a rabbit monoclonal IgG isotype control (**ab172730**; black) and an unlabelled control (cells without incubation with primary antibody and secondary antibody; blue). Goat anti rabbit IgG (Alexa Fluor® 488) at 1/2000 dilution was used as the secondary antibody.

This data was developed using the same antibody clone in a different buffer formulation containing PBS, BSA, glycerol, and sodium azide (**ab178518**).

### Why choose a recombinant antibody?



**Research with confidence**  
Consistent and reproducible results



**Long-term and scalable supply**  
Recombinant technology



**Success from the first experiment**  
Confirmed specificity



**Ethical standards compliant**  
Animal-free production

Anti-SESN2/Sestrin-2 antibody [EPR18907] - BSA  
and Azide free (ab236025)

**Please note:** All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

### Our Abpromise to you: Quality guaranteed and expert technical support

---

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <https://www.abcam.com/abpromise> or contact our technical team.

### Terms and conditions

---

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors