

Product datasheet

Anti-SPINK1/P12 antibody [EPR17585-30] ab206294

Recombinant RabMAb

[4 Images](#)

Overview

Product name	Anti-SPINK1/P12 antibody [EPR17585-30]
Description	Rabbit monoclonal [EPR17585-30] to SPINK1/P12
Host species	Rabbit
Tested applications	Suitable for: IHC-P, WB
Species reactivity	Reacts with: Human
Immunogen	Synthetic peptide. This information is proprietary to Abcam and/or its suppliers.
Positive control	WB; Human pancreas lysate. IHC-P; Human pancreatic carcinoma tissue, Human liver carcinoma tissue.
General notes	<p>This product is a recombinant monoclonal antibody, which offers several advantages including:</p> <ul style="list-style-type: none">- High batch-to-batch consistency and reproducibility- Improved sensitivity and specificity- Long-term security of supply- Animal-free production <p>For more information see here.</p> <p>Our RabMAb[®] technology is a patented hybridoma-based technology for making rabbit monoclonal antibodies. For details on our patents, please refer to RabMAb[®] patents.</p>

Properties

Form	Liquid
Storage instructions	Shipped at 4°C. Store at +4°C short term (1-2 weeks). Upon delivery aliquot. Store at -20°C long term. Avoid freeze / thaw cycle.
Storage buffer	pH: 7.2 Preservative: 0.01% Sodium azide Constituents: 59% PBS, 40% Glycerol (glycerin, glycerine), 0.05% BSA
Purity	Protein A purified
Clonality	Monoclonal
Clone number	EPR17585-30
Isotype	IgG

Applications


The Abpromise guarantee Our **Abpromise guarantee** covers the use of ab206294 in the following tested applications. The application notes include recommended starting dilutions; optimal dilutions/concentrations should be determined by the end user.

Application	Abreviews	Notes
IHC-P		1/100. Perform heat mediated antigen retrieval with Tris/EDTA buffer pH 9.0 before commencing with IHC staining protocol.
WB		1/5000. Detects a band of approximately 11 kDa (predicted molecular weight: 8 kDa).

Target

Function	Serine protease inhibitor which exhibits anti-trypsin activity (PubMed:7142173). In the pancreas, protects against trypsin-catalyzed premature activation of zymogens. In the male reproductive tract, binds to sperm heads where it modulates sperm capacitance by inhibiting calcium uptake and nitrogen oxide (NO) production.
Involvement in disease	Pancreatitis, hereditary Tropical calcific pancreatitis
Sequence similarities	Contains 1 Kazal-like domain.
Cellular localization	Secreted.

Images




Immunohistochemical analysis of paraffin-embedded human pancreatic carcinoma tissue labeling SPINK1/P12 with ab206294 at 1/100 dilution followed by Goat Anti-Rabbit IgG H&L (HRP) ([ab97051](#)) at 1/500 dilution.

Cytoplasmic staining on human pancreatic carcinoma tissue is observed.

Counter stained with Hematoxylin.

Negative control: Used PBS instead of primary ab, secondary ab is Goat Anti-Rabbit IgG H&L (HRP) ([ab97051](#)).

Perform heat mediated antigen retrieval with Tris/EDTA buffer pH 9.0 before commencing with IHC staining protocol.



Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-SPINK1/P12 antibody [EPR17585-30] (ab206294)


Immunohistochemical analysis of paraffin-embedded human liver carcinoma tissue labeling SPINK1/P12 with ab206294 at 1/100 dilution followed by Goat Anti-Rabbit IgG H&L (HRP) ([ab97051](#)) at 1/500 dilution.

Cytoplasmic staining on human liver carcinoma tissue is observed.

Counter stained with Hematoxylin.

Negative control: Used PBS instead of primary ab, secondary ab is Goat Anti-Rabbit IgG H&L (HRP) ([ab97051](#)).

Perform heat mediated antigen retrieval with Tris/EDTA buffer pH 9.0 before commencing with IHC staining protocol.



Western blot - Anti-SPINK1/P12 antibody [EPR17585-30] (ab206294)

All lanes : Anti-SPINK1/P12 antibody [EPR17585-30] (ab206294) at 1/5000 dilution

Lane 1 : Human pancreas lysate

Lane 2 : Human fetal kidney tissue lysate

Lane 3 : HepG2 (Human liver hepatocellular carcinoma cell line) whole cell lysate

Lane 4 : BxPC-3 (Human pancreas adenocarcinoma cell line) whole cell lysate

Lysates/proteins at 20 µg per lane.

Secondary

All lanes : Anti-Rabbit IgG (HRP), specific to the non-reduced form of IgG at 1/10000 dilution

Predicted band size: 8 kDa

Observed band size: 11 kDa

Blocking/dilution buffer: 5% NFDM/TBST.

Exposure time; Lane 1:5 seconds;

Lane 2,3 and 4: 3 minutes.

Why choose a recombinant antibody?



Research with confidence
Consistent and reproducible results



Long-term and scalable supply
Recombinant technology



Success from the first experiment
Confirmed specificity



Ethical standards compliant
Animal-free production

Anti-SPINK1/P12 antibody [EPR17585-30]
(ab206294)

Please note: All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <https://www.abcam.com/abpromise> or contact our technical team.

Terms and conditions

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors