

Anti-Steroidogenic Factor 1/SF-1 antibody [EPR19744] ab217317

Recombinant RabMAb

[7 References](#) [4 Images](#)

Overview

Product name	Anti-Steroidogenic Factor 1/SF-1 antibody [EPR19744]
Description	Rabbit monoclonal [EPR19744] to Steroidogenic Factor 1/SF-1
Host species	Rabbit
Tested applications	Suitable for: IHC-P
Species reactivity	Reacts with: Human
Immunogen	Synthetic peptide. This information is proprietary to Abcam and/or its suppliers.
Positive control	IHC-P: Human ovary, testis and adrenal cancer tissues.
General notes	<p>This product is a recombinant monoclonal antibody, which offers several advantages including:</p> <ul style="list-style-type: none"> - High batch-to-batch consistency and reproducibility - Improved sensitivity and specificity - Long-term security of supply - Animal-free production <p>For more information see here.</p> <p>Our RabMAb[®] technology is a patented hybridoma-based technology for making rabbit monoclonal antibodies. For details on our patents, please refer to RabMAb[®] patents.</p>

Properties

Form	Liquid
Storage instructions	Shipped at 4°C. Store at +4°C short term (1-2 weeks). Upon delivery aliquot. Store at -20°C long term. Avoid freeze / thaw cycle.
Storage buffer	<p>pH: 7.2</p> <p>Preservative: 0.01% Sodium azide</p> <p>Constituents: 59% PBS, 40% Glycerol (glycerin, glycerine), 0.05% BSA</p>
Purity	Protein A purified
Clonality	Monoclonal
Clone number	EPR19744
Isotype	IgG

Applications

The Abpromise guarantee

Our **Abpromise guarantee** covers the use of ab217317 in the following tested applications.

The application notes include recommended starting dilutions; optimal dilutions/concentrations should be determined by the end user.

Application	Abreviews	Notes
IHC-P		1/2000. Perform heat mediated antigen retrieval with Tris/EDTA buffer pH 9.0 before commencing with IHC staining protocol.

Target

Function

Transcriptional activator. Seems to be essential for sexual differentiation and formation of the primary steroidogenic tissues. Binds to the Ad4 site found in the promoter region of steroidogenic P450 genes such as CYP11A, CYP11B and CYP21B. Also regulates the AMH/Muellerian inhibiting substance gene as well as the AHCH and STAR genes. 5'-YCAAGGYC-3' and 5'-RRAGGTCA-3' are the consensus sequences for the recognition by NR5A1. The SFPQ-NONO-NR5A1 complex binds to the CYP17 promoter and regulates basal and cAMP-dependent transcriptional activity. Binds phosphatidylcholine (By similarity). Binds phospholipids with a phosphatidylinositol (PI) headgroup, in particular PI(3,4)P2 and PI(3,4,5)P3.

Involvement in disease

Defects in NR5A1 are a cause of 46,XY disorder of sex development (46,XY DSD) [MIM:612965]; also known as XY sex reversal with or without adrenal failure. A congenital condition in which development of chromosomal, gonadal, or anatomic sex is atypical. 46,XY DSD is a disorder of gonadal (testicular) development, which may be complete or partial. The complete form includes streak gonads, normal mullerian structures, and normal female external genitalia. The partial form includes ambiguous external genitalia and partial development of mullerian and wolffian structures.

Defects in NR5A1 are a cause of adrenocortical insufficiency without ovarian defect (ACIWOD) [MIM:184757]. ACIWOD is characterized by severe 'slackness' muscular hypotonia. There is decreased sodium, increased potassium and elevated ACTH.

Defects in NR5A1 are the cause of premature ovarian failure type 7 (POF7) [MIM:612964]. An ovarian disorder defined as the cessation of ovarian function under the age of 40 years. It is characterized by oligomenorrhea or amenorrhea, in the presence of elevated levels of serum gonadotropins and low estradiol.

Sequence similarities

Belongs to the nuclear hormone receptor family. NR5 subfamily.
Contains 1 nuclear receptor DNA-binding domain.


Post-translational modifications

Acetylation stimulates the transcriptional activity.

Cellular localization

Nucleus.

Images




Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-Steroidogenic Factor 1/SF-1 antibody [EPR19744] (ab217317)

Immunohistochemical analysis of paraffin-embedded human ovary tissue labeling Steroidogenic Factor 1/SF-1 with ab217317 at 1/2000 dilution, followed by Goat Anti-Rabbit IgG H&L (HRP) Ready to use. Nuclear staining on human ovary is observed [PMID: 20660055] [PMID: 10370224]. Counter stained with Hematoxylin.

Secondary antibody only control: Used PBS instead of primary antibody, secondary antibody is Goat Anti-Rabbit IgG H&L (HRP) Ready to use.

Perform heat mediated antigen retrieval with Tris/EDTA buffer pH 9.0 before commencing with IHC staining protocol.




Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-Steroidogenic Factor 1/SF-1 antibody [EPR19744] (ab217317)

Immunohistochemical analysis of paraffin-embedded human testis tissue labeling Steroidogenic Factor 1/SF-1 with ab217317 at 1/2000 dilution, followed by Goat Anti-Rabbit IgG H&L (HRP) Ready to use. Nuclear staining on human testis is observed [PMID: 10370224]. Counter stained with Hematoxylin.

Secondary antibody only control: Used PBS instead of primary antibody, secondary antibody is Goat Anti-Rabbit IgG H&L (HRP) Ready to use.

Perform heat mediated antigen retrieval with Tris/EDTA buffer pH 9.0 before commencing with IHC staining protocol.







Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-Steroidogenic Factor 1/SF-1 antibody [EPR19744] (ab217317)

Immunohistochemical analysis of paraffin-embedded human adrenal cancer tissue labeling Steroidogenic Factor 1/SF-1 with ab217317 at 1/2000 dilution, followed by Goat Anti-Rabbit IgG H&L (HRP) Ready to use. Nuclear staining on human adrenal cancer is observed [PMID: 20660055]. Counter stained with Hematoxylin.

Secondary antibody only control: Used PBS instead of primary antibody, secondary antibody is Goat Anti-Rabbit IgG H&L (HRP) Ready to use.

Perform heat mediated antigen retrieval with Tris/EDTA buffer pH 9.0 before commencing with IHC staining protocol.

Why choose a recombinant antibody?

 <p>Research with confidence Consistent and reproducible results</p>	 <p>Long-term and scalable supply Recombinant technology</p>
 <p>Success from the first experiment Confirmed specificity</p>	 <p>Ethical standards compliant Animal-free production</p>

Anti-Steroidogenic Factor 1/SF-1 antibody
[EPR19744] (ab217317)

Please note: All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

Our Abpromise to you: Quality guaranteed and expert technical support

-
- Replacement or refund for products not performing as stated on the datasheet
 - Valid for 12 months from date of delivery
 - Response to your inquiry within 24 hours

 - We provide support in Chinese, English, French, German, Japanese and Spanish
 - Extensive multi-media technical resources to help you
 - We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <https://www.abcam.com/abpromise> or contact our technical team.

Terms and conditions

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors